Oncogenes and Tumor Suppressor Genes

Eva Y.H.P. Lee¹ and William J. Muller²

¹Department of Biological Chemistry and Department of Developmental and Cell Biology, University of California, Irvine, California 92697-4037

²Goodman Cancer Center, McGill University, Montreal, Quebec H3A 1A3, Canada

Correspondence: william.muller@mcgill.ca

Breast cancer progression involves multiple genetic events, which can activate dominant-acting oncogenes and disrupt the function of specific tumor suppressor genes. This article describes several key oncogene and tumor suppressor signaling networks that have been implicated in breast cancer progression. Among the tumor suppressors, the article emphasizes *BRCA1/2* and *p53* tumor suppressors. In addition to these well characterized tumor suppressors, the article highlights the importance of PTEN tumor suppressor in counteracting PI3K signaling from activated oncogenes such as ErbB2. This article discusses the use of mouse models of human breast that recapitulate the key genetic events involved in the initiation and progression of breast cancer. Finally, the therapeutic potential of targeting these key tumor suppressor and oncogene signaling networks is discussed.

Karyotypic and epidemiological analyses of mammary tumors at various stages suggest that breast carcinomas become increasingly aggressive through the stepwise accumulation of genetic changes. The majority of genetic changes found in human breast cancer fall into two categories: gain-of-function mutations in proto-oncogenes, which stimulate cell growth, division, and survival; and loss-of-function mutations in tumor suppressor genes that normally help prevent unrestrained cellular growth and promote DNA repair and cell cycle checkpoint activation. Epigenetic deregulation also contributes to the abnormal expression of these genes. For example, genes that encode enzymes involved in histone modification are mutated in primary renal cell carcinoma (Dalglish et al. 2010; van Haaften et al. 2009).

In addition, the involvement of noncoding RNAs in tumorigenesis and tumor metastasis has been recently documented (Croce 2009; Shimono et al. 2009). These can act as oncogenes or tumor suppressor genes, depending on the context. Here, we discuss genes that are frequently altered in breast cancer, focusing on *ErbB2*, *PI3K* (phosphatidylinositol 3 kinase) pathways, *TP53*, *BRCA1/2*, and *PTEN* (phosphatase and tensin homolog deleted on chromosome 10). Genetically engineered mouse models are emphasized because these provide a wealth of biological information. We consider in detail genetic and biochemical studies that have shown that oncogenic proteins and tumor suppressors provide a critical balance in regulation of key pathways that control cell number and cell behavior.

Editors: Mina J. Bissell, Kornelia Polyak, and Jeffrey Rosen

Additional Perspectives on The Mammary Gland as an Experimental Model available at www.cshperspectives.org

Copyright © 2010 Cold Spring Harbor Laboratory Press; all rights reserved; doi: 10.1101/cshperspect.a003236

Cite this article as *Cold Spring Harb Perspect Biol* 2010;2:a003236

E.Y.H.P. Lee and W.J. Muller

ONCOGENES AND TUMOR SUPPRESSOR GENES

The identification of oncogenes such as *H-RAS* and tumor suppressor genes such as that encoding retinoblastoma protein (*RB*) involved a combination of functional cloning, linkage analyses, positional cloning, or mutational analyses of genetically predisposed individuals. Comparative genomic hybridization has since revealed various genes that can be amplified or deleted in cancer. The human genome projects further increased the speed of discovery (Bell 2010). Breast cancer genome analyses indicate that there are only a few genes that are frequently mutated but many that are infrequently mutated, providing an explanation for the observed cancer heterogeneity. Complex somatic DNA rearrangements, mostly intrachromosomal tandem duplications, have been found in the breast cancer genomes that have been sequenced. Many of these rearrangements are because of nonhomologous end joining (Stephens et al. 2009). Comparison of genome sequences of primary breast cancer and cancer metastasis show limited de novo mutations in metastasis in addition to significantly shared mutations with the primary tumor (Ding et al. 2010). In a case of metastasis evolved over 9 years, a large number of mutations are novel when compared to the primary tumor (Shah et al. 2009). These studies provide information on tumor evolution and identify pathways critical to breast cancer metastasis.

The *ErbB2*, *PI3KCA*, *MYC*, and *CCND1* (encodes cyclin D1) oncogenes are frequently deregulated in breast cancer. Loss-of-function mutations of *RB* in breast cancer cell lines and primary tumors were reported in 1988 (Lee et al. 1988; T'Ang et al. 1988), but a full appreciation of the tumor suppressor genes affected came much later from studies of hereditary breast cancer. To date, ten tumor suppressor genes, all of which are involved in the regulation of genomic integrity, have been associated with hereditary breast cancer (Walsh and King 2007). *BRCA1* and *BRCA2* mutations are associated with a significantly elevated risk for breast and ovarian cancers. Rare germ-line mutations in

TP53 (the gene encoding p53) and *PTEN* are associated with high risk for various cancers, including breast cancer. Germ-line mutations in *ATM*, *CHK2*, *NBS1*, *RAD50*, *PALB2*, and *BRIP1* all moderately increase breast cancer risk. Among these tumor suppressors, *ATM* and *CHK2* are kinases involved in the DNA damage response. *RAD50* is a component of protein complex critical to DNA double-stranded-break end processing. *PALB2* encodes a *BRCA2*-interacting protein and *BRIP1* encodes a *BRCA1*-interacting protein with DNA helicase activity. These genes are also mutated in pancreatic, prostate, and other tumors. Tumors develop because of the loss-of-heterozygosity mutations in the remaining normal allele plus other somatic mutations. For example, *BRCA1*-driven breast cancers frequently harbor somatic mutations in *TP53* and *PTEN* (Holstege et al. 2009; Saal et al. 2008).

Several pathways are frequently deregulated in breast cancer as a consequence of mutations in these genes. These include pathways involving key lipid and protein kinases that function in cell growth and survival, the cell cycle machinery, DNA damage response pathways, and apoptosis.

THE PHOSPHATIDYLINOSITOL 3 KINASE (PI3K) PATHWAY

PI3K signaling influences cell growth, survival, metabolism, and metastasis. PI3K is activated by growth factors and signals from the extracellular matrix (Dillon et al. 2007b). Alterations in upstream components of the PI3K pathway, such as receptor tyrosine kinases, and downstream components such as AKT are frequent in breast cancers. The *PTEN* tumor suppressor antagonizes the PI3K pathway. Loss of *PTEN* function, and activating mutations in or amplification of the gene that encodes the PIK3 catalytic subunit (*PIK3CA*) are both common.

The Akt (PKB) serine kinase family—AKT1, AKT2, and AKT3—are downstream molecules in the PI3K pathway. These have been implicated in a number of cellular processes, including control of cell proliferation, cell survival, and metabolism (Dillon et al. 2007b). Direct

evidence supporting a role for AKT1 in mammary tumor progression came from studies using transgenic mice that expressed different activated forms of AKT1. Expression of these in the mammary epithelium, although incapable of inducing mammary tumors, resulted in a profound involution defect (Ackler et al. 2002; Dillon et al. 2009; Hutchinson et al. 2004). However, coexpression of an activated AKT1 mutant (AKT1-DD) with an activated ErbB2 mutant (NDL) or a PI3K defective middle T oncogene resulted in a decrease in tumor latency in these tumor models (Dillon et al. 2009; Hutchinson et al. 2004). AKT1 coexpression decreased lung metastases, however, in tumor-bearing animals in the MMTV ErbB2 model (Dillon et al. 2009; Hutchinson et al. 2004). Conversely, germ-line deletion of AKT1 profoundly reduced mammary tumor formation in this ErbB2 mouse model of human breast cancer (Ju et al. 2007; Maroulakou et al. 2007). Germ-line deletion of AKT2 accelerated mammary tumor induction in this ErbB2 tumor models (Maroulakou et al. 2007), whereas expression of activated AKT2 in mammary epithelium had little impact on tumor induction but dramatically increased tumor metastasis (Dillon et al. 2009). The enhanced metastatic phenotypes that have been observed in various AKT models have recently been linked to AKT-mediated suppression of the miR 200 micro-RNA, which suppresses the epithelial–mesenchymal transition in epithelial cells (Iliopoulos et al. 2009). Knocking out another kinase in this pathway—the Snf1-related serine kinase Hunk—abrogates the metastatic phenotype in an ErbB2 mouse tumor model (Wertheim et al. 2009). Oncogenic activation of the different serine kinases may thus have distinct effects on the induction and metastatic phases of mammary tumor development.

Recent studies have also examined the impact of mammary-specific disruption of PTEN. Conditional ablation of PTEN in the mammary epithelium of this MMTV/activated ErbB2 or ErbB2^{KI} strains dramatically accelerated mammary tumor progression (Dourdin et al. 2008; Schade et al. 2009). Gene expression profiling of PTEN-deficient ErbB2 tumors

further showed they had acquired many of the salient transcriptional features of the basal breast cancer subtype (Dourdin et al. 2008). Inactivation of a key tumor suppressor pathway thus appears to impact on the breast cancer subtype.

TRANSGENIC MODELS OF ErbB2-INDUCED TUMOR PROGRESSION

ErbB2 is a member of the EGFR family of receptor tyrosine kinases (RTKs). This family comprises four closely related members: EGFR, ErbB2 (Neu, HER-2), ErbB3 (HER-3), and ErbB4 (HER-4) (Hynes and Stern 1994). The importance of ErbB2 in primary human breast cancer is highlighted by the fact that 20%–30% of human breast cancers show elevated levels of ErbB2 because of genomic amplification of the *ErbB2* proto-oncogene (Slamon 1987, 1989). The overexpression of ErbB2 strongly correlates with a negative clinical prognosis in both lymph-node-positive (Antoniotti et al. 1994; Mansour et al. 1994; Ravdin and Chamness 1995) and node-negative (Andrulis et al. 1998) breast cancer patients. Further evidence that overexpression of ErbB2 results in an aggressive tumor type stems from studies showing that elevated ErbB2 expression occurs in many in situ and invasive human ductal carcinomas but is rarely observed in benign breast disorders, such as hyperplasias and dysplasias (Allred et al. 1992; Mansour et al. 1994). Significantly, ErbB-2 may be useful not only as a prognostic marker but also as a predictive marker, given that elevated expression of HER2 predicts tamoxifen resistance of the primary tumor (Pegram et al. 1998) and the response to anti-HER2 targeted therapy such as Herceptin.

Direct evidence supporting a role for the various EGFR family members and their ligands in mammary tumorigenesis comes from studies of transgenic mice. Expression of EGFR or its ligand TGF α in mammary epithelium in transgenic mice resulted in the frequent induction of mammary adenocarcinomas (Brandt et al. 2000; Matsui et al. 1990; Sandgren et al. 1990, 1995). Mammary-epithelium-specific expression of activated ErbB2 similarly rapidly

E.Y.H.P. Lee and W.J. Muller

induced metastatic multifocal mammary tumors (Bouchard et al. 1989; Guy et al. 1996; Lucchini et al. 1992; Muller et al. 1988; Siegel et al. 1999). Whereas expression of the activated *ErbB2* oncogene in murine mammary epithelium is capable of efficiently inducing multifocal mammary tumors, no comparable activating mutations have been detected in human *ErbB2* (Lemoine et al. 1990). Thus, the primary mechanism by which *ErbB2* induces mammary tumorigenesis in human breast cancer must be through overexpression of the wild-type receptor. To assess whether elevated expression of the wild-type *ErbB2* product can indeed induce mammary tumors, Guy et al. generated transgenic mice carrying the *ErbB2* proto-oncogene under the transcriptional control of the MMTV promoter/enhancer (Guy et al. 1992). In contrast to the rapid tumor progression observed in the activated *ErbB2*-bearing strains, focal mammary tumors arose in these strains after a long latency period (Guy et al. 1992). Tumor progression in these strains is associated with the activation of the *ErbB2* tyrosine kinase by somatic mutations in the transgene in at least 70% of the mammary tumors analyzed (Siegel et al. 1994). Significantly, these mutations are confined to a cysteine-rich region of the receptor in the juxta-transmembrane domain and comprise either deletion or insertion of single cysteine residues (Siegel et al. 1994). Further genetic and biochemical analyses have revealed that these cysteine alterations promote the formation intermolecular cysteine bridges between *ErbB2* monomers, which results in constitutive receptor dimerization and activation (Siegel and Muller 1996).

Genetic and biochemical analyses of these various transgenic mouse models have also revealed potential roles for other members of the EGFR family. For example, coexpression of the EGFR ligand TGF α and the *ErbB2* proto-oncogene resulted in a dramatic acceleration of tumor progression that correlated with the tyrosine phosphorylation of EGFR and *ErbB2* (Muller et al. 1996). In contrast to the parental MMTV/*ErbB2* strains, sporadic mutations were not detected in the *ErbB2* transgene in

these mice, presumably because activated EGFR can transphosphorylate *ErbB2*. Conversely, inhibition of EGFR function can dramatically impair mammary tumor formation in transgenic animals expressing activated *ErbB2* in the mammary epithelium (Gillgrass et al. 2003; Lenferink et al. 2000). Another EGFR family member that has been implicated in *ErbB2*-induced tumor progression is *ErbB3*. Immunoblot analyses revealed that *ErbB3* levels were dramatically elevated during tumor progression in *ErbB2* transgenic strains. In fact, quantitative measurement of *ErbB3* levels revealed a 10- to 20-fold increase in the levels of *ErbB3* protein in tumors compared with adjacent normal tissue (Siegel et al. 1999). Although a dramatic increase in the amount of *ErbB3* protein was observed in these strains, equivalent levels of *ErbB3* transcript were detected in normal and tumor tissues (Siegel et al. 1999). Thus, the increased levels of *ErbB3* observed in the tumors are the result of an increase in either the stability or the translation of *ErbB3*.

Tumor progression in transgenic mice expressing the *ErbB2* proto-oncogene is associated with mutations in the cysteine-rich juxta-transmembrane region (Siegel and Muller 1996; Siegel et al. 1994, 1999). Given the large proportion of the *ErbB2*-induced mammary tumors that possess these activating mutations, there appears to be strong biological selection for them. Although comparable somatic mutations in *ERBB2* have not been detected in human breast cancer, several studies have reported the expression of an alternatively spliced *ErbB2* isoform that carries a 16-residue in-frame deletion in the juxtatransmembrane domain (herein referred to *ErbB2* Δ Ex16) that closely resembles the protein generated by the sporadic mutations in the *ErbB2* transgenic mice (Kwong and Hung 1998; Siegel et al. 1999). This splice variant is constitutively active because of its capacity to form disulfide-bonded dimers (Kwong and Hung 1998; Siegel et al. 1999). Because the *ErbB2* cDNA used in the transgenic mice generates an RNA that is incapable of undergoing alternative splicing, the selection for sporadic mutations in these mice may reflect a need to mimic this alternatively

spliced form. Indeed, recent crystal studies with the ErbB2 extracellular domain suggest that ErbB2 does not readily form homodimers owing to electrostatic repulsion (Garrett et al. 2003). Thus, this splice isoform may be required for the production of ErbB2 homodimers through formation of covalent bonds between cysteine residues. Although these genetic studies suggest that the ErbB2 Δ Ex16 splice variant plays a critical role in ErbB2-driven tumor induction, the biological significance remains to be established.

One limitation of the transgenic mouse models is that expression of ErbB2 is driven by a strong viral promoter. In an attempt to more closely mimic the events involved in ErbB2-induced mammary tumor progression, we have recently derived transgenic mice that carry a Cre-inducible activated ErbB2 under the transcriptional control of the endogenous *ErbB2* promoter (herein referred as the ErbB2^{KI} model) (Andrechek et al. 2000). In contrast to the rapid tumor progression observed in the MMTV-activated ErbB2 strains, focal mammary tumors arose only after an extended latency period. Tumor progression in these strains is associated with a dramatic elevation in the levels of both ErbB2 mRNA and protein. Remarkably, the elevated expression of ErbB2 also correlates with selective genomic amplification of the activated *ErbB2* allele (Andrechek et al. 2000; Montagna et al. 2002; Hodgson et al. 2005). Thus, as in human breast cancers, amplification of *ErbB2* appears to be a critical event in mammary tumor progression in this mouse model. Its similarity to ERBB2-initiated human breast cancer has been further highlighted by detailed comparative genome hybridization (CGH) analyses of the *ErbB2* amplicon. The results of these studies revealed that both human and mouse *ErbB2* amplicons contain the same core 10 genes closely linked to the *ErbB2* locus (Hodgson et al. 2005).

Tumorigenesis in the ErbB2^{KI} model is also associated with a number of other alterations, including centrosome abnormalities and recurrent deletions of chromosome 4 (Montagna et al. 2002). Further refined mapping of chromosome 4 region has revealed that these tumors

have frequently loss of expression of the tumor suppressor 14-3-3 σ (Hodgson et al. 2005). Interestingly, loss of 14-3-3 σ expression has been noted in many primary human breast cancers (Ferguson et al. 2000; Urano et al. 2002; Vercoutter-Edouart et al. 2001). One potential explanation for selection for loss of 14-3-3 σ expression in ErbB2-induced tumors is that 14-3-3 σ is involved in cytoplasmic sequestration of the EGR2 transcription factor, which plays a critical role in up-regulating ErbB2 expression (Dillon et al. 2007a). Future studies involving the targeted disruption of 14-3-3 σ in mammary epithelium should establish the relative contribution loss that 14-3-3 σ plays in ErbB2-driven mammary tumor progression.

Another important issue influencing breast cancer morbidity is recurrence of the cancer after a period of remission. It has been hypothesized that recurrence of a cancer is a consequence of emergence of tumor cells from dormancy (White et al. 2006), and this phenomenon has recently been experimentally validated in several studies. For example, although down-regulation of ErbB2 in an inducible model of mammary tumorigenesis initially results in the regression of mammary tumors, these eventually re-emerge—presumably from dormant tumor cells—after a long latency period (Moody et al. 2002). This re-emergence correlates with up-regulation of the Snail transcription factor in these cells (Moody et al. 2005). Interestingly, Snail controls normal development, epithelial-mesenchymal transitions, and cancer stem cell properties (reviewed in Cobaleda et al. 2007).

IDENTIFICATION OF SIGNALING NETWORKS CRITICAL FOR ONCOGENE-INDUCED MAMMARY TUMOR PROGRESSION

The *in vivo* importance of oncogene-coupled signaling in mammary tumor progression has recently been shown by experiments using transgenic mice that express phosphorylation mutants of the ErbB2 or PyV mT oncogenes that decouple these oncogenes from particular signaling pathways in the mammary epithelium

E.Y.H.P. Lee and W.J. Muller

(Dankort et al. 2001; Schade et al. 2007). Although mammary epithelial expression of ErbB2 mutants coupled specifically to either the adaptor protein ShcA or Grb2 were capable of efficiently inducing mammary tumors (Dankort et al. 2001), they had dramatically different metastatic outcomes that correlated with distinct transcriptional profiles (Dankort et al. 2001; Schade et al. 2007).

Direct evidence for the importance of ShcA in mammary tumor progression has recently been obtained from studies interbreeding the MMTV/ErbB2 IRES Cre strain (NIC) and a separate strain of mice harboring a conditional knockout of ShcA (Ursini-Siegel et al. 2008). Remarkably, mammary-specific disruption of ShcA in MMTV/NIC mice completely blocked mammary tumor progression in virgin females, suggesting that ShcA signaling is critical for mammary tumor induction. However, the precise role of the ShcA-coupled signaling pathways in ErbB2-induced mammary tumor progression remains to be addressed.

Another important signaling pathway that plays a critical role in ErbB2-induced mammary tumor progression is the c-Src tyrosine kinase. We have previously shown that Src family kinases are activated in mammary tumors derived from MMTV-activated *ErbB-2* transgenic strains (Guy et al. 1994; Muthuswamy and Muller 1995a,b; Muthuswamy et al. 1994). In the case of ErbB-2, activation of c-Src occurred through the direct interaction of the catalytic domain of c-Src with the catalytic domain of ErbB-2 (Kim et al. 2005; Marcotte et al. 2009). Recruitment of c-Src to chimeric EGFR/ErbB-2 receptors possessing the ErbB-2 catalytic domain resulted in disruption of epithelial polarity (Kim et al. 2005; Marcotte et al. 2009).

ONCOGENE-INTEGRIN CROSS TALK IN MAMMARY TUMOR PROGRESSION

An increasing body of evidence suggests oncogenes can activate other classes of transmembrane receptors. For instance, genetic and biochemical analyses have revealed that the $\alpha\beta 1$ integrin receptor can associate with activated members of the EGFR family (Weaver

et al. 1997). Stimulation of $\beta 1$ integrin function can thereby result in the enhanced phosphorylation of EGFR family members (Adelsman et al. 1999; Mariotti et al. 2001; Moro et al. 1998). Moreover, activation of TGF- β signaling can induce coclustering of $\beta 1$ -integrin and ErbB2, and thereby promote metastatic invasion (Wang et al. 2009a). Indeed, coexpression of activated TGF- β receptor or its ligand and ErbB2 in the mammary epithelium of transgenic mice dramatically accelerates metastatic progression (Muraoka-Cook et al. 2004; Muraoka et al. 2003; Siegel et al. 2003). It has also been shown that antibodies directed against $\beta 1$ integrin down-regulate EGFR phosphorylation and can interfere with the morphological transformation of breast tumor cells (Wang et al. 2002; Weaver et al. 1997).

Direct evidence for a role for integrin in mammary tumor progression comes from a recent study involving the conditional ablation of $\beta 1$ integrin in the PyV mT model. Lack of $\beta 1$ integrin causes PyV-mT-induced mammary tumors to disappear (Andrechek et al. 2005). Interestingly, this study also showed that tyrosine phosphorylation and subcellular localization of the integrin-associated kinase FAK was compromised in the $\beta 1$ -integrin-null tumor cells (White et al. 2004). This block in tumor progression was associated with a complete block in cell cycle progression without any overt evidence of apoptotic cell death (Andrechek et al. 2005). It was not because of an indirect effect on normal mammary development, because the initial stages of mammary epithelial ductal outgrowth were not perturbed (White et al. 2004). The $\beta 1$ -integrin-deficient PyV mT tumor cells were not able to contribute to tumor development and showed many of the hallmarks of tumor dormancy (White et al. 2004). The mammary tumors that eventually arose were identified as “escapee” populations of epithelial cells: These cells failed to express Cre recombinase owing to the stochastic nature of its expression in MMTV/Cre transgenic strains and thus retained $\beta 1$ -integrin function (White et al. 2004). Conditional disruption of the integrin-coupled FAK pathway also resulted in a profound block in tumor progression

(Lahlou et al. 2007; Luo et al. 2009; Provenzano et al. 2008; Pylayeva et al. 2009). Further evidence for importance of integrin signaling stems from a collaborative study of mice carrying a mutant $\beta 4$ integrin lacking the carboxy-terminal signaling domain as well as the MMTV/activated *ErbB2* transgene (Guo et al. 2006). The mutant $\beta 4$ integrin retains the capacity to form hemidesmosomes but cannot signal, owing to truncation of the carboxy-terminal domain (Guo et al. 2006). These mice had a moderate impairment of *ErbB2*-induced tumor progression, delaying tumor onset 5 weeks (Guo et al. 2006).

ErbB2 SIGNALING IMPACTS ON CELL CYCLE PROGRESSION AND TRANSCRIPTION FACTOR NETWORKS

Ultimately, signaling from activated *ErbB2* affects transcription factors and the machinery that controls the cell cycle. Germ-line loss of cyclin D1 in the MMTV/*ErbB2* transgenic strains results in a complete block in tumor progression (Yu et al. 2001). Functional Cdk4, a known cyclin D1 partner, is also required for *ErbB2*-induced tumor progression (Landis et al. 2006; Yu et al. 2006). Conversely, modulation of the levels of the p16 and p27 inhibitors of Cdk function can also have profound effect on *ErbB2*-driven mammary tumor progression (Bulavin et al. 2004; Muraoka et al. 2002). Key elements of the cell cycle machinery can thus be modulated by oncogenic *ErbB2* signaling.

ErbB2-mediated activation of key transcription factors can also influence tumor progression. For example, disruption of Stat3 in mammary epithelium can have a profound impact on *ErbB2*-driven tumor progression. Although *ErbB2* can induce tumors in the absence of Stat3, these tumors fail to metastasize (Ranger et al. 2009). Examination of the gene expression profile of Stat3-deficient tumors revealed that they failed to induce the expression of a number of Stat3 target genes involved in inflammation (Ranger et al. 2009). Disruption of certain key signaling networks thus appears to have selective effects on the

Oncogenes and Tumor Suppressor Genes

metastatic phase of *ErbB2*-driven tumor progression.

RB AND CELL CYCLE REGULATION

RB regulates the G1-S phase transition by repressing E2F transcription factors that stimulate the expression of genes required for cell cycle progression (reviewed in Burkhardt and Sage 2008). Cyclin D1-CDK4 complex phosphorylates RB, leading to its dissociation from the RB-E2F complex, which frees E2F for activation of cyclin D1 and S-phase genes transcription. E2F targets include cell cycle regulatory genes such as cyclin D1 and S-phase genes. The positive feedback loop among RB, E2F, and cyclin D1 allows for cell cycle progression through G1/S and S phases. Deregulation of RB pathways occurs in most cancer and is mediated either by loss-of-function mutation of negative players including RB and CDK inhibitors (CKIs, p15, p16, p21, etc.) or by amplification or overexpression of cyclin D1. Many of these regulators have additional functions; for example, RB plays a role in adipocyte differentiation (Chen et al. 1996) and cyclin D1 has a critical role in development (Bienvenu et al. 2010).

THE p53 PATHWAY

Rare germ-line mutation of *TP53* leads to the Li-Fraumeni familial cancer syndrome. On the other hand, somatic mutations of *TP53* occur in the majority of sporadic cancers. p53 responds to various cellular stress signals and is hailed as the “guardian of genome” (Lane 1992). p53 alteration is common in spontaneously immortalized murine embryonic fibroblasts (Harvey and Levine 1991).

In normal cells, p53 is kept at very low level by p53-interacting protein, MDM2. MDM2 is an E3 ubiquitin ligase that promotes ubiquitination and subsequent proteasomal degradation of p53. Cellular stress induces post-translational modification of p53 and MDM2, leading to stabilization and activation of p53 (reviewed in Levine and Oren 2009; Brown et al. 2009). Mdm2 appears to be intrinsically

E.Y.H.P. Lee and W.J. Muller

active and its deletion results in embryonic lethality (reviewed in Donehower and Lozano 2009; Kruse and Gu 2009). Deletion of *TP53* rescues embryonic lethality in *Mdm2*- and *Mdm4*-deficient mice (Jones et al. 1995; Montes de Oca Luna et al. 1995; Parant et al. 2001). Furthermore, mice carrying a knockin *TP53* allele that shows defective transactivation and *Mdm2* binding die during embryogenesis (Johnson et al. 2005). Thus, unrepressed *Mdm2* activity is a likely cause of embryonic lethality. Notably, a single nucleotide polymorphism within the first intron of *Mdm2* has been shown to affect promoter strength and affect the level and activities of p53 subsequently (Bond et al. 2004). These studies provide genetic and biochemical evidence of regulation of p53 by MDM2 and MDM4.

p53 binds to DNA in a sequence-specific manner to induce cell cycle checkpoint activation, cellular senescence, apoptosis, or autophagy (reviewed in Menendez et al. 2009; Zilfou and Lowe 2009). Expression of hundreds of genes is directly regulated by p53. For examples, CKI *p21* and Polo-like kinase 1 are important targets of p53 for G1/S and G2-M cell cycle checkpoint activation, respectively (Sur et al. 2009). Centrosome amplification occurs frequently in cells harboring *TP53* mutations, partly through overactivation of CDKs by cyclin E and cyclin A (reviewed in Fukasawa 2008). This in turn results in chromosome instability, a hallmark of solid tumors.

Loss of p53 allows efficient reprogramming of somatic cells (Kawamura et al. 2009). In mammary gland, loss of p53 result in aberrant asymmetric cell divisions of mammary stem cells (Cicalese et al. 2009). p53 also enhances the maturation of several microRNAs with growth-suppressive functions (Suzuki et al. 2009). These recent results continue to expand our understanding of the tumor suppressor mechanisms of p53.

p53 is inactivated in the cancer stem cells of ErbB2-associated mouse mammary tumors (Cicalese et al. 2009). In this model, cancer stem cells undergo frequent symmetric cell division and expansion. Reactivation of p53 by pharmacological intervention restores asymmetric cell

division (Cicalese et al. 2009). In a separate model, expression of the exon-5-6-less *TP53* allele in neural stem cells was followed by the expansion of transient-amplifying progenitor-like cells and subsequent development of malignant astrocytic glioma (Wang et al. 2009b). These studies indicate a role of p53 in the regulation of stem cell division in addition to its well-documented role in transformation.

The majority of cancer-associated mutations in *TP53* are missense mutations of the DNA binding domain. Some mutations induce genetic instability by inactivating ATM (reviewed in Xu 2008). Mouse models of missense mutations corresponding to those in human mutant *TP53* reveal a gain-of-function. These mutant p53 interact with Mre11 and suppress binding of Mre11-Rad50-NBS1 complex to DNA double-stranded breaks (Song et al. 2007). Although mutant p53 levels are increased in most tumors, only low levels of mutant p53 are present in normal mouse tissues initially, which indicates additional events are needed to allow mutant p53 to escape MDM2-dependent degradation (reviewed in Brosh and Rotter 2009).

THE BRCA PATHWAYS

Individuals with mutations in *BRCA1* are predisposed to breast and ovarian cancers. Earlier studies indicated that the majority of *BRCA1*-associated cancers are triple (ER α , PR, and HER-2) negative, basal-type breast cancers (reviewed in Lynch et al. 2008). However, recent reports show that *BRCA1*-associated breast cancers can be ER α - and PR-positive, especially in aged patients (Atchley et al. 2008; Tung et al. 2010). The *BRCA1* gene encodes a protein of 1863 amino acids with a predicted molecular weight of 220 kDa (Miki et al. 1994). At the amino terminus is a RING domain that mediates the interaction with another RING-domain-containing protein, BARD1. The *BRCA1*/BARD1 heterodimer shows ubiquitin ligase activity (Chen et al. 2002; Wu-Baer et al. 2003). Exon 11 of *Brca1* encodes approximately 60% of *BRCA1*. This central region interacts with the DNA repair protein complex Mre11-Rad50-NBS1 and

the transcriptional repressor ZBRK1 (Zhong et al. 1999; Zheng et al. 2000). The Mre11-Rad50-NBS1 complex binds to and processes DNA double stranded breaks. This complex is involved in both nonhomologous end joining and homologous recombinational repair. Mouse embryonic fibroblasts deficient in *Brca1* have impaired homologous recombinational repair (Moynahan et al. 1999). Loss of 53BP1 expression restores homologous recombination in *Brca1* mutant cells by promoting ATM-dependent processing of broken DNA ends (Bunting et al. 2010). The carboxyl terminus of BRCA1 contains tandem BRCA1 C-terminal (BRCT) repeats. This region binds to phospho-peptides (Manke et al. 2003; Yu et al. 2003) involved in cell-cycle checkpoints and DNA repair. Several proteins including BACH1, CtIP, Acetyl-CoA carboxylase, Abraxas/CCDC98, and RAP80 interact with the BRCT domain of BRCA1 in a phospho-dependent manner (reviewed in Rodriguez and Songyang 2008). These studies reveal how BRCA1 maintains genomic stability through DNA repair and cell cycle checkpoint activation.

Three BRCA1 protein complexes have been characterized. One complex contains BRCA2 and PALB2, a BRCA2-associated protein (Sy et al. 2009; Zhang et al. 2009). Germ-line mutations of *PALB2* not only associate with hereditary breast cancer, but also is found in Fanconi's anemia and familial pancreatic cancer syndrome. *BRCA2* mutations have been linked to a wide spectrum of cancers. In contrast, *BRCA1* mutation is primarily associated with breast and ovarian cancers. Plausible explanations include the connection between BRCA1, but not BRCA2, and steroid hormone receptors (reviewed in Lee 2008) and the role of BRCA1 in mammary luminal epithelial lineage determination (reviewed in Visvader 2009).

Developmental defects and early embryonic death have been observed in *Brca1* homozygous knockout mice (reviewed in Dasika et al. 1999; Drost and Jonkers 2009). Several further lines of evidence indicate that BRCA1 plays a role in mammary development and differentiation. Mouse mammary tissue harboring a conditional *Brca1* knockout displays abnormal ductal

morphogenesis (Xu et al. 1999). Depletion of BRCA1 by RNAi in 3-D cultured MECs impairs acinus formation and up-regulates genes that control proliferation but down-regulates those controlling differentiation (Furuta et al. 2005). Importantly, the epithelial progenitor cell population is expanded in *BRCA1* carriers. This highlights the haplo-insufficiency phenotype of BRCA1 in mammary epithelial differentiation (Lim et al. 2009). The EGFR pathway is activated in the mammary epithelial cells of *BRCA1* carriers (Burga et al. 2009), and may be linked to the expansion of luminal epithelial cells. In contrast, knocking down BRCA1 in primary mammary epithelial cells increases the number of stem/progenitor cells and decreases ER α expression. Loss of heterozygosity of *BRCA1* in the mammary epithelial cells of *BRCA1* carriers leads to the expansion of ALDH1-positive, ER- and PR-negative stem/progenitor cells (Liu et al. 2008). Thus, the effects of BRCA1 on mammary cell differentiation appear to be dosage-dependent (Ginestier et al. 2009).

BRCA2 carriers develop different subtypes of breast cancer. The *BRCA2* gene encodes a protein of 3418 amino acids with a predicted molecular weight of 375 kDa (Wooster et al. 1995). A repeat motif termed the BRC domain that comprises approximately 70 amino acids is present in the middle third of the BRCA2 protein. BRC repeats bind to the Rad51 recombinase (Chen et al. 1998; Wong et al. 1997). *BRCA2*-deficient cells show defective formation of IR-induced Rad51 foci (Yuan et al. 1999). At the carboxyl terminus of BRCA2 is a region with extensive secondary structure that interacts with the evolutionarily conserved protein DSS1. Based on the 3-D structure, it is predicted that the high-affinity ssDNA-binding and dsDNA-binding domains of BRCA2 play critical roles in homologous recombination (Yang et al. 2002). Indeed, both the formation of Rad51 nuclear filaments and Rad51-mediated strand exchange during homologous recombination are regulated by BRCA2 (reviewed in Thorslund and West 2007).

Brca2 homozygous knockout mice either die during embryogenesis or survive beyond birth, depending on the mutation (Dasika et al.

E.Y.H.P. Lee and W.J. Muller

1999). Mutations in either *p53* or *p21* prolong the survival of *Brca1*^{-/-} and *Brca2*^{-/-} embryos. Several conditional knockout models of these two tumor suppressor genes have been established (see the following).

TRANSGENIC MOUSE MODELS OF BRCA-ASSOCIATED BREAST CANCER

Several different *Brca1* alleles have been generated (Drost and Jonkers 2009). Because of differential splicing, the exon 11-less *Brca1* isoform is also present in wild-type mice and the protein product is located in the nucleus—like the full-length *Brca1* (Huber et al. 2001). All other alleles are predicted to produce aberrant protein products. Different mutant *TP53* alleles, including null and internally truncated alleles, are generated, which can result in different phenotype. Indeed, mutant but not null mutation promotes expansion of neuronal progenitor cells and subsequent astrocytic glioma formation (Wang et al. 2009b). *Cre* transgenes under the control of various different promoters have been used for the conditional inactivation of *Brca1*. Thus, these models differ in the nature of mutation as well as cell types expressing mutant alleles.

The WAPCre or MMTVCre*Brca1*^{Δ11} Model

Whey acidic protein (WAP) or MMTV promoter driven *Cre* expression leads to the deletion of exon 11 of the *Brca1* gene in the mammary gland (Xu et al. 1999). Mammary tumors developed in these mice at low frequency after the 10–13 months of latency. Introduction of heterozygous *p53* mutation significantly shortened tumor latency (Brodie et al. 2001). The absence of *Brca1* turns the proliferation of ERα-positive cells to an autocrine fashion in contrast to that of the wild-type mice in that estrogen mainly induces proliferation of ERα-negative cells in the mammary gland (Li et al. 2007). This model indicates that exon 11-less *Brca1* isoform is deficient in tumor suppression. Within exon 11, a unique ATM phosphorylation site exists. Mice homozygous for the phosphorylation mutant show

aging phenotype and have elevated irradiation-induced tumorigenesis (Kim et al. 2009).

The WAPCre^c*Brca1*^{Δ11}*p53*^{Δ5&6} Model

Transgenic mice have been generated that carry a constitutively active whey acidic protein promoter-*Cre* transgene (*WAPCre^c*) that is expressed in both basal myoepithelial and luminal epithelial cells (Lin et al. 2004). Because expression of this *Cre* transgene does not require pregnancy, the interaction between tumor suppressor genes and ovarian hormones can be addressed. In contrast to wild-type and *p53*-deficient mammary epithelial cells, proliferation of *Brca1*- and *p53*-deficient mammary epithelial cells was uniquely sensitive to progesterone. Furthermore, progesterone receptors were stabilized in these cells. Antiprogesterone treatment in pubertal mice prevented or delayed mammary tumorigenesis (Poole et al. 2006). These mice develop mammary tumors with full penetrancy and a median tumor latency of 6.3 months (Shafee et al. 2008). In line with recent reports (Atchley et al. 2008; Tung et al. 2010), both ERα-positive and ERα-negative tumors were identified and *p63* expression was up-regulated. These findings raise the possibility that the cells of origin in *BRCA1*-mediated breast carcinogenesis may be heterogeneous.

The K14-Cre*Brca1*^{Δ5-13}*p53*^{Δ2-10} Model

Keratin 14-*Cre* transgene mediated deletion of *Brca1* and *p53* genes affect epithelial tissues, including mammary myoepithelial cells and the skin epithelium (Liu et al. 2007). A high incidence of poorly differentiated ERα-negative basal tumors developed with a medium tumor latency of 7.0 months, which is consistent with the fact that the keratin 14 promoter drives robust expression in myoepithelial cells.

The β-lactoglobulin-Cre*Brca1*^{Δ22-24}*p53*^{+/-} Model

In this model, mammary tumors developed with a median latency of 7.0 months (McCarthy et al. 2007). The *β-lactoglobulin* is highly expressed in luminal epithelial cells of the mammary

gland but deletion of the C-terminus of *Brca1* leads to expression of basal-like markers in tumor.

The *Brca1* and *p53* mutations and the cell type(s) targeted differ in each of the above models. Together, the findings from these models and recent studies of *BRCA1*-associated breast cancer indicate that tumor cells can arise from multiple cell types in the mammary gland.

The K14-Cre*Brca2*^{Δ11}*p53*^{Δ2-10} Model

The exon 11-less *Brca2* embryos died before E9.5. The K14-Cre transgenic approach, however, restricts deletion of *Brca2* exon 11 and *p53* exons 2-10 to the skin, myoepithelial and luminal epithelial cells in the mammary gland, and some other tissues (Jonkers et al. 2001). Mammary and skin tumors developed at high frequencies in these mice (Jonkers et al. 2001). Overall, 5%–30% of adult mammary epithelial cells were targeted in this model; in contrast, 70%–90% were targeted in the *WAPCre*^c transgenic model (Lin et al. 2004).

The WAP-Cre*Brca2*^{Δ3-4/-} Model

Knocking a *Cre* gene into the *WAP* locus leads to a 77% incidence of mammary tumors after approximately 1.4-year latency (Ludwig et al. 2001). Surprisingly, invasive adenocarcinomas that are histologically uniform developed in these mice, in contrast to the different breast cancer subtypes seen in *BRCA2* carriers, in which loss of heterozygosity leads to tumorigenesis.

The MMTV-Cre*Brca2*^{Δ9&10} Model

This *Brca2* mutation did not affect pregnancy-induced mammary epithelial expansion and involution (Cheung et al. 2004). Mammary adenocarcinomas developed after approximately 1.6 years latency, and mutations of the *p53* gene were identified in the tumors. Introduction of a *Brca2* mutant allele into *p53*^{+/-} mice significantly skewed the tumor spectrum toward mammary adenocarcinoma.

Despite the fact that both *BRCA1* and *BRCA2* have a role in DNA double strand break repair,

Oncogenes and Tumor Suppressor Genes

studies of genetically engineered *Brca1* and *Brca2* transgenic mice clearly reveal unique biological functions of these two proteins.

TARGETING ABERRANT PATHWAYS IN BREAST CANCERS

BRCA-deficient mammary tumors are sensitive to cisplatin but not doxorubicin treatment, which is consistent with the observation that *BRCA*-deficient cells have double strand break repair defects (Shafee et al. 2008). In a small clinical trial, cisplatin treatment for *BRCA1*-associated breast cancers appears to be promising (Byrski et al. 2009). Similarly, *BRCA1*- and *BRCA2*-associated ovarian cancers were responsive to cisplatin or carboplatin treatment; however, resistance developed (Cass et al. 2003). Secondary mutations that restore the open reading frame of *BRCA2* and its DNA repair function were identified in carboplatin-resistant tumors (Edwards et al. 2008; Sakai et al. 2008). Using a synthetic lethality approach, several labs have concluded that *BRCA* deficiency sensitizes cells to poly ADP-ribose polymerase (PARP) inhibitors (see Ashworth and Bernards 2008). Indeed, *Brca1*-associated mammary tumors have been shown to be highly sensitive to PARP inhibitors (Rottenberg et al. 2008).

Breast cancer stem cells became enriched on chemotherapy (Creighton et al. 2009). The CD29^{hi}CD24^{med} cancer stem cells in tumors developed in *WAPCre^cBrca1^{Δ11}p53^{Δ5&6}* mice expanded in cisplatin-refractory tumors (Shafee et al. 2008). In several breast cancer cell lines, expansion of CD44^{hi}CD24^{med} cells was associated with herceptin resistance (Reim et al. 2009). Strategies to target specific pathways required for the maintenance of cancer stem cells are being developed. For example, metformin, a diabetes drug, (Hirsch et al. 2009) and salinomycin have been shown to target breast cancer stem cells selectively (Gupta et al. 2009). However, differentiation of CD29^{hi}CD24^{med} cancer stem cells to CD29⁻CD24⁺ cells is likely to be subject to feedback regulation similar to that observed in normal stem cells (Lander 2009). Understanding the mechanisms

E.Y.H.P. Lee and W.J. Muller

underlying such feedback control will provide additional insights into how one might eradicate cancer stem cells.

CONCLUSIONS

Breast cancers are known for their heterogeneity, which is also reflected in the expression of cancer stem cell surface markers (Hwang-Versluis et al. 2009). There are several schools of thoughts on the origin of heterogeneity in cancer: stochastic events (Nowell 1976), the cell of origin (Visvader 2009), expansion of cancer stem cells (Pece et al. 2010), and inter tumor genetic diversity (Park et al. 2010). Results from different cancer models indicate that *Brcal* inactivation in different cell populations can lead to different tumor subtypes. As the tumor progresses, its cancer stem cell content may vary.

Several pathways critical to breast tumorigenesis have been identified and effective pharmacological interventions, including the targeting of steroid hormone receptors and ErbB2 pathways, are now available. To prevent cancer recurrence, new strategies such as targeting of DNA repair pathways, reactivating p53 (reviewed in Brown et al. 2009; Shangary and Wang 2009), and inhibiting cancer stem cells, are being developed. Meanwhile, advances in our understanding of breast cancer oncogenes and tumor suppressor genes will continue to provide insights critical to the development of novel anti-cancer approaches.

REFERENCES

- Ackler S, Ahmad S, Tobias C, Johnson MD, Glazer RI. 2002. Delayed mammary gland involution in MMTV-AKT1 transgenic mice. *Oncogene* **21**: 198–206.
- Adelman MA, McCarthy JB, Shimizu Y. 1999. Stimulation of β 1-integrin function by epidermal growth factor and heregulin- β has distinct requirements for ErbB2 but a similar dependence on phosphoinositide 3-OH kinase. *Mol Biol Cell* **10**: 2861–2878.
- Allred DC, Clark GM, Molina R, Tandon AK, Schnitt SJ, Gilchrist KW, Osborne CK, Tormey DC, McGuire WL. 1992. Overexpression of HER-2/neu and its relationship with other prognostic factors change during the progression of in situ to invasive breast cancer. *Hum Pathol* **23**: 974–979.
- Andrechek ER, White D, Muller WJ. 2005. Targeted disruption of ErbB2/Neu in the mammary epithelium results in impaired ductal outgrowth. *Oncogene* **24**: 932–937.
- Andrechek ER, Hardy WR, Siegel PM, Rudnicki MA, Cardiff RD, Muller WJ. 2000. Amplification of the neu/erbB-2 oncogene in a mouse model of mammary tumorigenesis. *Proc Natl Acad Sci* **97**: 3444–3449.
- Andrulis IL, Bull SB, Blackstein ME, Sutherland D, Mak C, Sidlofsky S, Pritzker KP, Hartwick RW, Hanna W, Lickley L, et al. 1998. neu/erbB-2 amplification identifies a poor-prognosis group of women with node-negative breast cancer. Toronto Breast Cancer Study Group. *J Clin Oncol* **16**: 1340–1349.
- Antoniotti S, Taverna D, Maggiora P, Sapei ML, Hynes NE, De Bortoli M. 1994. Oestrogen and epidermal growth factor down-regulate erbB-2 oncogene protein expression in breast cancer cells by different mechanisms. *Br J Cancer* **70**: 1095–1101.
- Atchley DP, Albarracin CT, Lopez A, Valero V, Amos CI, Gonzalez-Angulo AM, Hortobagyi GN, Arun BK. 2008. Clinical and pathologic characteristics of patients with BRCA-positive and BRCA-negative breast cancer. *J Clin Oncol* **26**: 4282–4288.
- Bell DW. 2010. Our changing view of the genomic landscape of cancer. *J Pathol* **220**: 231–243.
- Bienvenu F, Jirawatnotai S, Elias JE, Meyer CA, Mizeracka K, Marson A, Frampton GM, Cole ME, Odom DT, Odajima J, et al. 2010. Transcriptional role of cyclin D1 in development revealed by a genetic-proteomic screen. *Nature* **463**: 374–378.
- Bond GL, Hu W, Bond EE, Robins H, Lutzker SG, Arva NC, Bargonetti J, Bartel F, Taubert H, Wuerl P, et al. 2004. A single nucleotide polymorphism in the MDM2 promoter attenuates the p53 tumor suppressor pathway and accelerates tumor formation in humans. *Cell* **119**: 591–602.
- Bouchard L, Lamarre L, Tremblay PJ, Jolicoeur P. 1989. Stochastic appearance of mammary tumors in transgenic mice carrying the MMTV/c-neu oncogene. *Cell* **57**: 931–936.
- Brandt R, Eisenbrandt R, Leenders F, Zschiesche W, Binas B, Juergensen C, Theuring F. 2000. Mammary gland specific hEGF receptor transgene expression induces neoplasia and inhibits differentiation. *Oncogene* **19**: 2129–2137.
- Brodie SG, Xu X, Qiao W, Li WM, Cao L, Deng CX. 2001. Multiple genetic changes are associated with mammary tumorigenesis in *Brcal* conditional knockout mice. *Oncogene* **20**: 7514–7523.
- Brosh R, Rotter V. 2009. When mutants gain new powers: News from the mutant p53 field. *Nat Rev Cancer* **9**: 701–713.
- Brown CJ, Lain S, Verma CS, Fersht AR, Lane DP. 2009. Awakening guardian angels: Drugging the p53 pathway. *Nat Rev Cancer* **9**: 862–873.
- Bulavin DV, Phillips C, Nannenga B, Timofeev O, Donehower LA, Anderson CW, Appella E, Fornace AJ Jr. 2004. Inactivation of the Wip1 phosphatase inhibits mammary tumorigenesis through p38 MAPK-mediated activation of the p16(Ink4a)-p19(Arf) pathway. *Nat Genet* **36**: 343–350.
- Bunting SE, Callén E, Wong N, Chen HT, Polato F, Gunn A, Bothmer A, Feldhahn N, Fernandez-Capetillo O, Cao L, et al. 2010. 53BP1 inhibits homologous recombination in *Brcal*-deficient cells by blocking resection of DNA breaks. *Cell* **141**: 243–254.

Oncogenes and Tumor Suppressor Genes

- Burga LN, Tung NM, Troyan SL, Bostina M, Konstantinopoulos PA, Fountzilas H, Spentzos D, Miron A, Yassin YA, Lee BT, et al. 2009. Altered proliferation and differentiation properties of primary mammary epithelial cells from BRCA1 mutation carriers. *Cancer Res* **69**: 1273–1278.
- Burkhardt DL, Sage J. 2008. Cellular mechanisms of tumour suppression by the retinoblastoma gene. *Nat Rev Cancer* **8**: 671–682.
- Byrski T, Huzarski T, Dent R, Gronwald J, Zuziak D, Cybulski C, Kladny J, Gorski B, Lubinski J, Narod SA. 2009. Response to neoadjuvant therapy with cisplatin in BRCA1-positive breast cancer patients. *Breast Cancer Res Treat* **115**: 359–363.
- Cass I, Baldwin RL, Varkey T, Moslehi R, Narod SA, Karlan BY. 2003. Improved survival in women with BRCA-associated ovarian carcinoma. *Cancer* **97**: 2187–2195.
- Chen PL, Chen CF, Chen Y, Xiao J, Sharp ZD, Lee WH. 1998. The BRC repeats in BRCA2 are critical for RAD51 binding and resistance to methyl methanesulfonate treatment. *Proc Natl Acad Sci* **95**: 5287–5292.
- Chen A, Kleiman FE, Manley JL, Ouchi T, Pan ZQ. 2002. Autoubiquitination of the BRCA1*BARD1 RING ubiquitin ligase. *J Biol Chem* **277**: 22085–22092.
- Chen PL, Riley DJ, Chen Y, Lee WH. 1996. Retinoblastoma protein positively regulates terminal adipocyte differentiation through direct interaction with C/EBPs. *Genes Dev* **10**: 2794–2804.
- Cheung AM, Elia A, Tsao MS, Done S, Wagner KU, Hennighausen L, Hakem R, Mak TW. 2004. Brca2 deficiency does not impair mammary epithelium development but promotes mammary adenocarcinoma formation in *p53*^{+/−} mutant mice. *Cancer Res* **64**: 1959–1965.
- Cicalese A, Bonizzi G, Pasi CE, Faretta M, Ronzoni S, Giulini B, Briskin C, Minucci S, Di Fiore PP, Pelicci PG. 2009. The tumor suppressor p53 regulates polarity of self-renewing divisions in mammary stem cells. *Cell* **138**: 1083–1095.
- Cobaleda C, Pérez-Caro M, Vicente-Dueñas C, Sánchez-García I. 2007. Function of the zinc-finger transcription factor SNAI2 in cancer and development. *Annu Rev Genet* **41**: 41–61.
- Creighton CJ, Li X, Landis M, Dixon JM, Neumeister VM, Sjolund A, Rimm DL, Wong H, Rodriguez A, Herschkowitz JJ, et al. 2009. Residual breast cancers after conventional therapy display mesenchymal as well as tumor-initiating features. *Proc Natl Acad Sci* **106**: 13820–13825.
- Croce CM. 2009. Causes and consequences of microRNA dysregulation in cancer. *Nat Rev Genet* **10**: 704–714.
- Dalglish GL, Furge K, Greenman C, Chen L, Bignell G, Butler A, Davies H, Edkins S, Hardy C, Latimer C, et al. 2010. Systematic sequencing of renal carcinoma reveals inactivation of histone modifying genes. *Nature* **463**: 360–363.
- Dankort D, Maslikowski B, Warner N, Kanno N, Kim H, Wang Z, Moran MF, Oshima RG, Cardiff RD, Muller WJ. 2001. Grb2 and Shc adapter proteins play distinct roles in Neu (ErbB-2)-induced mammary tumorigenesis: Implications for human breast cancer. *Mol Cell Biol* **21**: 1540–1551.
- Dasika GK, Lin SC, Zhao S, Sung P, Tomkinson A, Lee EY. 1999. DNA damage-induced cell cycle checkpoints and DNA strand break repair in development and tumorigenesis. *Oncogene* **18**: 7883–7899.
- Dillon RL, White DE, Muller WJ. 2007b. The phosphatidylinositol 3-kinase signaling network: Implications for human breast cancer. *Oncogene* **26**: 1338–1345.
- Dillon RL, Brown ST, Ling C, Shioda T, Muller WJ. 2007a. An EGR2/CITED1 transcription factor complex and the 14-3-3σ tumor suppressor are involved in regulating ErbB2 expression in a transgenic-mouse model of human breast cancer. *Mol Cell Biol* **27**: 8648–8657.
- Dillon RL, Marcotte R, Hennessy BT, Woodgett JR, Mills GB, Muller WJ. 2009. Akt1 and Akt2 play distinct roles in the initiation and metastatic phases of mammary tumor progression. *Cancer Res* **69**: 5057–5064.
- Donehower LA, Lozano G. 2009. 20 years studying p53 functions in genetically engineered mice. *Nat Rev Cancer* **9**: 831–841.
- Ding L, Ellis MJ, Li S, Larson DE, Chen K, Wallis JW, Harris CC, McLellan MD, Fulton RS, Fulton LL, et al. 2010. Genome remodelling in a basal-like breast cancer metastasis and xenograft. *Nature* **464**: 999–1005.
- Dourdin N, Schade B, Lesurf R, Hallett M, Munn RJ, Cardiff RD, Muller WJ. 2008. Phosphatase and tensin homologue deleted on chromosome 10 deficiency accelerates tumor induction in a mouse model of ErbB-2 mammary tumorigenesis. *Cancer Res* **68**: 2122–2131.
- Drost RM, Jonkers J. 2009. Preclinical mouse models for BRCA1-associated breast cancer. *Br J Cancer* **101**: 1651–1657.
- Edwards SL, Brough R, Lord CJ, Natrajan R, Vatcheva R, Levine DA, Boyd J, Reis-Filho JS, Ashworth A. 2008. Resistance to therapy caused by intragenic deletion in BRCA2. *Nature* **451**: 1111–1115.
- Ferguson AT, Evron E, Umbricht CB, Pandita TK, Chan TA, Hermeking H, Marks JR, Lambers AR, Futreal PA, Stampfer MR, et al. 2000. High frequency of hypermethylation at the 14-3-3σ locus leads to gene silencing in breast cancer. *Proc Natl Acad Sci* **97**: 6049–6054.
- Fukasawa K. 2008. P53, cyclin-dependent kinase and abnormal amplification of centrosomes. *Biochim Biophys Acta* **1786**: 15–23.
- Furuta S, Jiang X, Gu B, Cheng E, Chen PL, Lee WH. 2005. Depletion of BRCA1 impairs differentiation but enhances proliferation of mammary epithelial cells. *Proc Natl Acad Sci* **102**: 9176–9181.
- Garrett TP, McKern NM, Lou M, Elleman TC, Adams TE, Lovrecz GO, Kofler M, Jorissen RN, Nice EC, Burgess AW, et al. 2003. The crystal structure of a truncated ErbB2 ectodomain reveals an active conformation, poised to interact with other ErbB receptors. *Mol Cell* **11**: 495–505.
- Gillgrass A, Cardiff RD, Sharan N, Kannan S, Muller WJ. 2003. Epidermal growth factor-dependent activation of Gab1 is involved in ErbB-2-mediated mammary tumor progression. *Oncogene* **11**: 9151–9155.
- Ginestier C, Liu S, Wicha MS. 2009. Getting to the root of BRCA1-deficient breast cancer. *Cell Stem Cell* **5**: 229–230.
- Guo W, Pylayeva Y, Pepe A, Yoshioka T, Muller WJ, Inghirami G, Giancotti FG. 2006. β4 integrin amplifies ErbB2 signaling to promote mammary tumorigenesis. *Cell* **126**: 489–502.

E.Y.H.P. Lee and W.J. Muller

- Gupta PB, Onder TT, Jiang G, Tao K, Kuperwasser C, Weinberg RA, Lander ES. 2009. Identification of selective inhibitors of cancer stem cells by high-throughput screening. *Cell* **138**: 645–659.
- Guy CT, Cardiff RD, Muller WJ. 1996. Activated neu induces rapid tumor progression. *J Biol Chem* **271**: 7673–7678.
- Guy CT, Muthuswamy SK, Cardiff RD, Soriano P, Muller WJ. 1994. Activation of the c-Src tyrosine kinase is required for the induction of mammary tumors in transgenic mice. *Genes Dev* **8**: 23–32.
- Guy CT, Webster MA, Schaller M, Parsons TJ, Cardiff RD, Muller WJ. 1992. Expression of the neu protooncogene in the mammary epithelium of transgenic mice induces metastatic disease. *Proc Natl Acad Sci* **89**: 10578–10582.
- Harvey DM, Levine AJ. 1991. p53 alteration is a common event in the spontaneous immortalization of primary BALB/c murine embryo fibroblasts. *Genes Dev* **5**: 2375–2385.
- Hirsch HA, Iliopoulos D, Tschlis PN, Struhl K. 2009. Metformin selectively targets cancer stem cells, and acts together with chemotherapy to block tumor growth and prolong remission. *Cancer Res* **69**: 7507–7511.
- Hodgson JG, Malek T, Bornstein S, Hariono S, Ginzinger DG, Muller WJ, Gray JW. 2005. Copy number aberrations in mouse breast tumors reveal loci and genes important in tumorigenic receptor tyrosine kinase signaling. *Cancer Res* **65**: 9695–9704.
- Holstege H, Joosse SA, van Oostrom CT, Nederlof PM, de Vries A, Jonkers J. 2009. High incidence of protein-truncating TP53 mutations in BRCA1-related breast cancer. *Cancer Res* **69**: 3625–3633.
- Huber LJ, Yang TW, Sarkisian CJ, Master SR, Deng CX, Chodosh LA. 2001. Impaired DNA damage response in cells expressing an exon 11-deleted murine Brca1 variant that localizes to nuclear foci. *Mol Cell Biol* **21**: 4005–4015.
- Hutchinson JN, Jin J, Cardiff RD, Woodgett JR, Muller WJ. 2004. Activation of Akt-1 (PKB- α) can accelerate ErbB-2-mediated mammary tumorigenesis but suppresses tumor invasion. *Cancer Res* **64**: 3171–3178.
- Hwang-Versluis WW, Kuo WH, Chang PH, Pan CC, Wang HH, Tsai ST, Jeng YM, Shew JY, Kung JT, Chen CH, et al. 2009. Multiple lineages of human breast cancer stem/progenitor cells identified by profiling with stem cell markers. *PLoS One* **4**: e8377.
- Hynes NE, Stern DF. 1994. The biology of erbB-2/neu/HER-2 and its role in cancer. *Biochim Biophys Acta* **1198**: 165–184.
- Iliopoulos D, Polyarchou C, Hatzia Apostolou M, Kottakis F, Maroulakou IG, Struhl K, Tschlis PN. 2009. MicroRNAs differentially regulated by Akt isoforms control EMT and stem cell renewal in cancer cells. *Sci Signal* **2**: ra62.
- Johnson TM, Hammond EM, Giaccia A, Attardi LD. 2005. The p53QS transactivation-deficient mutant shows stress-specific apoptotic activity and induces embryonic lethality. *Nat Genet* **37**: 145–152.
- Jones SN, Roe AE, Donehower LA, Bradley A. 1995. Rescue of embryonic lethality in Mdm2-deficient mice by absence of p53. *Nature* **378**: 206–208.
- Jonkers J, Meuwissen R, van der Gulden H, Peterse H, van der Valk M, Berns A. 2001. Synergistic tumor suppressor activity of BRCA2 and p53 in a conditional mouse model for breast cancer. *Nat Genet* **29**: 418–425.
- Ju X, Katiyar S, Wang C, Liu M, Jiao X, Li S, Zhou J, Turner J, Lisanti MP, Russell RG, et al. 2007. Akt1 governs breast cancer progression in vivo. *Proc Natl Acad Sci* **104**: 7438–7443.
- Kawamura T, Suzuki J, Wang YV, Menendez S, Morera LB, Raya A, Wahl GM, Belmonte JC. 2009. Linking the p53 tumour suppressor pathway to somatic cell reprogramming. *Nature* **460**: 1140–1144.
- Kim SS, Cao L, Baek HJ, Lim SC, Li C, Wang RH, Xu X, Cho KH, Deng CX. 2009. Impaired skin and mammary gland development and increased γ -irradiation-induced tumorigenesis in mice carrying a mutation of S1152-ATM phosphorylation site in Brca1. *Cancer Res* **69**: 9291–9300.
- Kim H, Chan R, Dankort DL, Zuo D, Najoukas M, Park M, Muller WJ. 2005. The c-Src tyrosine kinase associates with the catalytic domain of ErbB-2: Implications for ErbB-2 mediated signaling and transformation. *Oncogene* **24**: 7599–7607.
- Kruse JB, Gu W. 2009. Modes of p53 regulation. *Cell* **137**: 609–622.
- Kwong KY, Hung MC. 1998. A novel splice variant of HER2 with increased transformation activity. *Mol Carcinog* **23**: 62–68.
- Lahlou H, Sanguin-Gendreau V, Zuo D, Cardiff RD, McLean GW, Frame MC, Muller WJ. 2007. Mammary epithelial-specific disruption of the focal adhesion kinase blocks mammary tumor progression. *Proc Natl Acad Sci* **104**: 20302–20307.
- Lander AD. 2009. The ‘stem cell’ concept: Is it holding us back? *J Biol* **8**: 70.
- Landis MW, Pawlyk BS, Li T, Sicinski P, Hinds PW. 2006. Cyclin D1-dependent kinase activity in murine development and mammary tumorigenesis. *Cancer Cell* **9**: 13–22.
- Lane DP. 1992. Cancer, p53, guardian of the genome. *Nature* **358**: 15–16.
- Lee EY. 2008. Promotion of BRCA1-associated triple-negative breast cancer by ovarian hormones. *Curr Opin Obstet Gynecol* **20**: 68–73.
- Lee EY, To H, Shew JY, Bookstein R, Scully P, Lee WH. 1988. Inactivation of the retinoblastoma susceptibility gene in human breast cancers. *Science* **241**: 218–221.
- Lemoine NR, Staddon S, Dickson C, Barnes DM, Gullick WJ. 1990. Absence of activating transmembrane mutations in the c-erbB-2 proto-oncogene in human breast cancer. *Oncogene* **5**: 237–239.
- Lenferink AE, Simpson JF, Shawver LK, Coffey RJ, Forbes JT, Arteaga CL. 2000. Blockade of the epidermal growth factor receptor tyrosine kinase suppresses tumorigenesis in MMTV/Neu + MMTV/TGF- α bigenic mice. *Proc Natl Acad Sci* **97**: 9609–9614.
- Levine AJ, Oren M. 2009. The first 30 years of p53: Growing ever more complex. *Nat Rev Cancer* **9**: 749–758.
- Li W, Xiao C, Vonderhaar BK, Deng CX. 2007. A role of estrogen/ER α signaling in BRCA1-associated tissue-specific tumor formation. *Oncogene* **26**: 7204–7212.
- Lim E, Vaillant F, Wu D, Forrest NC, Pal B, Hart AH, Asselin-Labat ML, Gyorki DE, Ward T, Partanen A, et al. 2009. Aberrant luminal progenitors as the candidate target

- population for basal tumor development in BRCA1 mutation carriers. *Nat Med* **15**: 907–913.
- Lin SC, Lee KF, Nikitin AY, Hilsenbeck SG, Cardiff RD, Li A, Kang KW, Frank SA, Lee WH, Lee EY. 2004. Somatic mutation of p53 leads to estrogen receptor α -positive and negative mouse mammary tumors with high frequency of metastasis. *Cancer Res* **64**: 3525–3532.
- Liu S, Ginestier C, Charafe-Jauffret E, Foco H, Kleer CG, Merajver SD, Dontu G, Wicha MS. 2008. BRCA1 regulates human mammary stem/progenitor cell fate. *Proc Natl Acad Sci* **105**: 1680–1685.
- Liu X, Holstege H, van der Gulden H, Treur-Mulder M, Zevenhoven J, Velds A, Kerkhoven RM, van Vliet MH, Wessels LF, Peterse JL, et al. 2007. Somatic loss of BRCA1 and p53 in mice induces mammary tumors with features of human BRCA1-mutated basal-like breast cancer. *Proc Natl Acad Sci* **104**: 12111–12116.
- Lucchini F, Sacco MG, Hu N, Villa A, Brown J, Cesano L, Mangiarini L, Rindi G, Kindl S, Sessa F, et al. 1992. Early and multifocal tumors in breast, salivary, hardierian and epididymal tissues developed in MMTV-Neu transgenic mice. *Cancer Lett* **64**: 203–209.
- Ludwig T, Fisher P, Murty V, Efstratiadis A. 2001. Development of mammary adenocarcinomas by tissue-specific knockout of Brca2 in mice. *Oncogene* **20**: 3937–3948.
- Luo M, Fan H, Nagy T, Wei H, Wang C, Liu S, Wicha MS, Guan JL. 2009. Mammary epithelial-specific ablation of the focal adhesion kinase suppresses mammary tumorigenesis by affecting mammary cancer stem/progenitor cells. *Cancer Res* **69**: 466–474.
- Lynch HT, Marcus JN, Rubinstein WS. 2008. Stemming the tide of cancer for BRCA1/2 mutation carriers. *J Clin Oncol* **26**: 4239–4243.
- Manke IA, Lowery DM, Nguyen A, Yaffe MB. 2003. BRCT repeats as phosphopeptide-binding modules involved in protein targeting. *Science* **302**: 636–639.
- Mansour EG, Ravdin PM, Dressler L. 1994. Prognostic factors in early breast carcinoma. *Cancer* **74**: 381–400.
- Marcotte R, Zhou L, Kim H, Roskelly CD, Muller WJ. 2009. c-Src associates with ErbB2 through an interaction between catalytic domains and confers enhanced transforming potential. *Mol Cell Biol* **29**: 5858–5871.
- Mariotti A, Kedeshian PA, Dans M, Curatola AM, Gagnoux-Palacios L, Giancotti FG. 2001. EGF-R signaling through Fyn kinase disrupts the function of integrin $\alpha 6 \beta 4$ at hemidesmosomes: Role in epithelial cell migration and carcinoma invasion. *J Cell Biol* **155**: 447–458.
- Maroulakou IG, Oemler W, Naber SP, Tschlis PN. 2007. Akt1 ablation inhibits, whereas Akt2 ablation accelerates, the development of mammary adenocarcinomas in mouse mammary tumor virus (MMTV)-ErbB2/neu and MMTV-polyoma middle T transgenic mice. *Cancer Res* **67**: 167–177.
- Matsui Y, Halter SA, Holt JT, Hogan BL, Coffey RJ. 1990. Development of mammary hyperplasia and neoplasia in MMTV-TGF α transgenic mice. *Cell* **61**: 1147–1155.
- McCarthy A, Savage K, Gabriel A, Naceur C, Reis-Filho JS, Ashworth A. 2007. A mouse model of basal-like breast carcinoma with metaplastic elements. *J Pathol* **211**: 389–398.
- Menendez D, Inga A, Resnick MA. 2009. The expanding universe of p53 targets. *Nat Rev Cancer* **9**: 724–737.
- Miki Y, Swensen J, Shattuck-Eidens D, Futreal PA, Harshman K, Tavtigian S, Liu Q, Cochran C, Bennett LM, Ding W, et al. 1994. A strong candidate for the breast and ovarian cancer susceptibility gene BRCA1. *Science* **266**: 66–71.
- Montagna C, Andrechek ER, Padilla-Nash H, Muller WJ, Ried T. 2002. Centrosome abnormalities, recurring deletions of chromosome 4, and genomic amplification of HER2/neu define mouse mammary gland adenocarcinomas induced by mutant HER2/neu. *Oncogene* **21**: 890–898.
- Montes de Oca Luna R, Wagner DS, Lozano G. 1995. Rescue of early embryonic lethality in mdm2-deficient mice by deletion of p53. *Nature* **378**: 203–206.
- Moody SE, Perez D, Pan TC, Sarkisian CJ, Portocarrero CP, Sterner CJ, Notorfrancesco KL, Cardiff RD, Chodosh LA. 2005. The transcriptional repressor Snail promotes mammary tumor recurrence. *Cancer Cell* **8**: 197–209.
- Moody SE, Sarkisian CJ, Hahn KT, Gunther EJ, Pickup S, Dugan KD, Innocent N, Cardiff RD, Schnall MD, Chodosh LA. 2002. Conditional activation of Neu in the mammary epithelium of transgenic mice results in reversible pulmonary metastasis. *Cancer Cell* **2**: 451–461.
- Moro L, Venturino M, Bozzo C, Silengo L, Altruda F, Beguinot L, Tarone G, Defilippi P. 1998. Integrins induce activation of EGF receptor: Role in MAP kinase induction and adhesion-dependent cell survival. *Embo J* **17**: 6622–6632.
- Moynahan ME, Chiu JW, Koller BH, Jasin M. 1999. Brca1 controls homology-directed DNA repair. *Mol Cell* **4**: 511–518.
- Muller WJ, Arteaga CL, Muthuswamy SK, Siegel PM, Webster MA, Cardiff RD, Meise KS, Li F, Halter SA, Coffey RJ. 1996. Synergistic interaction of the Neu proto-oncogene product and transforming growth factor α in the mammary epithelium of transgenic mice. *Mol Cell Biol* **16**: 5726–5736.
- Muller WJ, Sinn E, Pattengale PK, Wallace R, Leder P. 1988. Single-step induction of mammary adenocarcinoma in transgenic mice bearing the activated c-neu oncogene. *Cell* **54**: 105–115.
- Muraoka-Cook RS, Kurokawa H, Koh Y, Forbes JT, Roebuck LR, Barcellos-Hoff MH, Moody SE, Chodosh LA, Arteaga CL. 2004. Conditional overexpression of active transforming growth factor $\beta 1$ in vivo accelerates metastases of transgenic mammary tumors. *Cancer Res* **64**: 9002–9011.
- Muraoka RS, Koh Y, Roebuck LR, Sanders ME, Brantley-Sieders D, Gorska AE, Moses HL, Arteaga CL. 2003. Increased malignancy of Neu-induced mammary tumors overexpressing active transforming growth factor $\beta 1$. *Mol Cell Biol* **23**: 8691–8703.
- Muraoka RS, Lenferink AE, Law B, Hamilton E, Brantley DM, Roebuck LR, Arteaga CL. 2002. ErbB2/Neu-induced, cyclin D1-dependent transformation is accelerated in p27-haploinsufficient mammary epithelial cells but impaired in p27-null cells. *Mol Cell Biol* **22**: 2204–2219.
- Muthuswamy SK, Muller WJ. 1995a. Activation of Src family kinases in Neu-induced mammary tumors correlates with their association with distinct sets of tyrosine phosphorylated proteins in vivo. *Oncogene* **11**: 1801–1810.
- Muthuswamy SK, Muller WJ. 1995b. Direct and specific interaction of c-Src with Neu is involved in signaling

E.Y.H.P. Lee and W.J. Muller

- by the epidermal growth factor receptor. *Oncogene* **11**: 271–279.
- Muthuswamy SK, Siegel PM, Dankort DL, Webster MA, Muller WJ. 1994. Mammary tumors expressing the neu proto-oncogene possess elevated c-Src tyrosine kinase activity. *Mol Cell Biol* **14**: 735–743.
- Nowell PC. 1976. The clonal evolution of tumor cell populations. *Science* **194**: 23–28.
- Parant J, Chavez-Reyes A, Little NA, Yan W, Reinke V, Jochemsen AG, Lozano G. 2001. Rescue of embryonic lethality in Mdm4-null mice by loss of Trp53 suggests a nonoverlapping pathway with MDM2 to regulate p53. *Nat Genet* **29**: 92–95.
- Park SY, Gonen M, Kim HJ, Michor F, Polyak K. 2010. Cellular and genetic diversity in the progression of in situ human breast carcinomas to an invasive phenotype. *J Clin Invest* **120**: 636–644.
- Pece S, Tosoni D, Confalonieri S, Mazzarol G, Vecchi M, Ronzoni S, Bernard L, Viale G, Pelicci PG, Di Fiore PP. 2010. Biological and molecular heterogeneity of breast cancers correlates with their cancer stem cell content. *Cell* **140**: 62–73.
- Pegram MD, Pauletti G, Slamon DJ. 1998. HER-2/neu as a predictive marker of response to breast cancer therapy. *Breast Cancer Res Treat* **52**: 65–77.
- Poole AJ, Li Y, Kim Y, Lin SC, Lee WH, Lee EY. 2006. Prevention of Brca1-mediated mammary tumorigenesis in mice by a progesterone antagonist. *Science* **314**: 1467–1470.
- Provenzano PP, Inman DR, Eliceiri KW, Beggs HE, Keely PJ. 2008. Mammary epithelial-specific disruption of focal adhesion kinase retards tumor formation and metastasis in a transgenic mouse model of human breast cancer. *Am J Pathol* **173**: 1551–1565.
- Pylayeva Y, Gillen KM, Gerald W, Beggs HE, Reichardt LF, Giancotti FG. 2009. Ras- and PI3K-dependent breast tumorigenesis in mice and humans requires focal adhesion kinase signaling. *J Clin Invest* **119**: 252–266.
- Ranger JJ, Levy DE, Shahalizadeh S, Hallet M, Muller WJ. 2009. Identification of a Stat3-dependent transcription regulatory network involved in metastatic progression. *Cancer Res* **69**: 6823–6830.
- Ravdin PM, Chamness GC. 1995. The c-erbB-2 proto-oncogene as a prognostic and predictive marker in breast cancer: A paradigm for the development of other macromolecular markers—a review. *Gene* **159**: 19–27.
- Reim F, Dombrowski Y, Ritter C, Buttman M, Hausler S, Ossadnik M, Krockenberger M, Beier D, Beier CP, Dietl J, et al. 2009. Immunoselection of breast and ovarian cancer cells with trastuzumab and natural killer cells: Selective escape of CD44^{high}/CD24^{low}/HER2^{low} breast cancer stem cells. *Cancer Res* **69**: 8058–8066.
- Rodriguez MC, Songyang Z. 2008. BRCT domains: Phosphopeptide binding and signaling modules. *Front Biosci* **13**: 5905–5915.
- Rottenberg S, Jaspers JE, Kersbergen A, van der Burg E, Nygren AO, Zander SA, Derksen PW, de Bruin M, Zevenhoven J, Lau A, et al. 2008. High sensitivity of BRCA1-deficient mammary tumors to the PARP inhibitor AZD2281 alone and in combination with platinum drugs. *Proc Natl Acad Sci* **105**: 17079–17084.
- Saal LH, Gruvberger-Saal SK, Persson C, Lovgren K, Juntanen M, Staaf J, Jonsson G, Pires MM, Maurer M, Holm K, et al. 2008. Recurrent gross mutations of the PTEN tumor suppressor gene in breast cancers with deficient DSB repair. *Nat Genet* **40**: 102–107.
- Sakai W, Swisher EM, Karlan BY, Agarwal MK, Higgins J, Friedman C, Villegas E, Jacquemont C, Farrugia DJ, Couch FJ, et al. 2008. Secondary mutations as a mechanism of cisplatin resistance in BRCA2-mutated cancers. *Nature* **451**: 1116–1120.
- Sandgren EP, Luetke NC, Palmiter RD, Brinster RL, Lee DC. 1990. Overexpression of TGF α in transgenic mice: Induction of epithelial hyperplasia, pancreatic metaplasia, and carcinoma of the breast. *Cell* **61**: 1121–1135.
- Sandgren EP, Schroeder JA, Qui TH, Palmiter RD, Brinster RL, Lee DC. 1995. Inhibition of mammary gland involution is associated with transforming growth factor α but not c-myc-induced tumorigenesis in transgenic mice. *Cancer Res* **55**: 3915–3927.
- Schade B, Lam SH, Cernea D, Sanguin-Gendreau V, Cardiff RD, Jung BL, Hallett M, Muller WJ. 2007. Distinct ErbB-2 coupled signaling pathways promote mammary tumors with unique pathologic and transcriptional profiles. *Cancer Res* **67**: 7579–7588.
- Schade B, Rao T, Dourdin N, Lesurf R, Halle M, Cardiff RD, Muller WJ. 2009. PTEN deficiency in a luminal ErbB-2 mouse model results in dramatic acceleration of mammary tumorigenesis and metastasis. *J Biol Chem* **284**: 19018–19026.
- Shafee N, Smith CR, Wei S, Kim Y, Mills GB, Hortobagyi GN, Stanbridge EJ, Lee EY. 2008. Cancer stem cells contribute to cisplatin resistance in Brca1/p53-mediated mouse mammary tumors. *Cancer Res* **68**: 3243–3250.
- Shah SP, Morin RD, Khattri J, Prentice L, Pugh T, Burleigh A, Delaney A, Gelmon K, Guliany R, Senz J, et al. 2009. Mutational evolution in a lobular breast tumour profiled at single nucleotide resolution. *Nature* **461**: 809–813.
- Shangary S, Wang S. 2009. Small-molecule inhibitors of the MDM2-p53 protein–protein interaction to reactivate p53 function: A novel approach for cancer therapy. *Annu Rev Pharmacol Toxicol* **49**: 223–241.
- Shimono Y, Zabala M, Cho RW, Lobo N, Dalerba P, Qian D, Diehn M, Liu H, Panula SP, Chiao E, et al. 2009. Down-regulation of miRNA-200c links breast cancer stem cells with normal stem cells. *Cell* **138**: 592–603.
- Siegel PM, Muller WJ. 1996. Mutations affecting conserved cysteine residues within the extracellular domain of Neu promote receptor dimerization and activation. *Proc Natl Acad Sci* **93**: 8878–8883.
- Siegel PM, Dankort DL, Hardy WR, Muller WJ. 1994. Novel activating mutations in the neu proto-oncogene involved in induction of mammary tumors. *Mol Cell Biol* **14**: 7068–7077.
- Siegel PM, Ryan ED, Cardiff RD, Muller WJ. 1999. Elevated expression of activated forms of Neu/ErbB-2 and ErbB-3 are involved in the induction of mammary tumors in transgenic mice: Implications for human breast cancer. *Embo J* **18**: 2149–2164.
- Siegel PM, Shu W, Cardiff RD, Muller WJ, Massague J. 2003. Transforming growth factor $\{\beta\}$ signaling impairs

Oncogenes and Tumor Suppressor Genes

- Neu-induced mammary tumorigenesis while promoting pulmonary metastasis. *Proc Natl Acad Sci* **100**: 8430–8435.
- Slamon DJ, Clark GM, Wong SG, Levin WJ, Ullrich A, McGuire WL. 1987. Human breast cancer: Correlation of relapse and survival with the amplification of the HER2/neu oncogene. *Science* **235**: 177–182.
- Slamon DJ, Godolphin W, Jones LA, Holt JA, Wong SG, Keith DE, Levin WJ, Stuart SG, Udove J, Ullrich A et al. 1989. Studies of the HER-2/neu proto-oncogene in human breast and ovarian cancer. *Science* **244**: 707–712.
- Song H, Hollstein M, Xu Y. 2007. p53 gain-of-function cancer mutants induce genetic instability by inactivating ATM. *Nat Cell Biol* **9**: 573–580.
- Stephens PJ, McBride DJ, Lin ML, Varela I, Pleasance ED, Simpson JT, Stebbings LA, Leroy C, Edkins S, Mudie LJ, et al. 2009. Complex landscapes of somatic rearrangement in human breast cancer genomes. *Nature* **462**: 1005–1010.
- Sur S, Pagliarini R, Bunz F, Rago C, Diaz LA Jr, Kinzler KW, Vogelstein B, Papadopoulos N. 2009. A panel of isogenic human cancer cells suggests a therapeutic approach for cancers with inactivated p53. *Proc Natl Acad Sci* **106**: 3964–3969.
- Suzuki HI, Yamagata K, Sugimoto K, Iwamoto T, Kato S, Miyazono K. 2009. Modulation of microRNA processing by p53. *Nature* **460**: 529–533.
- Sy SM, Huen MS, Chen J. 2009. PALB2 is an integral component of the BRCA complex required for homologous recombination repair. *Proc Natl Acad Sci* **106**: 7155–7160.
- T'Ang A, Varley JM, Chakraborty S, Murphree AL, Fung YK. 1988. Structural rearrangement of the retinoblastoma gene in human breast carcinoma. *Science* **242**: 263–266.
- Thorslund T, West SC. 2007. BRCA2: A universal recombinase regulator. *Oncogene* **26**: 7720–7730.
- Tung N, Wang Y, Collins LC, Kaplan J, Li H, Gelman R, Comander AH, Gallagher B, Fettes K, Krag K, et al. 2010. Estrogen receptor positive breast cancers in BRCA1 mutation carriers: Clinical risk factors and pathologic features. *Breast Cancer Res* **12**: R12.
- Urano T, Saito T, Tsukui T, Fujita M, Hosoi T, Muramatsu M, Ouchi Y, Inoue S. 2002. Efp targets 14-3-3 σ for proteolysis and promotes breast tumour growth. *Nature* **417**: 871–875.
- Ursini-Siegel J, Hardy WR, Zuo D, Lam SH, Sanguin-Gendreau V, Cardiff RD, Pawson T, Muller WJ. 2008. ShcA signalling is essential for tumour progression in mouse models of human breast cancer. *Embo J* **27**: 910–920.
- van Haaften G, Dalgleish GL, Davies H, Chen L, Bignell G, Greenman C, Edkins S, Hardy C, O'Meara S, Teague J, et al. 2009. Somatic mutations of the histone H3K27 demethylase gene *UTX* in human cancer. *Nat Genet* **41**: 521–523.
- Vercoutter-Edouart AS, Lemoine J, Le Bourhis X, Louis H, Boilly B, Nurcombe V, Revillion F, Peyrat JB, Hondermarck H. 2001. Proteomic analysis reveals that 14-3-3 σ is down-regulated in human breast cancer cells. *Cancer Res* **61**: 76–80.
- Visvader JE. 2009. Keeping abreast of the mammary epithelial hierarchy and breast tumorigenesis. *Genes Dev* **23**: 2563–2577.
- Walsh T, King MC. 2007. Ten genes for inherited breast cancer. *Cancer Cell* **11**: 103–105.
- Wang F, Hansen RK, Radisky D, Yoneda T, Barcellos-Hoff MH, Petersen OW, Turley EA, Bissell MJ. 2002. Phenotypic reversion or death of cancer cells by altering signaling pathways in three-dimensional contexts. *J Natl Cancer Inst* **94**: 1494–1503.
- Wang SE, Xiang B, Zent R, Quaranta V, Pozzi A, Arteaga CL. 2009a. Transforming growth factor β induces clustering of HER2 and integrins by activating Src-focal adhesion kinase and receptor association to the cytoskeleton. *Cancer Res* **69**: 475–482.
- Wang Y, Yang J, Zheng H, Tomasek GJ, Zhang P, McKeever PE, Lee EY, Zhu Y. 2009b. Expression of mutant p53 proteins implicates a lineage relationship between neural stem cells and malignant astrocytic glioma in a murine model. *Cancer Cell* **15**: 514–526.
- Weaver VM, Petersen OW, Wang F, Larabell CA, Briand P, Damsky C, Bissell MJ. 1997. Reversion of the malignant phenotype of human breast cells in three-dimensional culture and in vivo by integrin blocking antibodies. *J Cell Biol* **137**: 231–245.
- Wertheim GB, Yang TW, Pan TC, Ramne A, Liu Z, Gardner HP, Dugan KD, Kristel P, Kreike B, van de Vijver MJ, et al. 2009. The Snf1-related kinase, Hunk, is essential for mammary tumor metastasis. *Proc Natl Acad Sci* **106**: 15855–15860.
- White DE, Rayment JH, Muller WJ. 2006. Addressing the role of cell adhesion in tumor cell dormancy. *Cell Cycle* **5**: 1756–1759.
- White DE, Kurpios NA, Zuo D, Hassell JA, Blaess S, Mueller U, Muller WJ. 2004. Targeted disruption of $\beta 1$ -integrin in a transgenic mouse model of human breast cancer reveals an essential role in mammary tumor induction. *Cancer Cell* **6**: 159–170.
- Wong AK, Pero R, Ormonde PA, Tavtigian SV, Bartel PL. 1997. RAD51 interacts with the evolutionarily conserved BRC motifs in the human breast cancer susceptibility gene *brca2*. *J Biol Chem* **272**: 31941–31944.
- Wooster R, Bignell G, Lancaster J, Swift S, Seal S, Mangion J, Collins N, Gregory S, Gumbs C, Micklem G. 1995. Identification of the breast cancer susceptibility gene BRCA2. *Nature* **378**: 789–792.
- Wu-Baer F, Lagazon K, Yuan W, Baer R. 2003. The BRCA1/BARD1 heterodimer assembles polyubiquitin chains through an unconventional linkage involving lysine residue K6 of ubiquitin. *J Biol Chem* **278**: 34743–34746.
- Xu Y. 2008. Induction of genetic instability by gain-of-function p53 cancer mutants. *Oncogene* **27**: 3501–3507.
- Xu X, Wagner KU, Larson D, Weaver Z, Li C, Ried T, Hennighausen L, Wynshaw-Boris A, Deng CX. 1999. Conditional mutation of *Brcal* in mammary epithelial cells results in blunted ductal morphogenesis and tumour formation [see comments]. *Nat Genet* **22**: 37–43.
- Yang H, Jeffrey PD, Miller J, Kinnucan E, Sun Y, Thoma NH, Zheng N, Chen PL, Lee WH, Pavletich NP. 2002. BRCA2

E.Y.H.P. Lee and W.J. Muller

- function in DNA binding and recombination from a BRCA2-DSS1-ssDNA structure. *Science* **297**: 1837–1848.
- Yu X, Chini CC, He M, Mer G, Chen J. 2003. The BRCT domain is a phospho-protein binding domain. *Science* **302**: 639–642.
- Yu Q, Geng Y, Sicinski P. 2001. Specific protection against breast cancers by cyclin D1 ablation. *Nature* **411**: 1017–1021.
- Yu Q, Sicinska E, Geng Y, Ahnstrom M, Zagozdzon A, Kong Y, Gardner H, Kiyokawa H, Harris LN, Sta O, et al. 2006. Requirement for CDK4 kinase function in breast cancer. *Cancer Cell* **9**: 23–32.
- Yuan SS, Lee SY, Chen G, Song M, Tomlinson GE, Lee EY. 1999. BRCA2 is required for ionizing radiation-induced assembly of Rad51 complex in vivo. *Cancer Res* **59**: 3547–3551.
- Zhang F, Ma J, Wu J, Ye L, Cai H, Xia B, Yu X. 2009. PALB2 links BRCA1 and BRCA2 in the DNA-damage response. *Curr Biol* **19**: 524–529.
- Zheng L, Pan H, Li S, Flesken-Nikitin A, Chen PL, Boyer TG, Lee WH. 2000. Sequence-specific transcriptional corepressor function for BRCA1 through a novel zinc finger protein, ZBRK1. *Mol Cell* **6**: 757–768.
- Zhong Q, Chen CF, Li S, Chen Y, Wang CC, Xiao J, Chen PL, Sharp ZD, Lee WH. 1999. Association of BRCA1 with the hRad50-hMre11-p95 complex and the DNA damage response. *Science* **285**: 747–750.
- Zilfou JT, Lowe SW. 2009. Tumor suppressive functions of p53. *Cold Spring Harb Perspect Biol* **1**: a001883.

Cold Spring Harbor Perspectives in Biology

Oncogenes and Tumor Suppressor Genes

Eva Y.H.P. Lee and William J. Muller

Cold Spring Harb Perspect Biol 2010; doi: 10.1101/cshperspect.a003236 originally published online August 18, 2010

Subject Collection [The Mammary Gland as an Experimental Model](#)

On the Role of the Microenvironment in Mammary Gland Development and Cancer

Derek Radisky

On Using Functional Genetics to Understand Breast Cancer Biology

Kornelia Polyak

On Oncogenes and Tumor Suppressor Genes in the Mammary Gland

Rushika M. Perera and Nabeel Bardeesy

On Leukocytes in Mammary Development and Cancer

Cyrus M. Ghajar

On Chromatin Remodeling in Mammary Gland Differentiation and Breast Tumorigenesis

Kornelia Polyak

On Hormone Action in the Mammary Gland

J.M. Rosen

TGF- β Biology in Mammary Development and Breast Cancer

Harold Moses and Mary Helen Barcellos-Hoff

A Compendium of the Mouse Mammary Tumor Biologist: From the Initial Observations in the House Mouse to the Development of Genetically Engineered Mice

Robert D. Cardiff and Nicholas Kenney

On How Mammary Gland Reprogramming Metalloproteinases Couple Form with Function

Bonnie F. Sloane

On Molecular Mechanisms Guiding Embryonic Mammary Gland Development

Gertraud W. Robinson

On Stem Cells in the Human Breast

Mark A. LaBarge

On Murine Mammary Epithelial Stem Cells: Discovery, Function, and Current Status

Jeffrey M. Rosen

On In Vivo Imaging in Cancer

David Piwnicka-Worms

Choosing a Mouse Model: Experimental Biology in Context—The Utility and Limitations of Mouse Models of Breast Cancer

Alexander D. Borowsky

Mammary Gland ECM Remodeling, Stiffness, and Mechanosignaling in Normal Development and Tumor Progression

Pepper Schedin and Patricia J. Keely

Molecular Mechanisms Guiding Embryonic Mammary Gland Development

Pamela Cowin and John Wysolmerski

For additional articles in this collection, see <http://cshperspectives.cshlp.org/cgi/collection/>

**All Modifications and
Oligo Types Synthesized**

Long Oligos • Fluorescent • Chimeric • DNA • RNA • Antisense

Oligo Modifications?
Your wish is our command.

For additional articles in this collection, see <http://cshperspectives.cshlp.org/cgi/collection/>

**All Modifications and
Oligo Types Synthesized**
Long Oligos • Fluorescent • Chimeric • DNA • RNA • Antisense

Oligo Modifications?
Your wish is our command.