IP₃ Receptors: Toward Understanding Their Activation

Colin W. Taylor and Stephen C. Tovey

Department of Pharmacology, University of Cambridge, Tennis Court Road, Cambridge, CB2 1PD, United Kingdom

Correspondence: cwt1000@cam.ac.uk

Inositol 1,4,5-trisphosphate receptors (IP₃R) and their relatives, ryanodine receptors, are the channels that most often mediate Ca²⁺ release from intracellular stores. Their regulation by Ca²⁺ allows them also to propagate cytosolic Ca²⁺ signals regeneratively. This brief review addresses the structural basis of IP₃R activation by IP₃ and Ca²⁺. IP₃ initiates IP₃R activation by promoting Ca²⁺ binding to a stimulatory Ca²⁺-binding site, the identity of which is unresolved. We suggest that interactions of critical phosphate groups in IP₃ with opposite sides of the clam-like IP₃-binding core cause it to close and propagate a conformational change toward the pore via the adjacent N-terminal suppressor domain. The pore, assembled from the last pair of transmembrane domains and the intervening pore loop from each of the four IP₃R subunits, forms a structure in which a luminal selectivity filter and a gate at the cytosolic end of the pore control cation fluxes through the IP₃R.

A BRIEF HISTORY OF IP₃ RECEPTORS

Sidney Ringer, in his famous correction to an earlier paper, showed that Ca²⁺ entry can evoke a physiological response by demonstrating that beating of the frog heart requires extracellular Ca²⁺ (Ringer 1883). Almost a century passed before it became clear that this Ca²⁺ entry, via voltage-gated Ca²⁺ channels, was not directly responsible for contraction, but instead provided the trigger for a much larger release of Ca²⁺ from stores within the sarcoplasmic reticulum (SR). The latter is mediated by type-2 ryanodine receptors (RyR) (Fabiato 1983; Cheng et al. 1993), which like many Ca²⁺ channels, are able both to transport Ca²⁺ through an open pore and respond to it. These observations highlight

two general points. First, cells call upon two sources of Ca²⁺ to evoke increases in cytosolic Ca²⁺ concentration; second, interactions between these Ca²⁺ fluxes across the plasma membrane and the membranes of intracellular stores are important determinants of the physiological response. The same points apply to the Ca²⁺ signals evoked by receptors that stimulate phospholipase C (PLC) and, thereby, formation of inositol 1,4,5-trisphosphate (IP₃).

The biochemical sequence linking these receptors to formation of IP₃ emerged in the 1980s (Michell et al. 1989; Berridge 2005), but work in the decade before had established that many receptors regulate many different responses by increasing the cytosolic Ca²⁺ concentration (Rasmussen 1970; Berridge 1975). In his

Editors: Martin D. Bootman, Michael J. Berridge, James W. Putney, and H. Llewelyn Roderick
Additional Perspectives on Calcium Signaling available at www.cshperspectives.org

Copyright © 2010 Cold Spring Harbor Laboratory Press; all rights reserved; doi: 10.1101/cshperspect.a004010
Cite this article as *Cold Spring Harb Perspect Biol* 2010;2:a004010

C.W. Taylor and S.C. Tovey

influential review, Bob Michell (Michell 1975), building on work showing that many of these receptors also stimulate phospholipid turnover (Hokin and Hokin 1953), had suggested a causal link between phosphoinositide hydrolysis and Ca^{2+} signals. Here, as in many studies, the emphasis was on Ca^{2+} entry, with a consensus only slowly emerging that Ca^{2+} fluxes across both the plasma membrane and the membranes of intracellular stores contribute to cytosolic Ca^{2+} signals (Rasmussen 1970; Berridge 1975; Williams 1980; Putney et al. 1981). In the years following Michell's review, decisive evidence, much of it coming from Mike Berridge's elegant studies of blowfly salivary gland, established that phosphoinositide hydrolysis is, as predicted by Michell, required for PLC-linked receptors to evoke Ca^{2+} signals (Berridge and Fain 1979). The same preparation was used to show that IP_3 is the first water-soluble product of the signaling pathway (Berridge 1983). IP_3 , thus, emerged as a prime candidate for the cytosolic messenger linking events at the plasma membrane to release of Ca^{2+} from intracellular stores. Paradoxically, it was to be many years before the links between receptors that stimulate PLC and Ca^{2+} entry were resolved. These came with elaboration of the pathways linking empty Ca^{2+} stores to Ca^{2+} entry, the so-called store-operated Ca^{2+} entry pathway (Putney 1997; Park et al. 2009), and recognition that many trp channels are regulated by products of PLC activity (Nilius et al. 2007). IP_3 receptors (IP_3R) also contribute more directly to Ca^{2+} entry across the plasma membrane either because, at least in some cells, IP_3R are functionally expressed in the plasma membrane (Dellis et al. 2006; Dellis et al. 2008), or perhaps through their direct interactions with other plasma membrane Ca^{2+} channels (Kiselyov et al. 1999). Here, we focus solely on Ca^{2+} release from the endoplasmic reticulum (ER) by IP_3R . Some of the key steps in the evolution of our current understanding of IP_3R are listed in Table 1.

The role of the SR as the intracellular source of Ca^{2+} signals in striated muscle was long-established (Endo et al. 1970), but there was no such agreement on the identity of the organelle from which Ca^{2+} was released in other cells.

Competing claims suggested roles for mitochondria or the ER. Evidence that in resting hepatocytes only the ER contains appreciable amounts of Ca^{2+} (Burgess et al. 1983) was quickly followed by the demonstration that IP_3 evoked Ca^{2+} release from a non-mitochondrial Ca^{2+} store in permeabilized pancreatic acinar cells (Streb et al. 1983). Countless groups quickly replicated these findings in many cells, and within months it was universally accepted that the ER is the major Ca^{2+} store from which IP_3 stimulates Ca^{2+} release in most animal cells (Berridge and Irvine 1984; Berridge and Irvine 1989). Subsequent work has suggested that IP_3 may also stimulate Ca^{2+} release from the Golgi apparatus (Pinton et al. 1998), from within the nucleus (Gerasimenko et al. 1995; Echevarria et al. 2003; Marchenko et al. 2005), and perhaps also from secretory vesicles (Gerasimenko et al. 1996), but ER remains the major IP_3 -sensitive Ca^{2+} store. Evidence that IP_3 stimulates Ca^{2+} efflux from the ER (rather than inhibiting Ca^{2+} uptake) and the first single channel recordings (Ehrlich and Watras 1988) established that the IP_3R is an IP_3 -gated, Ca^{2+} -permeable channel. The first studies of ^{32}P - IP_3 binding (Spät et al. 1986) were followed by purification of IP_3R from cerebellum (Maeda et al. 1988; Supattapone et al. 1988) and then cloning of the first IP_3R subtype ($\text{IP}_3\text{R1}$) (Furuichi et al. 1989; Mignery et al. 1989). Subsequent studies identified two additional genes encoding vertebrate IP_3R ($\text{IP}_3\text{R2}$ and $\text{IP}_3\text{R3}$) and a single gene in invertebrates (Taylor et al. 1999). It remains far from clear whether plants express related IP_3R (Krinke et al. 2007). These studies established that IP_3R are unusually large proteins, comprising tetramers of closely-related subunits, each with about 2700 amino acid residues. RyR are even larger: they, too, are tetramers, but the subunits are almost twice the size of IP_3R (~ 5000 residues). This progress with identifying IP_3R together with single channel recordings of IP_3R , initially in artificial lipid bilayers and later in native membranes (Foskett et al. 2007; Rahman et al. 2009), provided the foundations from which to explore the structural determinants of IP_3R behavior. The advances toward understanding the molecular mechanisms

Table 1. Landmarks en route to a structural analysis of IP₃ receptor behavior.

	RyR	IP ₃ R
1883	Ca ²⁺ entry required for heart contraction. ¹	
1953		Acetylcholine stimulates turnover of phospholipids. ²
1975		Phosphoinositide hydrolysis proposed to cause Ca ²⁺ signals. ³
1977	Ca ²⁺ waves occur at fertilization. ⁴	
1977	Ca ²⁺ -induced Ca ²⁺ release in SR. ⁵	
1979		Phosphoinositide hydrolysis required for receptor-stimulated Ca ²⁺ signals. ⁶
1980	Introduction of Quin 2 ⁷ and facile loading methods. ⁸	
1983		IP ₃ is first water-soluble product of PLC. ⁹
1983		IP ₃ stimulates Ca ²⁺ release from a non-mitochondrial store. ¹⁰
1985	Ryanodine, selective RyR ligand. ¹¹	
1985	Single channel records of RyR. ¹²	
1986		Frequency-coded Ca ²⁺ spikes. ¹³
1987		Ca ²⁺ regulates IP ₃ R. ^{14,15}
1987	RyR1 purified. ¹⁶	IP ₃ R1 purified. ¹⁷
1988		Single channel records of IP ₃ R. ¹⁸
1989	Cloning of RyR1. ¹⁹	Cloning of IP ₃ R1. ^{20,21}
1990		Elementary Ca ²⁺ -release events. ²²
1993	Elementary Ca ²⁺ -release events. ²³	
2002		Atomic structure of IBC. ²⁴
2005		Atomic structure of SD. ²⁵
2009	Atomic structure of N-terminal of RyR. ^{26,27}	

¹Ringer (1883).²Hokin & Hokin (1953).³Michell (1975).⁴Ridgeway et al. (1977).⁵Endo (1977).⁶Berridge & Fain (1979).⁷Tsien (1980).⁸Tsien (1981).⁹Berridge (1983).¹⁰Streb et al. (1983).¹¹Sutko et al. (1985).¹²Smith et al. (1985).¹³Woods et al. (1986).¹⁴Iino (1987).¹⁵Iino (1990).¹⁶Imagawa et al. (1987).¹⁷Supattapone et al. (1988).¹⁸Ehrlich & Watras (1988).¹⁹Takeshima et al. (1989).²⁰Mignery et al. (1989).²¹Furuichi et al. (1989).²²Parker & Ivorra (1990).²³Cheng et al. (1993).²⁴Bosanac et al. (2002).²⁵Bosanac et al. (2005).²⁶Amador et al. (2009).²⁷Lobo & Van Petegem (2009).

C.W. Taylor and S.C. Tovey

of IP₃R behavior were accompanied by similar progress with RyR (Table 1). Recurrent themes, to which we return, are the similarities between RyR and IP₃R, and the many instances where observations of one channel family have informed further analysis of the other. Very recently, a third family of intracellular Ca²⁺ channels, unrelated to RyR and IP₃R, has been implicated in Ca²⁺ signaling. These are the two-pore channels (TPC) that are activated by NAADP and release Ca²⁺ from acidic Ca²⁺ stores, including lysosomes and endosomes (Patel et al. 2010; Zhu et al. 2010). Several trp (transient receptor protein) channels, in addition to their roles in the plasma membrane, may also mediate release of Ca²⁺ from intracellular stores (Gees et al. 2010).

Parallel to work addressing the workings of IP₃R, there was growing interest in the spatio-temporal complexity of cytosolic Ca²⁺ signals. Ca²⁺ waves were first observed during fertilization. These waves were proposed to result from Ca²⁺-induced Ca²⁺ release (CICR) and were followed by smaller repetitive Ca²⁺ transients (Ridgway et al. 1977; Gilkey 1983). It was, however, the work of Peter Cobbold that focused most attention on the complexity of intracellular Ca²⁺ signals (Woods et al. 1986). Just as the activity of a nerve is conveyed by the frequency of its action potentials, Cobbold demonstrated that in hepatocytes the concentration of the extracellular stimulus determined the frequency of the cytosolic Ca²⁺ transients. As these ideas gathered momentum (Berridge 1995), evidence accumulated in support of cells using the information provided by frequency-encoded Ca²⁺ spikes as an efficient means of regulating cellular activity (Dolmetsch et al. 1997; Li et al. 1998; Berridge et al. 2000; Dupont et al. 2003). The single greatest contributor to progress in understanding the genesis of these intracellular Ca²⁺ signals was the introduction, by Roger Tsien in 1980, of simple, minimally disruptive methods for measuring the free cytosolic Ca²⁺ concentration in intact cells (Tsien 1980; Tsien 1981). These methods, in combination with improved optical microscopy, allowed Ian Parker to begin to resolve the subcellular organization of IP₃-evoked Ca²⁺ signals (Parker and Ivorra 1990;

Parker et al. 1996). He showed that as the IP₃ concentration increases, it triggers a hierarchy of elementary Ca²⁺ release events, beginning with the openings of single IP₃R (Ca²⁺ blips), progressing to the coordinated openings of a cluster of several IP₃R (Ca²⁺ puffs) and finally, with sufficient IP₃, culminating in a regenerative Ca²⁺ wave invading the entire cell (Bootman et al. 1997; Demuro and Parker 2007). The demonstration, in 1987 by Masamitsu Iino, that IP₃R are stimulated by cytosolic Ca²⁺ (Iino 1987), and the later widespread recognition that all IP₃R are biphasically regulated by cytosolic Ca²⁺ (Iino 1990; Taylor and Laude 2002), provided what has become the most widely accepted explanation for the recruitment of elementary Ca²⁺-release events. Namely, that CICR, already an established feature of RyR (Endo et al. 1970), allows an active IP₃R to propagate its activity to neighboring IP₃R.

These observations and accumulating evidence that local Ca²⁺ signals can selectively regulate local events (Rizzuto et al. 1993; Berridge et al. 2000; Dyer et al. 2005; Willoughby and Cooper 2007) prompted a re-assessment of the ways in which Ca²⁺ signals convey information. It became untenable to think of responses to graded changes in the intensity of the extracellular stimulus as being simply encoded in graded changes in global cytosolic Ca²⁺ concentration. Ca²⁺ entering the cytosol via one channel can regulate different proteins to Ca²⁺ entering via another (Berridge et al. 2000; Dyer et al. 2005; Willoughby and Cooper 2007). Hence, the spatial organization of the changes in cytosolic Ca²⁺ concentration profoundly affects the physiological response, and that presents many opportunities for delivering different Ca²⁺ signals in response to different stimuli or different stimulus intensities. The duration of each Ca²⁺ increase, whether local or global, is also important in determining not only the amplitude of the response, but also its nature, because Ca²⁺-binding proteins differ in their responses to transient and sustained signals. Finally, the frequency with which Ca²⁺ signals are delivered can determine both the nature and amplitude of the cellular response. The key point is that the versatility of Ca²⁺ as an intracellular messenger

capable of regulating diverse cellular events depends largely on the spatiotemporal complexity of cytosolic Ca²⁺ signals (Berridge et al. 2000). If we are to understand how Ca²⁺ functions as a ubiquitous intracellular messenger, we must explain how IP₃-evoked Ca²⁺ signals grow from the opening of a single IP₃R to much larger events. That explanation depends, ultimately, on putting IP₃R into appropriate places within the cell, and on the interactions between IP₃ and Ca²⁺ in regulating the opening of IP₃R. In recent reviews (Taylor et al. 2009a; Taylor et al. 2009b) and original reports, we have described how IP₃R are co-translationally targeted to the ER and then retained there by sequences within their transmembrane domains (TMD) (Parker et al. 2004; Pantazaka and Taylor 2010). We have also suggested that within the ER, IP₃ causes IP₃R to assemble into small clusters within which their regulation by both IP₃ and Ca²⁺ is retuned to facilitate the Ca²⁺-mediated recruitment of IP₃R activity by an active neighbor (Rahman and Taylor 2009; Rahman et al. 2009). Here, we focus entirely on the interactions between Ca²⁺ and IP₃ in regulating IP₃R activity, and the extent to which we can explain those interactions at the structural level.

REGULATION OF IP₃ RECEPTORS BY Ca²⁺ AND IP₃

Activation of IP₃R requires both IP₃ and its permeating ion, Ca²⁺ (Finch et al. 1991; Marchant and Taylor 1997; Adkins and Taylor 1999; Taylor and Laude 2002; Foskett et al. 2007). There are reports of IP₃-independent activation of IP₃R by CaBP1 (Yang et al. 2002), a member of the neuronal Ca²⁺-sensor family, and by Gβγ subunits (Zeng et al. 2003), but the physiological relevance is unclear (Haynes et al. 2004; Nadif Kasri et al. 2004). The current consensus is that binding of IP₃ to the IP₃R is essential for its activation, but whether all four IP₃-binding sites of the tetrameric IP₃R must be occupied is unresolved. Positively cooperative responses to IP₃ in some (Dufour et al. 1997; Marchant and Taylor 1997; Tu et al. 2005a), though not all, studies (Finch et al. 1991; Watras et al. 1991;

Laude et al. 2005), and delays before the first response to IP₃ that decrease with increasing IP₃ concentration (Marchant and Taylor 1997), indicate that channel opening requires occupancy of more than one IP₃-binding site. However, gating by IP₃ of heteromeric IP₃R in which at least one subunit is mutated to prevent IP₃ binding suggests that occupancy of fewer than four IP₃-binding sites may be sufficient to cause some channel opening (Boehning and Joseph 2000a). IP₃R subtypes differ in their affinities for IP₃, with the general consensus being that IP₃R2 is more sensitive than IP₃R1, and both are considerably more sensitive than IP₃R3 (Tu et al. 2005b; Iwai et al. 2007). In the cellular context, however, differences in expression level (Dellis et al. 2006; Tovey et al. 2010), subcellular distribution (Petersen et al. 1999), post-transcriptional and post-translational modifications, and association of IP₃R with accessory proteins (Patterson et al. 2004) may be more important determinants of sensitivity.

Soon after the first report of IP₃-evoked Ca²⁺ release, cytosolic Ca²⁺ was shown also to regulate IP₃R (Suematsu et al. 1984; Jean and Klee 1986); thereafter, it emerged that the effects of Ca²⁺ were biphasic, with modest increases in cytosolic Ca²⁺ concentration enhancing responses to IP₃, while higher concentrations were inhibitory (Iino 1987; Iino 1990; Finch et al. 1991; Parys et al. 1992; Marshall and Taylor 1993). This provided yet another parallel with RyR, which are also biphasically regulated by Ca²⁺ (Hamilton 2005). The coregulation of IP₃R by IP₃ and Ca²⁺ in permeabilized cells was confirmed by single-channel recordings of IP₃R1 reconstituted into lipid bilayers (Bezprozvanny et al. 1991; Striggow and Ehrlich 1996; Kaftan et al. 1997; Ramos-Franco et al. 1998a; Ramos-Franco et al. 1998b; Tu et al. 2002; Tu et al. 2005b) and in native nuclear membranes (Stehno-Bittel et al. 1995; Mak et al. 1998; Boehning et al. 2001a; Marchenko et al. 2005). In each case, the single-channel open probability (*P*_o) of IP₃-activated channels displayed a bell-shaped dependence on cytosolic Ca²⁺ concentration. Evidence that purified IP₃R1 could be stimulated, but not inhibited, by cytosolic Ca²⁺ (Thrower et al. 1998; Michikawa et al.

C.W. Taylor and S.C. Tovey

1999) raised the possibility that Ca^{2+} inhibition might be mediated by an accessory protein, although it has yet to be identified. The same explanation perhaps accounts for some reports, often derived from bilayer recordings, in which Ca^{2+} was suggested not to inhibit $\text{IP}_3\text{R2}$ or $\text{IP}_3\text{R3}$ (Horne and Meyer 1995; Hagar et al. 1998; Miyakawa et al. 1999; Ramos-Franco et al. 2000). The balance of opinion, supported by numerous studies of all three IP_3R subtypes and using both single-channel and Ca^{2+} -efflux studies, is that all three IP_3R subtypes are biphasically regulated by cytosolic Ca^{2+} (Marshall and Taylor 1993; Oancea and Meyer 1996; Dufour et al. 1997; Missiaen et al. 1998; Miyakawa et al. 1999; Swatton et al. 1999; Boehning and Joseph 2000b; Mak et al. 2000; Mak et al. 2001; Tu et al. 2005a). Two independent Ca^{2+} -binding sites, which differ in their interactions with different bivalent cations and in their affinities for Ca^{2+} , mediate the stimulatory and inhibitory effects of cytosolic Ca^{2+} (Marshall and Taylor 1994; Striggow and Ehrlich 1996; Hajnóczky and Thomas 1997). Both sites are essential elements of many models proposed to explain regenerative Ca^{2+} signals (Lechleiter et al. 1991; Berridge 1997). This core biphasic pattern of regulation by cytosolic Ca^{2+} may be modulated by other intracellular signals (and these, too, may have contributed to some of the disparate findings) and by processing of IP_3R . Ca^{2+} -dependent inhibition of $\text{IP}_3\text{R3}$, for example, is very sensitive to cytoplasmic ATP (Tu et al. 2005b), and the neuronal S2^+ splice variant of $\text{IP}_3\text{R1}$ has a broader Ca^{2+} -dependence than the peripheral S2^- form (Tu et al. 2002). However, IP_3 is the major influence on what Ca^{2+} does to IP_3R : The two ligands are essential co-agonists of IP_3R (Finch et al. 1991). Activation of $\text{IP}_3\text{R1}$ by Ca^{2+} is positively cooperative, enabling P_o to reach its maximum value over a narrow range of Ca^{2+} concentrations, suggesting that $\text{IP}_3\text{R1}$ may be well suited to mediating CICR and regenerative Ca^{2+} signals. Activation of $\text{IP}_3\text{R3}$ is less cooperative, occurs over a broader range of Ca^{2+} concentrations, and requires lesser activation, making it well suited as a trigger for Ca^{2+} release as the level of IP_3 increases (Mak et al. 2001; Foskett et al. 2007).

Foskett and colleagues have argued, from their analyses of patch-clamp recordings of nuclear IP_3R , that IP_3 decreases the sensitivity of the IP_3R to inhibition by cytosolic Ca^{2+} , and that this alone is the means whereby IP_3 stimulates channel opening (Mak et al. 1998; Mak et al. 2001; Ionescu et al. 2006). This simple explanation, where IP_3 serves only to relieve tonic inhibition by resting Ca^{2+} concentrations, is impossible to reconcile with their observation that pretreatment of cells with Ca^{2+} -free media abolishes Ca^{2+} inhibition without preventing IP_3 from activating IP_3R (Mak et al. 2003). This simple model was later elaborated to include at least three different Ca^{2+} sensors (Mak et al. 2003), but at the core of this revised scheme is a single Ca^{2+} -binding site that switches from being inhibitory in the absence of IP_3 to stimulatory in its presence (Mak et al. 2003). The essential feature of this scheme is consistent with our initial model, derived from rapid superfusion analysis, which suggests that IP_3 both relieves Ca^{2+} inhibition and promotes binding of Ca^{2+} to a stimulatory site (Marchant and Taylor 1997; Adkins and Taylor 1999). The latter is essential for the channel to open. We, however, argue that the stimulatory and inhibitory Ca^{2+} -binding sites are distinct (Marshall and Taylor 1994). We suggest, therefore, that the essential role of IP_3 is to promote Ca^{2+} binding to a stimulatory Ca^{2+} -binding site. IP_3 , by priming this site, allows Ca^{2+} to provide instantaneous control over whether the channel opens (Fig. 1A).

The structural basis for Ca^{2+} -regulation of IP_3R is unresolved: it may be either direct, via Ca^{2+} binding to a site intrinsic to the IP_3R or via an accessory Ca^{2+} -binding protein (Taylor et al. 2004). Stimulation of IP_3R by cytosolic Ca^{2+} is universally observed even with purified IP_3R reconstituted into lipid bilayers (Ferris et al. 1989; Hirota et al. 1995; Michikawa et al. 1999), suggesting that this essential Ca^{2+} -binding site probably resides within the primary sequence of the IP_3R . At least seven cytosolic Ca^{2+} -binding sites have been identified within $\text{IP}_3\text{R1}$ (Sienaert et al. 1996; Sienaert et al. 1997), but the physiological relevance of these sites is unresolved. Two of the sites (residues 304–381 and 378–450) are within the IP_3 -binding

Figure 1. Regulation of IP₃R by cytosolic and luminal Ca²⁺. (A) Binding of IP₃ (black circle) to the IP₃R determines whether a stimulatory (green) or inhibitory (red) Ca²⁺-binding site is available (Adkins and Taylor 1999). IP₃ binding causes the stimulatory site to become accessible and the inhibitory site to be concealed; binding of Ca²⁺ (blue circle) to the former then triggers opening of the channel. (B) Luminal Ca²⁺ is proposed to tune the sensitivity of the IP₃R to cytosolic IP₃ and Ca²⁺ such that full stores (i) are most sensitive to IP₃. As the IP₃R opens (ii) and the stores lose Ca²⁺, they are proposed to lose sensitivity to IP₃ until eventually the IP₃R closes, despite the continued presence of the cytosolic stimuli, trapping Ca²⁺ within the ER (iii). Conversely, stores regain their sensitivity to IP₃ as the stores refill, perhaps thereby determining the interval between Ca²⁺ spikes in stimulated cells (Berridge 2007).

core, for which there is a high-resolution structure (Bosanac et al. 2002). This structure shows two surface-exposed clusters of acidic residues that overlap with residues in the second N-terminal Ca²⁺-binding region. However, point mutations of several of these acidic residues had no effect on Ca²⁺-regulation of IP₃R (Joseph et al. 2005). The remaining Ca²⁺-binding sites fall within the central region of the IP₃R (Sienaert et al. 1996; Sienaert et al. 1997). The site between

residues 1347–1426 is interesting because its proximity to a calmodulin (CaM)-binding region is reminiscent of RyR, which have two CaM-binding regions within ~200 residues of high-affinity Ca²⁺-binding sites, and a third flanked by two high-affinity Ca²⁺-binding sites (Chen and MacLennan 1994). Interactions between these sites have been proposed to contribute to regulation of RyR by Ca²⁺ and CaM (Chen and MacLennan 1994). None of the Ca²⁺-binding sites within IP₃R contain EF-hands or any other known Ca²⁺-binding motif, and none have obvious sequence similarity with similar regions in RyR. However, each site has clusters of negatively charged residues that may coordinate Ca²⁺ (Sienaert et al. 1997). There is presently no evidence to link any of these sites directly to Ca²⁺ regulation of IP₃R. The only tangible link between specific residues and Ca²⁺ regulation comes from mutagenesis of a glutamate residue that is conserved in all IP₃R and RyR. Mutation of this residue in RyR massively reduced the Ca²⁺ sensitivity of the channel (Chen et al. 1998; Li and Chen 2001). Mutation of the same residue (Glu-2100) to another acidic residue (Asp) caused a ~5- to 10-fold decrease in the Ca²⁺-sensitivity of the IP₃R to both stimulation and inhibition, abolished oscillatory Ca²⁺ transients in response to agonist stimulation, and reduced the Ca²⁺-binding affinity of a large fragment that includes the residue (Miyakawa et al. 2001; Tu et al. 2003). A rather puzzling aspect of these results is the observation that mutation of a single residue similarly attenuates both stimulation and inhibition by Ca²⁺, when other evidence suggests that the two effects are mediated by distinct sites. This, together with the lack of direct evidence that Ca²⁺ is coordinated by the conserved glutamate, leaves open the possibility that rather than itself contributing to an essential Ca²⁺-binding site, this residue may be allosterically coupled to the site.

Ca²⁺-mediated inhibition of IP₃R is widely assumed to contribute to termination of local cytosolic Ca²⁺ signals, but it remains far from clear whether such inhibition is mediated by Ca²⁺ binding directly to IP₃R or to an associated protein (Taylor and Laude 2002). The effects of

C.W. Taylor and S.C. Tovey

Ca^{2+} on IP_3 binding differ between subtypes: It inhibits binding to $\text{IP}_3\text{R1}$ (Worley et al. 1987; Supattapone et al. 1988; Joseph et al. 1989; Varney et al. 1990; Richardson and Taylor 1993; Benevolensky et al. 1994; Cardy et al. 1997; Yoneshima et al. 1997), but the effects of Ca^{2+} on IP_3 binding to IP_3R from cells expressing predominantly $\text{IP}_3\text{R2}$ or $\text{IP}_3\text{R3}$ are confused (Pietri et al. 1990; Mohr et al. 1993; Marshall and Taylor 1994; Cardy et al. 1997; Yoneshima et al. 1997; Lin et al. 2000; Swatton and Taylor 2002). These conflicting results, and evidence that purified $\text{IP}_3\text{R1}$ is not inhibited by Ca^{2+} (Danoff et al. 1988; Richardson and Taylor 1993; Benevolensky et al. 1994; Lin et al. 2000), lend some support to the idea that Ca^{2+} inhibition may be mediated by an accessory protein. It is, however, noteworthy that deletion of the suppressor domain (SD, residues 1-223) of $\text{IP}_3\text{R1}$, which appears not to include a Ca^{2+} -binding site, abolishes inhibition of IP_3 binding by Ca^{2+} (Sienaert et al. 2002). This suggests that effective regulation by an accessory protein might require the SD.

Calmodulin (CaM) is one candidate for the accessory protein through which Ca^{2+} inhibition is exercised (Nadif Kasri et al. 2002; Taylor and Laude 2002). CaM is a ubiquitously expressed, EF-hand containing, Ca^{2+} -binding protein that serves as the Ca^{2+} -sensor for many cellular events (Gnegy 1993). All IP_3R subtypes are inhibited by Ca^{2+} -CaM (Hirota et al. 1999; Michikawa et al. 1999; Missiaen et al. 1999; Adkins et al. 2000; Missiaen et al. 2000), and CaM has been shown to restore Ca^{2+} inhibition to purified IP_3R (Hirota et al. 1999; Michikawa et al. 1999; Nosyreva et al. 2002). Yet, it has proven difficult to relate these functional effects of CaM to either its effects on IP_3 binding or to identified CaM-binding sites within IP_3R . CaM inhibits IP_3 binding to $\text{IP}_3\text{R1}$ in a Ca^{2+} -independent manner (Patel et al. 1997; Cardy and Taylor 1998), through a site that probably lies within the SD (Adkins et al. 2000; Sienaert et al. 2002). Its properties are clearly inconsistent with the ability of CaM to inhibit IP_3R function only in the presence of Ca^{2+} . There is a high-affinity Ca^{2+} -CaM-binding site within the central region of $\text{IP}_3\text{R1}$ and $\text{IP}_3\text{R2}$, but not

$\text{IP}_3\text{R3}$ (Yamada et al. 1995; Lin et al. 2000). However, mutations that prevented Ca^{2+} -CaM binding to this site had no effect on Ca^{2+} -dependent inhibition of IP_3R (Zhang and Joseph 2001; Nosyreva et al. 2002). This evidence and the absence of the site from $\text{IP}_3\text{R3}$ suggest that the central Ca^{2+} -CaM-binding site cannot be responsible for Ca^{2+} inhibition of IP_3R . An additional high-affinity Ca^{2+} -CaM-binding site is created in $\text{IP}_3\text{R1}$ after removal of the S2 splice region: While this may increase the Ca^{2+} -CaM sensitivity of peripheral S2⁻ $\text{IP}_3\text{R1}$, it is not a universal candidate for mediating Ca^{2+} inhibition of IP_3R (Islam et al. 1996; Lin et al. 2000). Recently, it was suggested that bound CaM is essential for IP_3R function because a peptide antagonist of CaM inhibited IP_3 -evoked Ca^{2+} release (Nadif Kasri et al. 2006). It is now clear that this peptide acts directly on IP_3R , with no requirement for CaM (Sun and Taylor 2008). While this eliminates an essential role for tethered CaM, it raises the intriguing possibility that an endogenous CaM-like structure might be essential for IP_3R activation (Sun and Taylor 2008). In summary, all IP_3R subtypes are inhibited by Ca^{2+} -CaM, but the molecular basis of this inhibition has not been established. It seems, on balance, that CaM is unlikely to be the accessory protein through which Ca^{2+} universally inhibits IP_3R . That need not preclude a role for CaM in modulating IP_3R function (Taylor and Laude 2002), just as it does for RyR (Chen et al. 1997; Fruen et al. 2000; Rodney et al. 2001), but we must look elsewhere for the site through which Ca^{2+} inhibits IP_3R .

We turn now to the luminal surface of the IP_3R , where, and again drawing parallels with RyR, we consider regulation by luminal Ca^{2+} . Persuasive evidence suggests that Ca^{2+} release by RyR may be terminated before Ca^{2+} stores are entirely depleted because luminal Ca^{2+} is required to maintain RyR activity (Györke and Györke 1998; Launikonis et al. 2006; Jiang et al. 2008), possibly via its interaction with calsequestrin, a luminal high-capacity Ca^{2+} -binding protein (Launikonis et al. 2006; Terentyev et al. 2006). A similar scheme has been proposed to account for two features of IP_3 -evoked Ca^{2+} release: the initiation of Ca^{2+} release after the

quiescent interspike interval during repetitive Ca²⁺ spikes (Berridge 2007) and quantal Ca²⁺ release via IP₃R. The latter describes the situation wherein unidirectional Ca²⁺ efflux from intracellular stores terminates before the stores have fully emptied after stimulation with sub-maximally effective concentrations of IP₃ without loss of their ability to respond to a further increase in IP₃ concentration (Muallem et al. 1989; Meyer and Stryer 1990; Taylor and Potter 1990; Oldershaw et al. 1991; Bootman et al. 1992; Brown et al. 1992; Combettes et al. 1992; Ferris et al. 1992; Hirota et al. 1995). The proposal is that luminal Ca²⁺ sets the gain on the regulation by cytosolic IP₃ and Ca²⁺, so that as the luminal free Ca²⁺ concentration falls, it causes the sensitivity of the IP₃R to IP₃ to fall until, as Ca²⁺ leaks from the ER, the IP₃R closes despite the continued presence of cytosolic IP₃ and residual Ca²⁺ within the ER (Irvine 1990). Conversely, as stores refill between Ca²⁺ spikes in an intact cell, the model predicts that the sensitivity of the IP₃R increases until it exceeds the threshold at which prevailing cytosolic IP₃ and Ca²⁺ concentrations become sufficient to trigger opening (Fig. 1B). Despite the enduring appeal of the model, evidence that luminal Ca²⁺ directly regulates IP₃R is not yet entirely convincing.

Stores loaded with Ca²⁺ have been shown to become more sensitive to IP₃ in some studies (Missiaen et al. 1992; Nunn and Taylor 1992; Oldershaw and Taylor 1993; Parys et al. 1993; Missiaen et al. 1994; Horne and Meyer 1995; Combettes et al. 1996; Tanimura and Turner 1996), but not in others (Combettes et al. 1992; Shuttleworth 1992; Combettes et al. 1993; van de Put et al. 1994). However, even the supportive results do not eliminate the possibility that the increased sensitivity to IP₃ arises from having Ca²⁺ pass through active IP₃R and increase their sensitivity from the cytosolic surface. Similar difficulties have plagued analyses of the effects of luminal Ca²⁺ on RyR (Tripathy and Meissner 1996; Laver 2007; Laver 2009). In bilayer recordings of IP₃R1, where essential accessory proteins may be lost, luminal Ca²⁺ failed to potentiate the Ca²⁺ release evoked by IP₃ (Bezprozvanny and Ehrlich 1994). Despite the caveats, regulation of

IP₃R by luminal Ca²⁺ deserves serious consideration. A high-affinity Ca²⁺-binding site within the luminal loop linking TMD 5 and 6 (Sienaert et al. 1996) contains conserved acidic residues that could mediate luminal Ca²⁺ regulation, although the sub-μM affinity of this site for Ca²⁺ would be ill-suited to detecting likely changes in luminal Ca²⁺ concentration. Luminal accessory proteins, akin to those that regulate RyR, are another possibility, with ERp44 being one candidate. ERp44 belongs to the thioredoxin protein family and regulates IP₃R in a pH- and luminal Ca²⁺-dependent manner (Higo et al. 2005). Binding of ERp44 to the TMD5-6 loop of IP₃R inhibits channel activity, and the interaction is disrupted by high concentrations of Ca²⁺ consistent with the suggestion that luminal Ca²⁺ might enhance IP₃R activity.

To summarize, IP₃ works by tuning the Ca²⁺ sensitivity of the IP₃R: It stimulates Ca²⁺ binding to a stimulatory site and inhibits Ca²⁺ binding to an inhibitory site (Fig. 1A). Binding to the stimulatory site is the trigger for opening of the pore. The identity of neither Ca²⁺-binding site is known: The stimulatory site probably resides within the IP₃R itself, but the inhibitory site may require an accessory protein, though this is unlikely to be CaM. Luminal Ca²⁺ may further tune the sensitivity of the IP₃R to regulation by its cytosolic ligands, but this remains unproven.

STRUCTURAL DETERMINANTS OF IP₃R ACTIVATION

Judged by their primary amino acid sequences, all known IP₃R subunits are assumed to have a similar architecture. Each subunit, of about 2700 residues, comprises three major regions: the N-terminal to which IP₃ binds, the C-terminal region with its six transmembrane regions (TMD) (Galvan et al. 1999), and a large intervening sequence (Fig. 2A). Functional IP₃Rs are tetrameric, assembled either from identical subunits or from mixtures of the three subtypes and their many splice variants (Taylor et al. 1999; Foskett et al. 2007). Several structures of the entire IP₃R1 have been published, each derived from single particle analysis of images from electron microscopy (Hamada and Mikoshiba

C.W. Taylor and S.C. Tovey

Figure 2. Major structural domains of IP₃R. (A) The three key regions defined by the primary sequence of a single IP₃R subunit are highlighted: the N-terminal with its SD and IBC, the C-terminal region with its six TMD and pore, and the large central region. Atomic structures of the SD (Bosanac et al. 2005) and IBC with IP₃ bound (Bosanac et al. 2002) are also shown. (B) Two views of the IP₃R derived from single particle analysis (da Fonseca et al. 2003) (*top*, from the cytosol; *bottom*, across the ER membrane with the ER lumen at the top). (C) A possible structure of the IP₃R pore, with a luminal selectivity filter and a constriction formed by the tepee-like structure of TMD6. Only two of the four IP₃R subunits are shown.

2002; Jiang et al. 2002a; da Fonseca et al. 2003; Hamada et al. 2003; Serysheva et al. 2003; Sato et al. 2004). These studies confirm the tetrameric state of IP₃R, but variability between the structures and their relatively low resolution (~ 30 Å) have, so far, limited any realistic interpretation of the structural basis of IP₃R activation (Taylor et al. 2004) (Fig. 2B). Whether structures of recombinant IP₃R will contribute to resolving this impasse remains to be seen (Wolfram et al. 2010).

There has been more progress with RyR, although only recently has the resolution of these structures (~ 30 Å) improved on that obtained for IP₃R. These structures of native RyR, and all three subtypes of recombinant RyR reveal a shape like a square mushroom with a very large, open cytoplasmic structure tethered

to a much smaller TMD region (the stalk). At ~ 30 Å resolution, the structures of the three RyR subtypes are almost indistinguishable, and because they, like the three subtypes of IP₃R, share about 65% sequence identity, it seems reasonable to suppose that the 3D structures of all IP₃R are also likely to be similar to each other. These studies of RyR have identified positions of critical residues within the 3D structure, the sites to which accessory proteins bind, and conformational changes associated with opening of the pore (Orlova et al. 1996; Serysheva et al. 2005; Wang et al. 2007; Jones et al. 2008). Activation of RyR is associated with considerable changes in both the pore and cytoplasmic regions: The four corners of the latter dip down toward the SR, while the central region lifts away from it (Samso et al. 2009). It is noteworthy, in

the context of schemes for activation of IP₃R (see below), that large movements of some cytoplasmic domains of RyR1 appear to occur around hinges that link them to relatively immobile domains.

The highest resolution maps ($\sim 10\text{\AA}$), although still insufficient to map 3D structure to primary sequence, have come close to defining the likely secondary structure of the pore of RyR1 (Ludtke et al. 2005; Samso et al. 2005; Samso et al. 2009). This region appears to have six α -helices (Samso et al. 2009), consistent with models of RyR that suggest six TMD (Meur et al. 2007). Along the central axis, it has a luminal constriction (probably the selectivity filter, see below) and a tepee-like assembly of four inner helices (likely to be TMD6), with the apex pointing into the cytoplasmic structure. By analogy with MthK channels, this constriction may form the gate of the RyR. Kinking of the inner helix around a central Gly residue causes splitting of the tepee and thereby opening of the channel for MthK (Jiang et al. 2002b). One structure (Samso et al. 2009) is consistent with a similar mechanism operating for RyR1, but another structure (Ludtke et al. 2005) and mutagenesis of the critical Gly (G4863 in RyR1) (Wang et al. 2003) contradict it. These insights into the possible workings of the RyR pore are significant for IP₃R, because it is within the pore region (TMD5-6) that RyR and IP₃R share the greatest sequence similarity. We turn, therefore, to the pore of the IP₃R to explore its properties and structure.

All IP₃R (like all RyR) are cation channels with extremely large conductance, but only modest selectivity for Ca²⁺ over monovalent cations (permeability ratio, $P_{\text{Ca}}/P_{\text{K}} \sim 6$) (Williams et al. 2001; Foskett et al. 2007). The voltage-gated and store-operated Ca²⁺ channels that mediate Ca²⁺ entry across the plasma membrane are vastly more selective ($P_{\text{Ca}}/P_{\text{K}} > 1000$). In the ER, where most IP₃Rs are located, this lack of selectivity is unlikely to be a problem because Ca²⁺ is probably the only cation with an appreciable electrochemical gradient across the ER membrane. In effect, the ER Ca²⁺ pump (SERCA), by creating a steep Ca²⁺ concentration gradient across the ER membrane, assumes responsibility

for determining which cations flow through an open IP₃R. Indeed, the K⁺ permeability of IP₃R and RyR may facilitate rapid Ca²⁺ release by allowing K⁺ to move into the ER to electrically compensate the efflux of Ca²⁺ (Gillespie and Fill 2008). The pore of the IP₃R, like that of RyR, is formed by the final pair of TMD (TMD5-6) and the luminal loop that links them from each of the four subunits (Ramos-Franco et al. 1999; Williams et al. 2001) (Fig. 2C). The loop includes a sequence (GGVGD in IP₃R) similar to that of the selectivity filter of K⁺ channels (Balshaw et al. 1999), consistent with the idea that the overall architecture of the pore region may be broadly similar to that of K⁺ channels (MacKinnon 2004). For both IP₃R and RyR, however, the pore must be larger and less-selective than for K⁺ channels, and probably able to accommodate only one cation at a time (Williams et al. 2001). This model for the IP₃R pore, where TMD5 (the outer helix) and TMD6 (inner helix) cradle a short pore helix and selectivity filter (Fig. 2C), is consistent with mutagenesis of residues within this region affecting ion permeation (Boehning et al. 2001b; Dellis et al. 2006; Dellis et al. 2008; Schug et al. 2008), with biophysical evidence that the narrowest region of the pore lies close to the luminal entrance of the RyR (Williams et al. 2001) and with the intermediate resolution structures of the pore region of RyR1 (Samso et al. 2009). A conserved acidic residue (D2550 in IP₃R1) at the luminal end of the selectivity filter (Fig. 2C) contributes to the modest Ca²⁺ selectivity of IP₃R (Boehning et al. 2001b; Dellis et al. 2008) and RyR (Gao et al. 2000; Wang et al. 2005; Gillespie 2008), but the structural determinants of ion selectivity and permeation by IP₃R are otherwise poorly understood. The changes in pore structure that allow it to open are also minimally understood. Indeed, mutation of the conserved Gly within TMD6 of IP₃R (G2586 in IP₃R1), which might have been thought to provide the gating hinge (Samso et al. 2009), appears not to prevent IP₃ from opening IP₃R (Schug et al. 2008). In short, aside from knowing the regions of primary sequence that form the IP₃R pore (TMD5-6) and a rather vague notion that its structure perhaps resembles that of K⁺ channels, we have only the most

C.W. Taylor and S.C. Tovey

rudimentary knowledge of the structural determinants of how the IP₃R pore opens and selects between ions.

The conformational changes in the IP₃R that lead to opening of its pore are initiated by IP₃ binding to the IP₃-binding core (IBC, residues 224–604 in IP₃R1) (Fig. 3A). Although IP₃ is the only endogenous ligand of the IBC, there are many synthetic agonists, all of which have structures equivalent to the equatorial 6-hydroxyl and the 4- and 5-phosphate groups of IP₃ (Fig. 3A) (Rossi et al. 2010). It is noteworthy that neither of the immediate products of IP₃ metabolism, IP₂ and IP₄, binds to the IBC; both metabolic pathways are therefore effective means of terminating activation of IP₃R by IP₃. An atomic structure of the IBC with IP₃ bound (Bosanac et al. 2002) shows IP₃ held within a clam-like structure in which the phosphate groups of IP₃ are coordinated by basic residues (Fig. 3A). The two sides of the clam,

the α - and β -domains, form a network of interactions with the essential groups of IP₃. The 4-phosphate is hydrogen-bonded with residues in the β -domain, the 5-phosphate forms hydrogen bonds with residues predominantly in the α -domain, and the 6-hydroxyl interacts with the backbone of a residue within the α -domain. It is easy to imagine how these interactions with IP₃ might pull the α - and β -domains together, causing the clam to close in a manner similar to glutamate binding to ionotropic glutamate receptors (Mayer 2006). Structures of the IBC without IP₃ bound are urgently needed to assess this proposal, but two lines of evidence lend circumstantial support. First, the IBC adopts a more constrained structure when it binds IP₃ (Chan et al. 2007). Second, adenophostins, which are high-affinity agonists of IP₃R (Rossi et al. 2010), retain some activity after loss of the 3-phosphate (analogous to the 5-phosphate of IP₃), probably because their adenine moiety

Figure 3. Initiation of IP₃R activation by IP₃. (A) The structure of IP₃, with its critical vicinal 4,5-bisphosphate and 6-hydroxyl groups, is shown alongside the structure of the IBC with IP₃ bound. The latter shows the 4- and 5-phosphates contacting the β - and α -domains, respectively (Bosanac et al. 2002), and thereby pulling the clam into a more closed state. (B) Structure of the SD (Bosanac et al. 2005) showing possible sites of interaction with the IBC and downstream domains through which it signals to the pore. See text for further details.

interacts strongly with a residue in the α -domain and thereby partially mimics the clam-closure that would otherwise require the 5-phosphate to bind to the α -domain (Sureshan et al. 2009). We envisage, therefore, that when IP₃ binds to the IBC, the essential vicinal phosphate groups through their contacts with the α - and β -domains effectively cross-bridge the two sides of the clam-like structure, causing it to close, and thereby initiate the processes that will culminate in opening of the pore.

It is worth commenting briefly on available antagonists of IP₃R because of their obvious value as experimental tools. There are no specific antagonists of IP₃R, although with appropriate caution some antagonists can yield useful insight (Michelangeli et al. 1995). Heparin is a competitive antagonist of IP₃ (Worley et al. 1987), although it is not membrane-permeant and, among many additional effects, it uncouples G-protein-coupled receptors from their G proteins (Dasso and Taylor 1991) and activates RyR (Ehrlich et al. 1994). 2-aminoethyl diphenylboronate (2-APB) is membrane-permeant and inhibits IP₃-evoked Ca²⁺ release without affecting IP₃ binding (Maruyama et al. 1997); its mechanism of action is unresolved. However, 2-APB also inhibits Ca²⁺ uptake and many other Ca²⁺ channels. It has recently aroused interest as a modulator of STIM and, therefore, store-operated Ca²⁺ entry (Goto et al. 2010). A screen of 2-APB analogues with selectivity for store-operated Ca²⁺ entry may yet also provide IP₃R-selective antagonists (Goto et al. 2010). Xestospongins, isolated from an Australian sponge, are high-affinity membrane-permeant inhibitors of IP₃-evoked Ca²⁺ release that do not affect IP₃ binding (Gafni et al. 1997), but they, too, have side effects (Solovyova et al. 2002). High concentrations of caffeine inhibit IP₃-evoked Ca²⁺ release (Parker and Ivorra 1991) without affecting IP₃ binding (Worley et al. 1987), but caffeine also stimulates RyR, inhibits cyclic nucleotide phosphodiesterases, and interferes with many Ca²⁺ indicators. Membrane-permeant peptide antagonists of IP₃R may provide another potential source of selective antagonists (Sun and Taylor 2008).

How IP₃ binding to the IBC leads to binding of Ca²⁺ to the IP₃R, and thereby opening of the pore, remains largely unknown, but it is clear that the suppressor domain (SD, residues 1-223 of IP₃R1), which is connected to the IBC by a flexible linkage (Chan et al. 2007), plays an essential role. The clearest evidence is that IP₃ binds to IP₃R without an SD, but it fails to open the pore (Uchida et al. 2003; Szlufcik et al. 2006). The name of this region derives from the observation that, although the SD itself is unlikely to make any direct contacts with IP₃, its presence decreases the affinity of IP₃R for IP₃ (Uchida et al. 2003). We have interpreted this effect to reflect the use of binding energy from the binding of IP₃ to the IBC to cause a conformational change within the SD. This interpretation gains considerable support from our analysis of partial agonists of the IP₃R (Rossi et al. 2009). The crux of our argument is that the energy provided by agonist binding drives both the conformational changes that lead to receptor activation and tighter binding of the ligand to its receptor. There is, therefore, a play-off between these two claims on the binding energy. Partial agonists, because they less effectively activate the receptor, divert more binding energy into stabilizing the binding, while full agonists evoke more substantial conformational changes; therefore, less binding energy remains to stabilize binding. Our results show that although full and partial agonists bind with similar affinities to the IBC, the SD causes the affinity of full agonists to decrease more than for partial agonists (Rossi et al. 2009). Quantitative analyses of these results lead to the conclusion that the most energetically costly conformational change in the IP₃R evoked by IP₃ occurs within its N-terminal (residues 1-604), and that these conformational changes pass entirely via the SD to the pore region (Rossi et al. 2009). We suggest, therefore, that the SD is the essential link between IP₃ binding to the IBC and the subsequent conformational changes that lead to opening of the pore. Without a structure of the entire N-terminal of the IP₃R, we can only speculate on the physical relationship between the IBC and SD, but our results with partial agonists and mutagenesis are

C.W. Taylor and S.C. Tovey

consistent with three exposed loops of the SD ($\beta 2$ – $\beta 3$, $\beta 5$ – $\beta 6$, and $\beta 7$ – $\beta 8$, blue in Fig. 3B) being the most likely sites of interaction with the IBC (Rossi et al. 2009).

Remarkably, and despite their rather low sequence identities ($\sim 30\%$), the crystal structures of the SD from IP₃R1 (Bosanac et al. 2005) and of the analogous N-terminal regions from RyR1 and RyR2 (Amador et al. 2009; Lobo and Van Petegem 2009) are extremely similar. Several mutations associated with malignant hyperthermia and central core disease (RyR1) or catecholaminergic polymorphic ventricular tachycardia (RyR2), all of which impair the normal regulation of gating, are clustered in an exposed loop ($\beta 8$ – $\beta 9$) of the N-terminal of RyR (Amador et al. 2009). Furthermore, and consistent with the N-terminal of the RyR mediating essential interdomain interactions, a peptide derived from this region causes RyR2 to open spontaneously, apparently by uncoupling an interaction between the endogenous loop and a central region of the RyR that includes residues 2460–2495 (Oda et al. 2005; Tateishi et al. 2009). In light of the conservation of structure between IP₃R and RyR, it is tempting to speculate that the same loop in the SD of the IP₃R ($\beta 8$ – $\beta 9$, red in Fig. 3B) may mediate transfer of conformational changes onward toward the pore. Co-immunoprecipitation studies have suggested an interaction between the N-terminal of IP₃R1 (most likely the SD) and the pore region of an adjacent subunit (Boehning and Joseph 2000a), perhaps mediated by the cytosolic loop linking TMD4 to TMD5 (Schug and Joseph 2006). An attractive possibility, therefore, is that the SD (perhaps its $\beta 8$ – $\beta 9$ loop) interacts directly with the short cytosolic helix linking TMD4 and TMD5, and thereby gates the pore (Schug and Joseph 2006; Rossi et al. 2009). Such an interaction would require that the SD comes very close to the pore, but the exact location of the SD within the 3D structure of either the IP₃R or RyR is unknown. The N-terminal of the RyR probably lies within the clamp region at the periphery of the large square cytoplasmic structure (Wang et al. 2007), and it does change shape during RyR activation (Samso et al. 2009). Yet, in this location the N-terminal is

too far from the pore to interact directly with the TMD4-5 loop, consistent perhaps with evidence that in RyR the N-terminal may interact directly with a neighboring domain that includes residues from the central part of the primary sequence (Wang et al. 2007). These observations and the evidence that the uncoupling peptide derived from the N-terminal of RyR2 appears to interact with residues remote from the pore (Oda et al. 2005; Tateishi et al. 2009), suggest that the links between the SD and pore may, at least for RyR, be indirect.

In summary, we suggest that IP₃R activation is initiated when IP₃ binds to the IBC, and perhaps thereby causes closure of its clam-like structure. That conformational change, which must also initiate the events that allow Ca²⁺ to bind to a stimulatory site, is passed to the rest of the IP₃R entirely via the SD. The location of that Ca²⁺-binding site and, therefore, the structural links between it and the SD, are unresolved. We speculate that one face of the SD interacts directly with the IBC, and the opposite face interacts with the structure through which conformational changes pass to the pore. The pore is a relatively nonselective, large-conductance cation channel formed by the tetrameric assembly of the TMD5-6 regions of each subunit. Its structure is unresolved but likely to be broadly similar to K⁺ channels with a selectivity filter and gate at opposite ends of its membrane-spanning structure.

ACKNOWLEDGMENTS

Work from the authors' laboratory is supported by the Wellcome Trust.

REFERENCES

- Adkins CE, Morris SA, De Smedt H, Török K, Taylor CW. 2000. Ca²⁺-calmodulin inhibits Ca²⁺ release mediated by type-1, -2 and -3 inositol trisphosphate receptors. *Biochem J* **345**: 357–363.
- Adkins CE, Taylor CW. 1999. Lateral inhibition of inositol 1,4,5-trisphosphate receptors by cytosolic Ca²⁺. *Curr Biol* **9**: 1115–1118.
- Amador FJ, Liu S, Ishiyama N, Plevin MJ, Wilson A, MacLennan DH, Ikura M. 2009. Crystal structure of type I ryanodine receptor amino-terminal β -trefoil domain reveals a disease-associated mutation “hot spot” loop. *Proc Natl Acad Sci USA* **106**: 11040–11044.

- Balshaw D, Gao L, Meissner G. 1999. Luminal loop of the ryanodine receptor: A pore-forming segment? *Proc Natl Acad Sci USA* **96**: 3345–3347.
- Benevolensky D, Moraru II, Watras J. 1994. Micromolar calcium decreases the affinity of inositol trisphosphate receptor in vascular smooth muscle. *Biochem J* **299**: 631–636.
- Berridge MJ. 1975. The interaction of cyclic nucleotides and calcium in the control of cellular activity. In *Advances in Cyclic Nucleotide and Protein Phosphorylation Research*, (ed. P Greengard, GA Robison), pp. 1–98. Raven Press, New York.
- Berridge MJ. 1983. Rapid accumulation of inositol trisphosphate reveals that agonists hydrolyse polyphosphoinositides instead of phosphatidylinositol. *Biochem J* **212**: 849–858.
- Berridge MJ, (ed.) 1995. *CIBA Foundation Symposium Calcium waves, gradients and oscillations*. John Wiley and Sons, Chichester.
- Berridge MJ. 1997. Elementary and global aspects of calcium signalling. *J Physiol* **499**: 291–306.
- Berridge MJ. 2005. Unlocking the secrets of cell signaling. *Annu Rev Physiol* **67**: 1–21.
- Berridge MJ. 2007. Inositol trisphosphate and calcium oscillations. *Biochem Soc Symp* **74**: 1–7.
- Berridge MJ, Fain JN. 1979. Inhibition of phosphatidylinositol synthesis and the inactivation of calcium entry after prolonged exposure of the blowfly salivary gland to 5-hydroxytryptamine. *Biochem J* **178**: 59–69.
- Berridge MJ, Irvine RF. 1984. Inositol trisphosphate, a novel second messenger in cellular signal transduction. *Nature* **312**: 315–321.
- Berridge MJ, Irvine RF. 1989. Inositol phosphates and cell signalling. *Nature* **341**: 197–205.
- Berridge MJ, Lipp P, Bootman MD. 2000. The versatility and universality of calcium signalling. *Nat Rev Mol Cell Biol* **1**: 11–21.
- Bezprozvanny I, Ehrlich BE. 1994. Inositol (1,4,5)-trisphosphate (InsP₃)-gated Ca channels from cerebellum: conduction properties for divalent cations and regulation by intraluminal calcium. *J Gen Physiol* **104**: 821–856.
- Bezprozvanny I, Watras J, Ehrlich BE. 1991. Bell-shaped calcium-response curves for Ins(1,4,5)P₃- and calcium-gated channels from endoplasmic reticulum of cerebellum. *Nature* **351**: 751–754.
- Boehning D, Joseph SK. 2000a. Direct association of ligand-binding and pore domains in homo- and heterotetrameric inositol 1,4,5-trisphosphate receptors. *EMBO J* **19**: 5450–5459.
- Boehning D, Joseph SK. 2000b. Functional properties of recombinant type I and type III inositol 1,4,5-trisphosphate receptor isoforms expressed in COS-7 cells. *J Biol Chem* **275**: 21492–21499.
- Boehning D, Joseph SK, Mak D-OD, Foskett JK. 2001a. Single-channel recordings of recombinant inositol trisphosphate receptors in mammalian nuclear envelope. *Biophys J* **81**: 117–124.
- Boehning D, Mak D-OD, Foskett JK, Joseph SK. 2001b. Molecular determinants of ion permeation and selectivity in inositol 1,4,5-trisphosphate receptor Ca²⁺ channels. *J Biol Chem* **276**: 13509–13512.
- Bootman MD, Berridge MJ, Lipp P. 1997. Cooking with calcium: the recipes for composing global signals from elementary events. *Cell* **91**: 367–373.
- Bootman MD, Berridge MJ, Taylor CW. 1992. All-or-nothing Ca²⁺ mobilization from the intracellular stores of single histamine-stimulated HeLa cells. *J Physiol* **450**: 163–178.
- Bosanac I, Alattia J-R, Mal TK, Chan J, Talarico S, Tong FK, Tong KI, Yoshikawa F, Furuichi T, Iwai M, et al. 2002. Structure of the inositol 1,4,5-trisphosphate receptor binding core in complex with its ligand. *Nature* **420**: 696–700.
- Bosanac I, Yamazaki H, Matsu-ura T, Michikawa M, Mikoshiba K, Ikura M. 2005. Crystal structure of the ligand binding suppressor domain of type 1 inositol 1,4,5-trisphosphate receptor. *Mol Cell* **17**: 193–203.
- Brown GR, Sayers LG, Kirk CJ, Michell RH, Michelangeli F. 1992. The opening of the inositol 1,4,5-trisphosphate-sensitive Ca²⁺ channel in rat cerebellum is inhibited by caffeine. *Biochem J* **282**: 309–312.
- Burgess GM, McKinney JS, Fabiato A, Leslie BA, Putney JW Jr. 1983. Calcium pools in saponin-permeabilized guinea pig hepatocytes. *J Biol Chem* **258**: 15336–15345.
- Cardy TJA, Taylor CW. 1998. A novel role for calmodulin: Ca²⁺-independent inhibition of type-1 inositol trisphosphate receptors. *Biochem J* **334**: 447–455.
- Cardy TJA, Traynor D, Taylor CW. 1997. Differential regulation of types 1 and 3 inositol trisphosphate receptors by cytosolic Ca²⁺. *Biochem J* **328**: 785–793.
- Chan J, Whitten AE, Jeffries CM, Bosanac I, Mal TK, Ito J, Porumb H, Michikawa T, Mikoshiba K, Trewella J, et al. 2007. Ligand-induced conformational changes via flexible linkers in the amino-terminal region of the inositol 1,4,5-trisphosphate receptor. *J Mol Biol* **373**: 1269–1280.
- Chen SRW, Ebisawa K, Li X, Zhang L. 1998. Molecular identification of the ryanodine receptor Ca²⁺ sensor. *J Biol Chem* **273**: 14675–14678.
- Chen SRW, Li X, Ebisawa K, Zhang L. 1997. Functional characterization of the recombinant type 3 Ca²⁺ release channel (ryanodine receptor) expressed in HEK293 cells. *J Biol Chem* **272**: 24234–24236.
- Chen SRW, MacLennan DH. 1994. Identification of calmodulin-, Ca²⁺- and ruthenium red-binding domains in the Ca²⁺ release channel (ryanodine receptor) of rabbit skeletal muscle sarcoplasmic reticulum. *J Biol Chem* **269**: 22698–22704.
- Cheng H, Lederer WJ, Cannell MB. 1993. Calcium sparks. Elementary events underlying excitation-contraction coupling in heart muscle. *Science* **262**: 740–744.
- Combettes L, Cheek TR, Taylor CW. 1996. Regulation of inositol trisphosphate receptors by luminal Ca²⁺ contributes to quantal Ca²⁺ mobilization. *EMBO J* **15**: 2086–2093.
- Combettes L, Claret M, Champeil P. 1992. Do submaximal InsP₃ concentrations only induce partial release discharge of permeabilized hepatocyte calcium pools because of the concomitant reduction of intraluminal Ca²⁺ concentration? *FEBS Lett* **301**: 287–290.
- Combettes L, Claret M, Champeil P. 1993. Calcium control of InsP₃-induced discharge of calcium from permeabilized hepatocyte pools. *Cell Calcium* **14**: 279–292.

C.W. Taylor and S.C. Tovey

- da Fonseca PCA, Morris SA, Nerou EP, Taylor CW, Morris EP. 2003. Domain organisation of the type 1 inositol 1,4,5-trisphosphate receptor as revealed by single-particle analysis. *Proc Natl Acad Sci USA* **100**: 3936–3941.
- Danoff SK, Supattapone S, Snyder SH. 1988. Characterization of a membrane protein from brain mediating the inhibition of inositol 1,4,5-trisphosphate receptor binding by calcium. *Biochem J* **254**: 701–705.
- Dasso LTT, Taylor CW. 1991. Heparin and other polyanions uncouple α_1 -adrenoceptors from G-proteins. *Biochem J* **280**: 791–795.
- Dellis O, Dedos S, Tovey SC, Rahman T-U, Dubel SJ, Taylor CW. 2006. Ca^{2+} entry through plasma membrane IP_3 receptors. *Science* **313**: 229–233.
- Dellis O, Rossi AM, Dedos SG, Taylor CW. 2008. Counting functional IP_3 receptors into the plasma membrane. *J Biol Chem* **283**: 751–755.
- Demuro A, Parker I. 2007. Multi-dimensional resolution of elementary Ca^{2+} signals by simultaneous multi-focal imaging. *Cell Calcium* **43**: 367–374.
- Dolmetsch RE, Lewis RS, Goodnow CC, Healy JJ. 1997. Differential activation of transcription factors induced by Ca^{2+} response amplitude and duration. *Nature* **386**: 855–858.
- Dufour J-F, Arias IM, Turner TJ. 1997. Inositol 1,4,5-trisphosphate and calcium regulate the calcium channel function of the hepatic inositol 1,4,5-trisphosphate receptor. *J Biol Chem* **272**: 2675–2681.
- Dupont G, Houart G, De Koninck P. 2003. Sensitivity of CaM kinase II to the frequency of Ca^{2+} oscillations: a simple model. *Cell Calcium* **34**: 485–497.
- Dyer JL, Liu Y, Pino de la Hueria I, Taylor CW. 2005. Long-lasting inhibition of adenylyl cyclase selectively mediated by inositol 1,4,5-trisphosphate-evoked calcium release. *J Biol Chem* **280**: 8936–8944.
- Echevarria W, Leite MF, Guerra MT, Zipfel WR, Nathanson MH. 2003. Regulation of calcium signals in the nucleus by a nucleoplasmic reticulum. *Nat Cell Biol* **5**: 440–446.
- Ehrlich BE, Kaftan E, Bezprozvannaya S, Bezprozvanny I. 1994. The pharmacology of intracellular Ca^{2+} -release channels. *Trends Pharmacol Sci* **15**: 145–149.
- Ehrlich BE, Watras J. 1988. Inositol 1,4,5-trisphosphate activates a channel from smooth muscle sarcoplasmic reticulum. *Nature* **336**: 583–586.
- Endo M. 1977. Calcium release from the sarcoplasmic reticulum. *Physiol Rev* **57**: 71–108.
- Endo M, Tanaka M, Ogawa Y. 1970. Calcium induced release of calcium from the sarcoplasmic reticulum of skinned skeletal muscle fibres. *Nature* **228**: 34–36.
- Fabiato A. 1983. Calcium-induced release of calcium from cardiac sarcoplasmic reticulum. *Am J Physiol* **245**: C1–C14.
- Ferris CD, Cameron AM, Haganir RL, Snyder SH. 1992. Quantal calcium release by purified reconstituted inositol 1,4,5-trisphosphate receptors. *Nature* **356**: 350–352.
- Ferris CD, Haganir RL, Supattapone S, Snyder SH. 1989. Purified inositol 1,4,5-trisphosphate receptor mediates calcium flux in reconstituted lipid vesicles. *Nature* **342**: 87–89.
- Finch EA, Turner TJ, Goldin SM. 1991. Calcium as a coagonist of inositol 1,4,5-trisphosphate-induced calcium release. *Science* **252**: 443–446.
- Foskett JK, White C, Cheung KH, Mak DO. 2007. Inositol trisphosphate receptor Ca^{2+} release channels. *Physiol Rev* **87**: 593–658.
- Fruen BR, Bardy JM, Byrem TM, Strasburg GM, Louis CF. 2000. Differential Ca^{2+} sensitivity of skeletal and cardiac muscle ryanodine receptors in the presence of calmodulin. *Am J Physiol* **279**: C724–C733.
- Furuichi T, Yoshikawa S, Miyawaki A, Wada K, Maeda M, Mikoshiba K. 1989. Primary structure and functional expression of the inositol 1,4,5-trisphosphate-binding protein P_{400} . *Nature* **342**: 32–38.
- Gafni J, Munsch JA, Lam TH, Catlin MC, Costa LG, Molinski TE, Pessah IN. 1997. Xestospongins: Potent membrane permeable blockers of the inositol 1,4,5-trisphosphate receptor. *Neuron* **19**: 723–733.
- Galvan DL, Borrego-Diaz E, Perez PJ, Mignery GA. 1999. Subunit oligomerization, and topology of the inositol 1,4,5-trisphosphate receptor. *J Biol Chem* **274**: 29483–29492.
- Gao L, Balshaw D, Xu L, Tripathy A, Xin C, Meissner G. 2000. Evidence for a role of the lumenal M3-M4 loop in skeletal muscle Ca^{2+} release channel (ryanodine receptor) activity and conductance. *Biophys J* **79**: 828–840.
- Gees M, Colsoul B, Nilius B. 2010. The Role of Transient Receptor Potential Cation Channels in Ca^{2+} Signaling. *Cold Spring Harb Perspect Biol* doi: 10.1101/cshperspect.a003962.
- Gerasimenko OV, Gerasimenko JV, Belan PV, Petersen OH. 1996. Inositol trisphosphate and cyclic ADP-ribose-mediated release of Ca^{2+} from single isolated pancreatic zymogen granules. *Cell* **84**: 473–480.
- Gerasimenko OV, Gerasimenko JV, Tepikin AV, Petersen OH. 1995. ATP-dependent accumulation and inositol trisphosphate- or cyclic ADP-ribose-mediated release of Ca^{2+} from the nuclear envelope. *Cell* **80**: 439–444.
- Gilkey JC. 1983. Roles of calcium and pH in activation of eggs of the medaka fish, *Oryzias latipes*. *J Cell Biol* **97**: 669–678.
- Gillespie D. 2008. Energetics of divalent selectivity in a calcium channel: the ryanodine receptor case study. *Biophys J* **94**: 1169–1184.
- Gillespie D, Fill M. 2008. Intracellular calcium release channels mediate their own countercurrent: the ryanodine receptor case study. *Biophys J* **95**: 3706–3714.
- Gnegy ME. 1993. Calmodulin in neurotransmitter and hormone action. *Annu Rev Pharmacol Toxicol* **33**: 45–70.
- Goto J, Suzuki AZ, Ozaki S, Matsumoto N, Nakamura T, Ebisui E, Fleig A, Penner R, Mikoshiba K. 2010. Two novel 2-aminoethyl diphenylborinate (2-APB) analogues differentially activate and inhibit store-operated Ca^{2+} entry via STIM proteins. *Cell Calcium* **47**: 1–10.
- Györke I, Györke S. 1998. Regulation of the cardiac ryanodine receptor channel by luminal Ca^{2+} involves luminal Ca^{2+} sensing sites. *Biophys J* **75**: 2801–2810.
- Hagar RE, Burgstahler AD, Nathanson MH, Ehrlich BE. 1998. Type III InsP_3 receptor channel stays open in the presence of increased calcium. *Nature* **296**: 81–84.

- Hajnóczky G, Thomas AP. 1997. Minimal requirements for calcium oscillations driven by the IP₃ receptor. *EMBO J* **16**: 3533–3543.
- Hamada K, Mikoshiba K. 2002. Two-state conformational changes in inositol 1,4,5-trisphosphate receptor regulated by calcium. *J Biol Chem* **277**: 21115–21118.
- Hamada K, Terauchi A, Mikoshiba K. 2003. Three-dimensional rearrangements with inositol 1,4,5-trisphosphate receptor by calcium. *J Biol Chem* **278**: 52881–52889.
- Hamilton SL. 2005. Ryanodine receptors. *Cell Calcium* **38**: 253–260.
- Haynes LP, Tepikin AV, Burgoyne RD. 2004. Calcium-binding protein 1 is an inhibitor of agonist-evoked, inositol 1,4,5-trisphosphate-mediated calcium signaling. *J Biol Chem* **279**: 547–555.
- Higo T, Hattori M, Nakamura T, Natsume T, Michikawa T, Mikoshiba K. 2005. Subtype-specific and ER luminal environment-dependent regulation of inositol 1,4,5-trisphosphate receptor type 1 by ERp44. *Cell* **120**: 85–98.
- Hirota J, Michikawa T, Miyawaki A, Furuichi T, Okura I, Mikoshiba K. 1995. Kinetics of calcium release by immunopurified inositol 1,4,5-trisphosphate receptor in reconstituted lipid vesicles. *J Biol Chem* **270**: 19046–19051.
- Hirota J, Michikawa T, Natsume T, Furuichi T, Mikoshiba K. 1999. Calmodulin inhibits inositol 1,4,5-trisphosphate-induced calcium release through the purified and reconstituted inositol 1,4,5-trisphosphate receptor type 1. *FEBS Lett* **456**: 322–326.
- Hokin MR, Hokin LE. 1953. Enzyme secretion and the incorporation of P³² into phospholipids of pancreas slices. *J Biol Chem* **203**: 967–977.
- Horne JH, Meyer T. 1995. Luminal calcium regulates the inositol trisphosphate receptor of rat basophilic leukemia cells at the cytosolic side. *Biochemistry* **34**: 12738–12746.
- Iino M. 1987. Calcium dependent inositol trisphosphate-induced calcium release in the guinea-pig taenia caeci. *Biochem Biophys Res Commun* **142**: 47–52.
- Iino M. 1990. Biphasic Ca²⁺ dependence of inositol 1,4,5-trisphosphate-induced Ca²⁺ release in smooth muscle cells of the guinea pig taenia caeci. *J Gen Physiol* **95**: 1103–1122.
- Imagawa T, Smith JS, Coronado R, Campbell KP. 1987. Purified ryanodine receptor from skeletal muscle sarcoplasmic reticulum is the Ca²⁺-permeable pore of the calcium release channel. *J Biol Chem* **262**: 16636–16643.
- Ionescu L, Cheung KH, Vais H, Mak DO, White C, Foskett JK. 2006. Graded recruitment and inactivation of single InsP₃ receptor Ca²⁺-release channels: implications for quantal Ca²⁺ release. *J Physiol* **573**: 645–662.
- Irvine RF. 1990. “Quantal” Ca²⁺ release and the control of Ca²⁺ entry by inositol phosphates - a possible mechanism. *FEBS Lett* **262**: 5–9.
- Islam MO, Yoshida Y, Koga T, Kojima M, Kanagawa K, Imai S. 1996. Isolation and characterization of vascular smooth muscle inositol 1,4,5-trisphosphate receptor. *Biochem J* **316**: 295–302.
- Iwai M, Michikawa T, Bosanac I, Ikura M, Mikoshiba K. 2007. Molecular basis of the isoform-specific ligand-binding affinity of inositol 1,4,5-trisphosphate receptors. *J Biol Chem* **282**: 12755–12764.
- Jean T, Klee CB. 1986. Calcium modulation of inositol 1,4,5-trisphosphate-induced calcium release from neuroblastoma x glioma hybrid (NG108-15) microsomes. *J Biol Chem* **261**: 16414–16420.
- Jiang D, Chen W, Xiao J, Wang R, Kong H, Jones PP, Zhang L, Fruen B, Chen SR. 2008. Reduced threshold for luminal Ca²⁺ activation of RyR1 underlies a causal mechanism of porcine malignant hyperthermia. *J Biol Chem* **283**: 20813–20820.
- Jiang Q-X, Thrower EC, Chester DW, Ehrlich BE, Sigworth FJ. 2002a. Three-dimensional structure of the type 1 inositol 1,4,5-trisphosphate receptor at 24 Å resolution. *EMBO J* **21**: 3575–3581.
- Jiang Y, Lee A, Chen J, Cadene M, Chait BT, MacKinnon R. 2002b. The open pore conformation of potassium channels. *Nature* **417**: 523–526.
- Jones PP, Meng X, Xiao B, Cai S, Bolstad J, Wagenknecht T, Liu Z, Chen SR. 2008. Localization of PKA phosphorylation site, Ser²⁰³⁰, in the three-dimensional structure of cardiac ryanodine receptor. *Biochem J* **410**: 261–270.
- Joseph SK, Brownell S, Khan MT. 2005. Calcium regulation of inositol 1,4,5-trisphosphate receptors. *Cell Calcium* **38**: 539–546.
- Joseph SK, Rice HL, Williamson JR. 1989. The effect of external calcium and pH on inositol trisphosphate-mediated calcium release from cerebellum microsomal fractions. *Biochem J* **258**: 261–265.
- Kaftan EJ, Ehrlich BE, Watras J. 1997. Inositol 1,4,5-trisphosphate (InsP₃) and calcium interact to increase the dynamic range of InsP₃ receptor-dependent calcium signaling. *J Gen Physiol* **110**: 529–538.
- Kiselyov K, Mignery GA, Zhu MX, Muallem S. 1999. The N-terminal domain of the IP₃ receptor gates store-operated hTrp3 channels. *Mol Cell* **4**: 423–429.
- Krinke O, Novotna Z, Valentova O, Martinek J. 2007. Inositol trisphosphate receptor in higher plants: is it real? *J Exp Bot* **58**: 361–376.
- Laude AJ, Tovey SC, Dedos S, Potter BVL, Lummis SCR, Taylor CW. 2005. Rapid functional assays of recombinant IP₃ receptors. *Cell Calcium* **38**: 45–51.
- Launikonis BS, Zhou J, Royer L, Shannon TR, Brum G, Rios E. 2006. Depletion “skraps” and dynamic buffering inside the cellular calcium store. *Proc Natl Acad Sci USA* **103**: 2982–2987.
- Laver DR. 2007. Ca²⁺ stores regulate ryanodine receptor Ca²⁺ release channels via luminal and cytosolic Ca²⁺ sites. *Biophys J* **92**: 3541–3555.
- Laver DR. 2009. Luminal Ca²⁺ activation of cardiac ryanodine receptors by luminal and cytoplasmic domains. *Eur J Biophys* **39**: 19–26.
- Lechleiter J, Girard S, Peralta E, Clapham D. 1991. Spiral calcium wave propagation and annihilation in *Xenopus laevis* oocytes. *Science* **252**: 123–126.
- Li P, Chen SR. 2001. Molecular basis of Ca²⁺ activation of the mouse cardiac Ca²⁺ release channel (ryanodine receptor). *J Gen Physiol* **118**: 33–44.
- Li W-H, Llopis J, Whitney M, Zlokarnik G, Tsien RY. 1998. Cell-permeant caged InsP₃ ester shows that Ca²⁺ spike frequency can optimize gene expression. *Nature* **392**: 936–941.

C.W. Taylor and S.C. Tovey

- Lin C, Widjaja J, Joseph SK. 2000. The interaction of calmodulin with alternatively spliced isoforms of the type-I inositol trisphosphate receptor. *J Biol Chem* **275**: 2305–2311.
- Lobo PA, Van Petegem F. 2009. Crystal structures of the N-terminal domains of cardiac and skeletal muscle ryanodine receptors: insights into disease mutations. *Structure* **17**: 1505–1514.
- Ludtke SJ, Serysheva II, Hamilton SL, Chiu W. 2005. The pore structure of the closed RYR1 channel. *Structure* **13**: 1203–1211.
- MacKinnon R. 2004. Potassium channels and the atomic basis of selective ion conduction (Nobel Lecture). *Angew Chem Int Edn Engl* **43**: 4265–4277.
- Maeda N, Niinobe M, Nakahira K, Mikoshiba K. 1988. Purification and characterization of P₄₀₀ protein, a glycoprotein characteristic of purkinje cell from mouse cerebellum. *J Neurochem* **51**: 1724–1730.
- Mak D-OD, McBride S, Foscett JK. 1998. Inositol 1,4,5-tris-phosphate activation of inositol tris-phosphate receptor Ca²⁺ channel by ligand tuning of Ca²⁺ inhibition. *Proc Natl Acad Sci USA* **95**: 15821–15825.
- Mak D-O, McBride S, Foscett JK. 2001. Regulation by Ca²⁺ and inositol 1,4,5-trisphosphate (InsP₃) of single recombinant type 3 InsP₃ receptor channels: Ca²⁺ activation uniquely distinguishes types 1 and 3 InsP₃ receptors. *J Gen Physiol* **117**: 435–446.
- Mak D-O, McBride SMJ, Petrenko NB, Foscett JK. 2003. Novel regulation of calcium inhibition of the inositol 1,4,5-trisphosphate receptor calcium-release channel. *J Gen Physiol* **122**: 569–581.
- Mak DO, McBride S, Raghuram V, Yue Y, Joseph SK, Foscett JK. 2000. Single-channel properties in endoplasmic reticulum membrane of recombinant type 3 inositol trisphosphate receptor. *J Gen Physiol* **115**: 241–256.
- Marchant JS, Taylor CW. 1997. Cooperative activation of IP₃ receptors by sequential binding of IP₃ and Ca²⁺ safeguards against spontaneous activity. *Curr Biol* **7**: 510–518.
- Marchenko SM, Yarotsky VV, Kovalenko TN, Kostyuk PG, Thomas RC. 2005. Spontaneously active and InsP₃-activated ion channels in cell nuclei from rat cerebellar Purkinje and granule neurones. *J Physiol* **565**: 897–910.
- Marshall ICB, Taylor CW. 1993. Biphasic effects of cytosolic calcium on Ins(1,4,5)P₃-stimulated Ca²⁺ mobilization in hepatocytes. *J Biol Chem* **268**: 13214–13220.
- Marshall ICB, Taylor CW. 1994. Two calcium-binding sites mediate the interconversion of liver inositol 1,4,5-trisphosphate receptors between three conformational states. *Biochem J* **301**: 591–598.
- Maruyama T, Kanaji T, Nakade S, Kanno T, Mikoshiba K. 1997. 2APB, 2-aminoethoxydiphenyl borate, a membrane-penetrable modulator of Ins(1,4,5)P₃-induced Ca²⁺ release. *J Biochem* **122**: 498–505.
- Mayer ML. 2006. Glutamate receptors at atomic resolution. *Nature* **440**: 456–462.
- Meur G, Parker AKT, Gergely FV, Taylor CW. 2007. Targeting and retention of type 1 ryanodine receptors to the endoplasmic reticulum. *J Biol Chem* **282**: 23096–23103.
- Meyer T, Stryer L. 1990. Transient calcium release induced by successive increments of inositol 1,4,5-trisphosphate. *Proc Natl Acad Sci USA* **87**: 3841–3845.
- Michelangeli F, Mezna M, Tovey S, Sayers LG. 1995. Pharmacological modulators of the inositol 1,4,5-trisphosphate receptor. *Neuropharmacol* **34**: 1111–1122.
- Michell RH. 1975. Inositol phospholipids and cell surface receptor function. *Biochim Biophys Acta* **415**: 81–147.
- Michell RH, Drummond AH, Downes CP. 1989. *Inositol Lipids in Cell Signalling*. Academic Press Ltd, London.
- Michikawa T, Hirota J, Kawano S, Hiraoka M, Yamada M, Furuichi T, Mikoshiba K. 1999. Calmodulin mediates calcium-dependent inactivation of the cerebellar type 1 inositol 1,4,5-trisphosphate receptor. *Neuron* **23**: 799–808.
- Mignery GA, Südhof TC, Takei K, De Camilli P. 1989. Putative receptor for inositol 1,4,5-trisphosphate similar to ryanodine receptor. *Nature* **342**: 192–195.
- Missiaen L, De Smedt H, Bultynck G, Vanlingen S, De Smet P, Callewaert G, Parys J. 2000. Calmodulin increases the sensitivity of type 3 inositol 1,4,5-trisphosphate receptors to Ca²⁺ inhibition in human bronchial mucosal cells. *Mol Pharmacol* **57**: 564–567.
- Missiaen L, De Smedt H, Parys JB, Casteels R. 1994. Co-activation of inositol trisphosphate-induced Ca²⁺ release by cytosolic Ca²⁺ is loading-dependent. *J Biol Chem* **269**: 7238–7242.
- Missiaen L, Parys JB, Sienaert I, Maes K, Kunzelmann K, Takahashi M, Tanzawa K, De Smedt H. 1998. Functional properties of the type-3 InsP₃ receptor in 16HBE140-bronchial mucosal cells. *J Biol Chem* **273**: 8983–8986.
- Missiaen L, Parys JB, Weidema AF, Sipma H, Vanlingen S, De Smet P, Callewaert G, De Smedt H. 1999. The bell-shaped Ca²⁺-dependence of the inositol 1,4,5-trisphosphate induced Ca²⁺ release is modulated by Ca²⁺/calmodulin. *J Biol Chem* **274**: 13748–13751.
- Missiaen L, Taylor CW, Berridge MJ. 1992. Luminal Ca²⁺ promoting spontaneous Ca²⁺ release from inositol trisphosphate-sensitive stores of rat hepatocytes. *J Physiol* **455**: 623–640.
- Miyakawa T, Maeda A, Yamazawa T, Hirose K, Kurosaki T, Iino M. 1999. Encoding of Ca²⁺ signals by differential expression of IP₃ receptor subtypes. *EMBO J* **18**: 1303–1308.
- Miyakawa T, Mizushima A, Hirose K, Yamazawa T, Bezprozvanny I, Kurosaki T, Iino M. 2001. Ca²⁺-sensor region of IP₃ receptor controls intracellular Ca²⁺ signaling. *EMBO J* **20**: 1674–1680.
- Mohr FC, Hershey PEC, Zimányi I, Pessah IN. 1993. Regulation of inositol 1,4,5-trisphosphate receptors in rat basophilic leukemia cells. I. Multiple conformational states of the receptor in a microsomal preparation. *Biochim Biophys Acta* **1147**: 105–114.
- Muallem S, Pandol SJ, Beeker TG. 1989. Hormone-evoked calcium release from intracellular stores is a quantal process. *J Biol Chem* **264**: 205–212.
- Nadif Kasri N, Bultynck G, Sienaert I, Callewaert G, Erneux C, Missiaen L, Parys JB, De Smedt H. 2002. The role of calmodulin for inositol 1,4,5-trisphosphate receptor function. *Biochim Biophys Acta* **1600**: 19–31.

- Nadif Kasri N, Holmes AM, Bultynck G, Parys JB, Bootman MD, Rietdorf K, Missiaen L, McDonald F, De Smedt H, Conway SJ, et al. 2004. Regulation of InsP₃ receptor activity by neuronal Ca²⁺-binding proteins. *EMBO J* **23**: 312–321.
- Nadif Kasri N, Torok K, Galione A, Garnham C, Callewaert G, Missiaen L, Parys JB, De Smedt H. 2006. Endogenously bound calmodulin is essential for the function of the inositol 1,4,5-trisphosphate receptor. *J Biol Chem* **281**: 8332–8338.
- Nilius B, Owsianik G, Voets T, Peters JA. 2007. Transient receptor potential cation channels in disease. *Physiol Rev* **87**: 165–217.
- Nosyreva E, Miyakawa T, Wang Z, Glouchankova L, Iino M, Bezprozvanny I. 2002. The high-affinity calcium-calmodulin-binding site does not play a role in the modulation of type 1 inositol 1,4,5-trisphosphate receptor function by calcium and calmodulin. *Biochem J* **365**: 659–667.
- Nunn DL, Taylor CW. 1992. Luminal Ca²⁺ increases the sensitivity of Ca²⁺ stores to inositol 1,4,5-trisphosphate. *Mol Pharmacol* **41**: 115–119.
- Oancea E, Meyer T. 1996. Reversible desensitization of inositol trisphosphate-induced calcium release provides a mechanism for repetitive calcium spikes. *J Biol Chem* **271**: 17253–17260.
- Oda T, Yano M, Yamamoto T, Tokuhisa T, Okuda S, Doi M, Ohkusa T, Ikeda Y, Kobayashi S, Ikemoto N, et al. 2005. Defective regulation of interdomain interactions within the ryanodine receptor plays a key role in the pathogenesis of heart failure. *Circulation* **111**: 3400–3410.
- Oldershaw KA, Nunn DL, Taylor CW. 1991. Quantal Ca²⁺ mobilization stimulated by inositol 1,4,5-trisphosphate in permeabilized hepatocytes. *Biochem J* **278**: 705–708.
- Oldershaw KA, Taylor CW. 1993. Luminal Ca²⁺ increases the affinity of inositol 1,4,5-trisphosphate for its receptor. *Biochem J* **292**: 631–633.
- Orlova EV, Serysheva II, van Heel M, Hamilton SL, Chiu W. 1996. Two structural configurations of the skeletal muscle calcium release channel. *Nat Struct Biol* **3**: 547–552.
- Pantazaka E, Taylor CW. 2010. Targeting of inositol 1,4,5-trisphosphate receptor to the endoplasmic reticulum by its first transmembrane domain. *Biochem J* **425**: 61–69.
- Park CY, Hoover PJ, Mullins FM, Bachhawat P, Covington ED, Raunser S, Walz T, Garcia KC, Dolmetsch RE, Lewis RS. 2009. STIM1 clusters and activates CRAC channels via direct binding of a cytosolic domain to Orai1. *Cell* **136**: 876–890.
- Parker AKT, Gergely FV, Taylor CW. 2004. Targeting of inositol 1,4,5-trisphosphate receptors to the endoplasmic reticulum by multiple signals within their transmembrane domains. *J Biol Chem* **279**: 23797–23805.
- Parker I, Choi J, Yao Y. 1996. Elementary events of InsP₃-induced Ca²⁺ liberation in *Xenopus* oocytes: hot spots, puffs and blips. *Cell Calcium* **20**: 105–121.
- Parker I, Ivorra I. 1990. Localized all-or-nothing calcium liberation by inositol trisphosphate. *Science* **250**: 977–979.
- Parker I, Ivorra I. 1991. Caffeine inhibits inositol trisphosphate-mediated liberation of intracellular calcium in *Xenopus* oocytes. *J Physiol* **433**: 229–240.
- Parys JB, Missiaen L, De Smedt H, Casteels R. 1993. Loading dependence of inositol 1,4,5-trisphosphate-induced Ca²⁺ release in the clonal cell line A7r5. *J Biol Chem* **268**: 25206–25212.
- Parys JB, Sernett SW, DeLisle S, Snyder PM, Welsh MJ, Campbell KP. 1992. Isolation, characterization, and localization of the inositol 1,4,5-trisphosphate receptor protein in *Xenopus laevis* oocytes. *J Biol Chem* **267**: 18776–18782.
- Patel S, Marchant JS, Brailoiu E. 2010. Two-pore channels: regulation by NAADP and customized roles in triggering calcium signals. *Cell Calcium* **47**: 480–490.
- Patel S, Morris SA, Adkins CE, O’Beirne G, Taylor CW. 1997. Ca²⁺-independent inhibition of inositol trisphosphate receptors by calmodulin: Redistribution of calmodulin as a possible means of regulating Ca²⁺ mobilization. *Proc Natl Acad Sci USA* **94**: 11627–11632.
- Patterson RL, Boehning D, Snyder SH. 2004. Inositol 1,4,5-trisphosphate receptors as signal integrators. *Annu Rev Biochem* **73**: 437–465.
- Petersen OH, Burdakov D, Tepikin AV. 1999. Polarity in intracellular calcium signalling. *Bioessays* **21**: 851–860.
- Pietri F, Hilly M, Mauger J-P. 1990. Calcium mediates the interconversion between two states of the liver inositol 1,4,5-trisphosphate receptor. *J Biol Chem* **265**: 17478–17485.
- Pinton P, Pozzan T, Rizzuto R. 1998. The Golgi apparatus is an inositol 1,4,5-trisphosphate-sensitive Ca²⁺ store, with functional properties distinct from those of the endoplasmic reticulum. *EMBO J* **17**: 5298–5308.
- Putney JW Jr. 1997. *Capacitative calcium entry*. RGLandes Company, Austin, Texas, USA.
- Putney JW, Poggioli J, Weiss SJ. 1981. Receptor regulation of calcium release and calcium permeability in parotid gland cells. *Philos Trans R Soc London [Biol]* **296**: 37–45.
- Rahman T-U, Skupin A, Falcke M, Taylor CW. 2009. Clustering of IP₃ receptors by IP₃ retunes their regulation by IP₃ and Ca²⁺. *Nature* **458**: 655–659.
- Rahman T, Taylor CW. 2009. Dynamic regulation of IP₃ receptor clustering and activity by IP₃. *Channels* **3**: 336–332.
- Ramos-Franco J, Bare D, Caenepeel S, Nani A, Fill M, Mignery G. 2000. Single-channel function of recombinant type 2 inositol 1,4,5-trisphosphate receptor. *Biophys J* **79**: 1388–1399.
- Ramos-Franco J, Caenepeel S, Fill M, Mignery G. 1998a. Single channel function of recombinant type-1 inositol 1,4,5-trisphosphate receptor ligand binding domain splice variants. *Biophys J* **75**: 2783–2793.
- Ramos-Franco J, Fill M, Mignery GA. 1998b. Isoform-specific function of single inositol 1,4,5-trisphosphate receptor channels. *Biophys J* **75**: 834–839.
- Ramos-Franco J, Galvan D, Mignery GA, Fill M. 1999. Location of the permeation pathway in the recombinant type-1 inositol 1,4,5-trisphosphate receptor. *J Gen Physiol* **114**: 243–250.
- Rasmussen H. 1970. Cell communication, calcium ion, and cyclic adenosine monophosphate. *Science* **170**: 404–412.
- Richardson A, Taylor CW. 1993. Effects of Ca²⁺ chelators on purified inositol 1,4,5-trisphosphate (InsP₃) receptors

C.W. Taylor and S.C. Tovey

- and InsP₃-stimulated Ca²⁺ mobilization. *J Biol Chem* **268**: 11528–11533.
- Ridgway EB, Gilkey JC, Jaffe LF. 1977. Free calcium increases explosively in activating medaka eggs. *Proc Natl Acad Sci USA* **74**: 623–627.
- Ringer S. 1883. A further contribution regarding the influence of the different constituents of the blood on the contraction of the heart. *J Physiol* **4**: 29–42.
- Rizzuto R, Brini M, Murgia M, Pozzan T. 1993. Microdomains with high Ca²⁺ close to IP₃-sensitive channels that are sensed by neighbouring mitochondria. *Science* **262**: 744–747.
- Rodney GG, Moore CP, Williams BY, Zhang J-Z, Krol J, Pedersen SE, Hamilton SL. 2001. Calcium binding to calmodulin leads to an N-terminal shift in its binding site on the ryanodine receptor. *J Biol Chem* **276**: 2069–2074.
- Rossi AM, Riley AM, Potter BVL, Taylor CW. 2010. Adenophostins: high-affinity agonists of IP₃ receptors. *Curr Topics Membr* **66**: 209–233.
- Rossi AM, Riley AM, Tovey SC, Rahman T, Dellis O, Taylor EJA, Veresov VG, Potter BVL, Taylor CW. 2009. Synthetic partial agonists reveal key steps in IP₃ receptor activation. *Nat Chem Biol* **5**: 631–639.
- Samso M, Feng W, Pessah IN, Allen PD. 2009. Coordinated movement of cytoplasmic and transmembrane domains of RyR1 upon gating. *PLoS Biol* **7**: e85.
- Samso M, Wagenknecht T, Allen PD. 2005. Internal structure and visualization of transmembrane domains of the RyR1 calcium release channel by cryo-EM. *Nat Struct Mol Biol* **6**: 539–544.
- Sato C, Hamada K, Ogura T, Miyazawa A, Iwasaki K, Hiroaki Y, Tani K, Terauchi A, Fujiyoshi Y, Mikoshiba K. 2004. Inositol 1,4,5-trisphosphate receptor contains multiple cavities and L-shaped ligand-binding domains. *J Mol Biol* **336**: 155–164.
- Schug ZT, da Fonseca PC, Bhanumathy CD, Wagner L 2nd, Zhang X, Bailey B, Morris EP, Yule DI, Joseph SK. 2008. Molecular characterization of the inositol 1,4,5-trisphosphate receptor pore-forming segment. *J Biol Chem* **283**: 2939–2948.
- Schug ZT, Joseph SK. 2006. The role of the S4-S5 linker and C-terminal tail in inositol 1,4,5-trisphosphate receptor function. *J Biol Chem* **281**: 24431–24440.
- Serysheva II, Bare DJ, Ludtke SJ, Kettlun CS, Chiu W, Mignery GA. 2003. Structure of the type 1 inositol 1,4,5-trisphosphate receptor revealed by cryomicroscopy. *J Biol Chem* **278**: 21319–21322.
- Serysheva II, Hamilton SL, Chiu W, Ludtke SJ. 2005. Structure of a Ca²⁺ release channel at 14 Å resolution. *J Struct Biol* **345**: 427–431.
- Shuttleworth TJ. 1992. Ca²⁺ release from inositol trisphosphate-sensitive stores is not modulated by intraluminal [Ca²⁺]. *J Biol Chem* **267**: 3573–3576.
- Sienaert I, De Smedt H, Parys JB, Missiaen L, Vanlingen S, Sipma H, Casteels R. 1996. Characterization of a cytosolic and a luminal Ca²⁺ binding site in the type I inositol 1,4,5-trisphosphate receptor. *J Biol Chem* **271**: 27005–27012.
- Sienaert I, Kasri NN, Vanlingen S, Parys J, Callewaert G, Missiaen L, De Smedt H. 2002. Localization and function of a calmodulin/apocalmodulin binding domain in the N-terminal part of the type I inositol 1,4,5-trisphosphate receptor. *Biochem J* **365**: 269–277.
- Sienaert I, Missiaen L, De Smedt H, Parys JB, Sipma H, Casteels R. 1997. Molecular and functional evidence for multiple Ca²⁺-binding domains on the type I inositol 1,4,5-trisphosphate receptor. *J Biol Chem* **272**: 25899–25906.
- Smith JS, Coronado R, Meissner G. 1985. Sarcoplasmic reticulum contains adenine nucleotide-activated calcium channels. *Nature* **316**: 446–449.
- Solovyova N, Fernyhough P, Glazner G, Verkhratsky A. 2002. Xestospongins empty the ER calcium store but do not inhibit InsP₃-induced Ca²⁺ release in cultured dorsal root ganglia neurones. *Cell Calcium* **32**: 49–52.
- Spät A, Bradford PG, McKinney JS, Rubin RP, Putney JW Jr. 1986. A saturable receptor for ³²P-inositol-1,4,5-trisphosphate in hepatocytes and neutrophils. *Nature* **319**: 514–516.
- Stehno-Bittel L, Lückhoff A, Clapham DE. 1995. Calcium release from the nucleus by InsP₃ receptor channels. *Neuron* **14**: 163–167.
- Streb H, Irvine RF, Berridge MJ, Schulz I. 1983. Release of Ca²⁺ from a nonmitochondrial store of pancreatic acinar cells by inositol-1,4,5-trisphosphate. *Nature* **306**: 67–69.
- Strigrow F, Ehrlich BE. 1996. The inositol 1,4,5-trisphosphate receptor of cerebellum. Mn²⁺ permeability and regulation by cytosolic Mn²⁺. *J Gen Physiol* **108**: 115–124.
- Suematsu E, Hirata M, Hashimoto T, Kuriyama H. 1984. Inositol 1,4,5-trisphosphate releases Ca²⁺ from intracellular store sites in skinned single cells of porcine coronary artery. *Biochem Biophys Res Commun* **120**: 481–485.
- Sun Y, Taylor CW. 2008. A calmodulin antagonist reveals a calmodulin-independent interdomain interaction essential for activation of inositol 1,4,5-trisphosphate receptors. *Biochem J* **416**: 243–253.
- Supattapone S, Worley PF, Baraban JM, Snyder SH. 1988. Solubilization, purification, and characterization of an inositol trisphosphate receptor. *J Biol Chem* **263**: 1530–1534.
- Sureshan KM, Riley AM, Rossi AM, Tovey SC, Dedos SG, Taylor CW, Potter BVL. 2009. Activation of IP₃ receptors by synthetic bisphosphate ligands. *Chem Commun* **14**: 1204–1206.
- Sutko JL, Ito K, Kenyon JL. 1985. Ryanodine: a modifier of sarcoplasmic reticulum calcium release in striated muscle. *FASEB J* **44**: 2984–2988.
- Swatton JE, Morris SA, Cardy TJA, Taylor CW. 1999. Type 3 inositol trisphosphate receptors in RINm5F cells are biphasically regulated by cytosolic Ca²⁺ and mediate quantal Ca²⁺ mobilization. *Biochem J* **344**: 55–60.
- Swatton JE, Taylor CW. 2002. Fast biphasic regulation of type 3 inositol trisphosphate receptors by cytosolic calcium. *J Biol Chem* **277**: 17571–17579.
- Szłufcik K, Bultynck G, Callewaert G, Missiaen L, Parys JB, De Smedt H. 2006. The suppressor domain of inositol 1,4,5-trisphosphate receptor plays an essential role in the protection against apoptosis. *Cell Calcium* **39**: 325–336.
- Takeshima H, Nishimura S, Matsumoto T, Ishida H, Kangawa K, Minamino N, Matsuo H, Ueda M, Hanaoka

- M, Hirose T, et al. 1989. Primary structure and expression from complementary DNA of skeletal muscle ryanodine receptor. *Nature* **339**: 439–445.
- Tanimura A, Turner RJ. 1996. Calcium release in HSY cells conforms to a steady-state mechanism involving regulation of the inositol 1,4,5-trisphosphate receptor Ca^{2+} channel by luminal $[\text{Ca}^{2+}]$. *J Cell Biol* **132**: 607–616.
- Tateishi H, Yano M, Mochizuki M, Suetomi T, Ono M, Xu X, Uchinoumi H, Okuda S, Oda T, Kobayashi S, et al. 2009. Defective domain-domain interactions within the ryanodine receptor as a critical cause of diastolic Ca^{2+} leak in failing hearts. *Cardiovasc Res* **81**: 536–545.
- Taylor CW, da Fonseca PCA, Morris EP. 2004. IP₃ receptors: the search for structure. *Trends Biochem Sci* **29**: 210–219.
- Taylor CW, Genazzani AA, Morris SA. 1999. Expression of inositol trisphosphate receptors. *Cell Calcium* **26**: 237–251.
- Taylor CW, Laude AJ. 2002. IP₃ receptors and their regulation by calmodulin and cytosolic Ca^{2+} . *Cell Calcium* **32**: 321–334.
- Taylor CW, Potter BVL. 1990. The size of inositol 1,4,5-trisphosphate-sensitive Ca^{2+} stores depends on inositol 1,4,5-trisphosphate concentration. *Biochem J* **266**: 189–194.
- Taylor CW, Prole DL, Rahman T. 2009a. Ca^{2+} channels on the move. *Biochemistry* **48**: 12062–12080.
- Taylor CW, Ur-Rahman T, Pantazaka E. 2009b. Targeting and clustering of IP₃ receptors: key determinants of spatially organized Ca^{2+} signals. *Chaos* **19**: 037102–037101–037102–037110.
- Terentyev D, Nori A, Santoro M, Viatchenko-Karpinski S, Kubalova Z, Gyorko I, Terentyeva R, Vedamoorthyrao S, Blom NA, Valle G, et al. 2006. Abnormal interactions of calsequestrin with the ryanodine receptor calcium release channel complex linked to exercise-induced sudden cardiac death. *Circ Res* **98**: 1151–1158.
- Thrower EC, Lea EJA, Dawson AP. 1998. The effects of free $[\text{Ca}^{2+}]$ on the cytosolic face of inositol (1,4,5)-trisphosphate receptor at the single channel level. *Biochem J* **330**: 559–564.
- Tovey SC, Dedos SG, Rahman T, Taylor EJA, Pantazaka E, Taylor CW. 2010. Regulation of inositol 1,4,5-trisphosphate receptors by cAMP independent of cAMP-dependent protein kinase. *J Biol Chem* **285**: 12979–12989.
- Tripathy A, Meissner G. 1996. Sarcoplasmic reticulum luminal Ca^{2+} has access to cytosolic activation and inactivation sites of skeletal muscle Ca^{2+} release channel. *Biophys J* **70**: 2600–2615.
- Tsien RY. 1980. New calcium indicators and buffers with high selectivity against magnesium and protons: design, synthesis, and properties of prototype structures. *Biochemistry* **19**: 2396–2404.
- Tsien RY. 1981. A non-disruptive technique for loading calcium buffers and indicators into cells. *Nature* **290**: 527–528.
- Tu H, Miyakawa T, Wang Z, Glouchankova L, Iino M, Bezprozvanny I. 2002. Functional characterization of the type 1 inositol 1,4,5-trisphosphate receptor coupling domain SII(±) splice variants and the *Opisthotonos* mutant form. *Biophys J* **82**: 1995–2004.
- Tu H, Nosyreva E, Miyakawa T, Wang Z, Mizushima A, Iino M, Bezprozvanny I. 2003. Functional and biochemical analysis of the type 1 inositol (1,4,5)-trisphosphate receptor calcium sensor. *Biophys J* **85**: 290–299.
- Tu H, Wang Z, Bezprozvanny I. 2005a. Modulation of mammalian inositol 1,4,5-trisphosphate receptor isoforms by calcium: a role of calcium sensor region. *Biophys J* **88**: 1056–1069.
- Tu H, Wang Z, Nosyreva E, De Smedt H, Bezprozvanny I. 2005b. Functional characterization of mammalian inositol 1,4,5-trisphosphate receptor isoforms. *Biophys J* **88**: 1046–1055.
- Uchida K, Miyauchi H, Furuichi T, Michikawa T, Mikoshiba K. 2003. Critical regions for activation gating of the inositol 1,4,5-trisphosphate receptor. *J Biol Chem* **278**: 16551–16560.
- van de Put FHMM, De Pont JHHM, Willems PHGM. 1994. Heterogeneity between intracellular Ca^{2+} stores as the underlying principle of quantal Ca^{2+} release by inositol 1,4,5-trisphosphate in permeabilized pancreatic acinar cells. *J Biol Chem* **269**: 12438–12443.
- Varney MA, Rivera J, Lopez Bernal A, Watson SP. 1990. Are there subtypes of the inositol 1,4,5-trisphosphate receptor? *Biochem J* **269**: 211–216.
- Wang R, Chen W, Cai S, Zhang J, Bolstad J, Wagenknecht T, Liu Z, Chen SR. 2007. Localization of an NH_2 -terminal disease-causing mutation hot spot to the “clamp” region in the three-dimensional structure of the cardiac ryanodine receptor. *J Biol Chem* **282**: 17785–17793.
- Wang R, Zhang L, Bolstad J, Diao N, Brown C, Ruest L, Welch W, Williams AJ, Chen SR. 2003. Residue Gln4863 within a predicted transmembrane sequence of the Ca^{2+} release channel (ryanodine receptor) is critical for ryanodine interaction. *J Biol Chem* **278**: 51557–51565.
- Wang Y, Xu L, Pasek DA, Gillespie D, Meissner G. 2005. Probing the role of negatively charged amino acid residues in ion permeation of skeletal muscle ryanodine receptor. *Biophys J* **89**: 256–265.
- Watras J, Bezprozvanny I, Ehrlich BE. 1991. Inositol 1,4,5-trisphosphate-gated channels in cerebellum: presence of multiple conductance states. *J Neurosci* **11**: 3239–3245.
- Williams AJ, West DJ, Sitsapesan R. 2001. Light at the end of the Ca^{2+} -release channel tunnel: structures and mechanisms involved in ion translocation in ryanodine receptor channels. *Quart Rev Biophys* **34**: 61–104.
- Williams JA. 1980. Regulation of pancreatic acinar cell function by intracellular calcium. *Am J Physiol* **238**: G269–G279.
- Willoughby D, Cooper DM. 2007. Organization and Ca^{2+} regulation of adenylyl cyclases in cAMP microdomains. *Physiol Rev* **87**: 965–1010.
- Wolfram F, Morris E, Taylor CW. 2010. Three-dimensional structure of recombinant type 1 inositol 1,4,5-trisphosphate receptor. *Biochem J* **428**: 483–489.
- Woods NM, Cuthbertson KSR, Cobbold PH. 1986. Repetitive transient rises in cytoplasmic free calcium in hormone-stimulated hepatocytes. *Nature* **319**: 600–602.
- Worley PF, Baraban JM, Supattapone S, Wilson VS, Snyder SH. 1987. Characterization of inositol trisphosphate

C.W. Taylor and S.C. Tovey

- receptor binding in brain. Regulation by pH and calcium. *J Biol Chem* **262**: 12132–12136.
- Yamada M, Miyawaki A, Saito K, Yamamoto-Hino M, Ryo Y, Furuichi T, Mikoshiba K. 1995. The calmodulin-binding domain in the mouse type 1 inositol 1,4,5-trisphosphate receptor. *Biochem J* **308**: 83–88.
- Yang J, McBride S, Mak D-OD, Vardi N, Palczewski K, Haeseleer F, Foscett JK. 2002. Identification of a family of calcium sensors as protein ligands of the inositol trisphosphate receptor Ca^{2+} release channels. *Proc Natl Acad Sci USA* **99**: 7711–7716.
- Yoneshima H, Miyawaki A, Michikawa T, Furuichi T, Mikoshiba K. 1997. Ca^{2+} differentially regulates ligand-affinity states of type 1 and 3 inositol 1,4,5-trisphosphate receptors. *Biochem J* **322**: 591–596.
- Zeng W, Mak DD, Li Q, Shin DM, Foscett JK, Muallem S. 2003. A new mode of Ca^{2+} signaling by G protein-coupled receptors: gating of IP_3 receptor Ca^{2+} release channels by $\text{G}\beta\gamma$. *Curr Biol* **13**: 872–876.
- Zhang X, Joseph SK. 2001. Effect of mutation of a calmodulin-binding sites on Ca^{2+} regulation of inositol trisphosphate receptors. *Biochem J* **360**: 395–400.
- Zhu MX, Ma J, Parrington J, Calcraft PJ, Galione A, Evans AM. 2010. Calcium signaling via two-pore channels: local or global, that is the question. *Am J Physiol* **298**: C430–C441.

Cold Spring Harbor Perspectives in Biology

IP₃ Receptors: Toward Understanding Their Activation

Colin W. Taylor and Stephen C. Tovey

Cold Spring Harb Perspect Biol 2010; doi: 10.1101/cshperspect.a004010 originally published online October 27, 2010

Subject Collection [Calcium Signaling](#)

The Endoplasmic Reticulum–Plasma Membrane Junction: A Hub for Agonist Regulation of Ca²⁺ Entry

Hwei Ling Ong and Indu Suresh Ambudkar

Calcium-Handling Defects and Neurodegenerative Disease

Sean Schrank, Nikki Barrington and Grace E. Stutzmann

Lysosomal Ca²⁺ Homeostasis and Signaling in Health and Disease

Emyr Lloyd-Evans and Helen Waller-Evans

Ca²⁺ Signaling in Exocrine Cells

Malini Ahuja, Woo Young Chung, Wei-Yin Lin, et al.

Functional Consequences of Calcium-Dependent Synapse-to-Nucleus Communication: Focus on Transcription-Dependent Metabolic Plasticity

Anna M. Hagenston, Hilmar Bading and Carlos Bas-Orth

Identifying New Substrates and Functions for an Old Enzyme: Calcineurin

Jagoree Roy and Martha S. Cyert

Fundamentals of Cellular Calcium Signaling: A Primer

Martin D. Bootman and Geert Bultynck

Primary Active Ca²⁺ Transport Systems in Health and Disease

Jialin Chen, Aljona Sitsel, Veronick Benoy, et al.

Signaling through Ca²⁺ Microdomains from Store-Operated CRAC Channels

Pradeep Barak and Anant B. Parekh

Structural Insights into the Regulation of Ca²⁺/Calmodulin-Dependent Protein Kinase II (CaMKII)

Moitrayee Bhattacharyya, Deepti Karandur and John Kuriyan

Store-Operated Calcium Channels: From Function to Structure and Back Again

Richard S. Lewis

Bcl-2-Protein Family as Modulators of IP₃ Receptors and Other Organellar Ca²⁺ Channels

Hristina Ivanova, Tim Vervliet, Giovanni Monaco, et al.

Calcium Signaling in Cardiomyocyte Function

Guillaume Gilbert, Kateryna Demydenko, Eef Dries, et al.

Cytosolic Ca²⁺ Buffers Are Inherently Ca²⁺ Signal Modulators

Beat Schwaller

For additional articles in this collection, see <http://cshperspectives.cshlp.org/cgi/collection/>

**All Modifications and
Oligo Types Synthesized**

Long Oligos • Fluorescent • Chimeric • DNA • RNA • Antisense

Oligo Modifications?
Your wish is our command.

Role of Two-Pore Channels in Embryonic Development and Cellular Differentiation

Sarah E. Webb, Jeffrey J. Kelu and Andrew L. Miller

Organellar Calcium Handling in the Cellular Reticular Network

Wen-An Wang, Luis B. Agellon and Marek Michalak

For additional articles in this collection, see <http://cshperspectives.cshlp.org/cgi/collection/>

**All Modifications and
Oligo Types Synthesized**

Long Oligos • Fluorescent • Chimeric • DNA • RNA • Antisense

Oligo Modifications?
Your wish is our command.