mTOR Signaling

Mathieu Laplante and David M. Sabatini

Centre de Recherche de l'Institut Universitaire de Cardiologie et de Pneumologie de Québec (CRIUCPQ), Université Laval, Québec G1V 4G5, Canada

Correspondence: mathieu.laplante@criucpq.ulaval.ca

The scarcity of nutrients/energy interspersed with sporadic periods of abundance means that cells must transition between anabolic and catabolic states. An important protein that has evolved to respond to this need is the target of rapamycin (TOR). TOR is a well-conserved serine/threonine kinase that plays an important role in the signaling network that controls growth and metabolism in response to environmental cues. As its name indicates, TOR

is the target of a molecule named rapamycin, an anti-fungal macrolide produced by the bacterial species *Streptomyces hygroscopicus*, which was isolated from a soil sample from the Eastern Islands in the 1970s (Vezina et al. 1975). In addition to its anti-fungal properties, rapamycin strongly inhibits cell growth and proliferation, making this molecule a valuable tool to study cell growth control. In the early 1990s, yeast genetic screens led to the identification of

Figure 1. A simplified view of the mTOR pathway.

Editors: Lewis Cantley, Tony Hunter, Richard Sever, and Jeremy W. Thorner
Additional Perspectives on Signal Transduction available at www.cshperspectives.org

Copyright © 2012 Cold Spring Harbor Laboratory Press; all rights reserved.
Advanced Online Article. Cite this article as *Cold Spring Harb Perspect Biol* doi: 10.1101/cshperspect.a011593

TOR as one mediator of the toxic effect of rapamycin in yeast (Heitman et al. 1991; Kunz et al. 1993). Shortly after, the mammalian TOR (mTOR), now officially known as the mechanistic TOR, was identified as the physical target of rapamycin (Fig. 1) (Brown et al. 1994; Sabatini et al. 1994; Sabers et al. 1995).

mTOR interacts with many proteins to form at least two distinct multiprotein complexes: mTOR complex 1 (mTORC1) and mTOR complex 2 (mTORC2) (Zoncu et al. 2011). In addition to their different protein compositions, the mTOR complexes have important differences in their sensitivities to rapamycin, in the upstream signals

Figure 2. The mTOR pathway.

they integrate, in the substrates they regulate, and in the biological processes they control (Laplanche and Sabatini 2009b). mTORC1 is sensitive to rapamycin and integrates inputs from at least five major signals—growth factors, genotoxic stress, energy status, oxygen, and amino acids—to regulate many processes involved in the promotion of cell growth and proliferation (Fig. 2). With the exception of amino acids, all of these inputs regulate mTORC1 activity by modulating the activity of TSC1–TSC2. TSC1–TSC2 negatively regulates mTORC1 activity by converting Rheb into its inactive GDP-bound state. Growth factors block TSC1–TSC2 activity and activate mTORC1, whereas genotoxic stress, energy deficit, and oxygen deprivation promote TSC1–TSC2 action and inhibit mTORC1. Conversely, amino acids activate mTORC1 through the action of the Rag GTPases. In the presence of amino acids, the Rag GTPases interact with mTORC1, which promotes its translocation from the cytoplasm to lysosomal membranes, where Rheb is thought to reside.

When activated, mTORC1 promotes protein synthesis mainly by phosphorylating the kinase S6K and the translation regulator 4E-BP1 (Ma and Blenis 2009). This complex also induces lipid biogenesis by activating SREBP1 and PPAR γ transcription factors (Laplanche and Sabatini 2009a). In addition to promoting anabolism, mTORC1 also inhibits catabolism by blocking autophagy through the phosphorylation of the ULK1–Atg13–FIP200 complex. Importantly, sustained mTORC1 activation blocks growth factor signaling by activating many negative feedback loops that inhibit PI3-kinase (PI3K) signaling.

Compared with mTORC1, our knowledge of mTORC2 biology is still very limited. mTORC2 is activated by growth factors through a mechanism that is not well understood.

It is insensitive to acute treatment with rapamycin and regulates cell survival, metabolism, and cytoskeletal organization by phosphorylating many AGC kinases, including Akt, SGK1, and PKC α (Laplanche and Sabatini 2009b; Zoncu et al. 2011).

Figures adapted from Laplanche and Sabatini (2009b).

REFERENCES

- Brown EJ, Albers MW, Shin TB, Ichikawa K, Keith CT, Lane WS, Schreiber SL. 1994. A mammalian protein targeted by G1-arresting rapamycin-receptor complex. *Nature* **369**: 756–758.
- Heitman J, Movva NR, Hall MN. 1991. Targets for cell cycle arrest by the immunosuppressant rapamycin in yeast. *Science* **253**: 905–909.
- Kunz J, Henriquez R, Schneider U, Deuter-Reinhard M, Movva NR, Hall MN. 1993. Target of rapamycin in yeast, TOR2, is an essential phosphatidylinositol kinase homolog required for G₁ progression. *Cell* **73**: 585–596.
- Laplanche M, Sabatini DM. 2009a. An emerging role of mTOR in lipid biosynthesis. *Curr Biol* **19**: R1046–R1052.
- Laplanche M, Sabatini DM. 2009b. mTOR signaling at a glance. *J Cell Sci* **122**: 3589–3594.
- Ma XM, Blenis J. 2009. Molecular mechanisms of mTOR-mediated translational control. *Nat Rev Mol Cell Biol* **10**: 307–318.
- Sabatini DM, Erdjument-Bromage H, Lui M, Tempst P, Snyder SH. 1994. RAFT1: A mammalian protein that binds to FKBP12 in a rapamycin-dependent fashion and is homologous to yeast TORs. *Cell* **78**: 35–43.
- Sabers CJ, Martin MM, Brunn GJ, Williams JM, Dumont FJ, Wiederrecht G, Abraham RT. 1995. Isolation of a protein target of the FKBP12–rapamycin complex in mammalian cells. *J Biol Chem* **270**: 815–822.
- Vezina C, Kudelski A, Sehgal SN. 1975. Rapamycin (AY-22,989), a new antifungal antibiotic. I. Taxonomy of the producing streptomycete and isolation of the active principle. *J Antibiot (Tokyo)* **28**: 721–726.
- Zoncu R, Efeyan A, Sabatini DM. 2011. mTOR: From growth signal integration to cancer, diabetes and ageing. *Nat Rev Mol Cell Biol* **12**: 21–35.

Cold Spring Harbor Perspectives in Biology

mTOR Signaling

Mathieu Laplante and David M. Sabatini

Cold Spring Harb Perspect Biol 2012; doi: 10.1101/cshperspect.a011593 originally published online November 30, 2011

Subject Collection [Signal Transduction](#)

Cell Signaling and Stress Responses

Gökhan S. Hotamisligil and Roger J. Davis

Protein Regulation in Signal Transduction

Michael J. Lee and Michael B. Yaffe

Synaptic Signaling in Learning and Memory

Mary B. Kennedy

Vertebrate Reproduction

Sally Kornbluth and Rafael Fissore

Signaling in Lymphocyte Activation

Doreen Cantrell

Signaling in Muscle Contraction

Ivana Y. Kuo and Barbara E. Ehrlich

Toll-Like Receptor Signaling

Kian-Huat Lim and Louis M. Staudt

Signaling Pathways that Regulate Cell Division

Nicholas Rhind and Paul Russell

Second Messengers

Alexandra C. Newton, Martin D. Bootman and John D. Scott

Signals and Receptors

Carl-Henrik Heldin, Benson Lu, Ron Evans, et al.

Cell Death Signaling

Douglas R. Green and Fabien Llambi

Signaling Networks that Regulate Cell Migration

Peter Devreotes and Alan Rick Horwitz

Signaling Networks: Information Flow, Computation, and Decision Making

Evren U. Azeloglu and Ravi Iyengar

Signal Transduction: From the Atomic Age to the Post-Genomic Era

Jeremy Thorner, Tony Hunter, Lewis C. Cantley, et al.

Signaling by the TGF β Superfamily

Jeffrey L. Wrana

Subversion of Cell Signaling by Pathogens

Neal M. Alto and Kim Orth

For additional articles in this collection, see <http://cshperspectives.cshlp.org/cgi/collection/>

**All Modifications and
Oligo Types Synthesized**
Long Oligos • Fluorescent • Chimeric • DNA • RNA • Antisense

Oligo Modifications?
Your wish is our command.