Helicase Loading at Chromosomal Origins of Replication

Stephen P. Bell¹ and Jon M. Kaguni²

¹Howard Hughes Medical Institute, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139

²Department of Biochemistry and Molecular Biology, Michigan State University, East Lansing, Michigan 48824

Correspondence: kaguni@msu.edu

Loading of the replicative DNA helicase at origins of replication is of central importance in DNA replication. As the first of the replication fork proteins assemble at chromosomal origins of replication, the loaded helicase is required for the recruitment of the rest of the replication machinery. In this work, we review the current knowledge of helicase loading at *Escherichia coli* and eukaryotic origins of replication. In each case, this process requires both an origin recognition protein as well as one or more additional proteins. Comparison of these events shows intriguing similarities that suggest a similar underlying mechanism, as well as critical differences that likely reflect the distinct processes that regulate helicase loading in bacterial and eukaryotic cells.

Replicative DNA helicases are ring-shaped molecules with a central cavity through which DNA passes as they unwind DNA. Their loading at replication origins is a critical and highly regulated event in chromosomal replication. The DNA helicase is the first of the replication fork proteins recruited to and loaded onto origins of replication, and the loaded helicase is required for the recruitment of the rest of the replication machinery (Remus and Difley 2009; Kaguni 2011). Indeed, the replicative DNA helicase links the replication machinery to the parental DNA (O'Donnell 2006). In *Escherichia coli* cells, the DnaB replicative helicase binds to primase (DnaG) and the sliding clamp loader, which in turn binds the DNA polymerases. Although the polymerases are also linked

to the template DNA by sliding clamps, when these interactions are broken, the polymerases' association with the sliding clamp loader and the helicase keeps them at the site of replication. The interactions that tether the DNA polymerases to the eukaryotic replication fork are less clear but very likely involve direct and indirect interactions with the Mcm2–7 replicative helicase (Calzada et al. 2005).

Helicase loading is carefully regulated to control the location and frequency of replication initiation. In eukaryotic cells, helicase loading is tightly restricted to the G₁ phase of the cell cycle. This constraint is a key part of the mechanisms that ensure that no origin can initiate more than once per cell cycle (Siddiqui et al. 2013; Zielke et al. 2013). In addition, the sites

S.P. Bell and J.M. Kaguni

of eukaryotic replicative helicase loading define the potential sites of replication initiation in the cell (but not all loaded helicases are used during a given S phase [Rhind and Gilbert 2012]). Although the central regulated event in bacterial chromosome duplication is the recruitment of the ATP-bound initiator protein DnaA (Skarstad and Katayama 2013), the loading of the replicative helicase represents a key committed step during initiation.

Here we will discuss the mechanism of helicase loading in bacteria and eukaryotic cells. Much of the discussion will focus on studies in the bacterium *E. coli*, the yeast *Saccharomyces cerevisiae*, and the frog *Xenopus laevis*, in which the events of helicase loading are best understood. Comparison of these mechanisms shows important similarities and differences between the domains of life. In both bacteria and eukaryotic cells, multiple AAA⁺ proteins use ATP binding and hydrolysis to direct helicase loading and both helicases are initially loaded in an inactive form. On the other hand, the eukaryotic helicase is loaded around double-stranded DNA (dsDNA) and as a double hexamer, whereas the bacterial helicase is loaded around single-stranded DNA (ssDNA) as a single hexamer. These distinctions are very likely due to the very different regulatory mechanisms of DNA replication in bacteria and eukaryotic cells.

Before describing the process of helicase loading, it is relevant to know that the essential function of replicative helicases is to unwind the parental duplex DNA using the energy provided by the hydrolysis of a nucleoside triphosphate (Table 1) (reviewed in Patel et al. 2011). Replicative DNA polymerases then copy each parental DNA strand to duplicate the genome. That replicative DNA helicases are hexameric (and

sometimes heptameric) structures that have a positively charged, central channel provides a framework for how these molecular machines work. Several models that describe the mechanism of unwinding have been considered. One is the steric or strand-exclusion model, in which the helicase excludes one strand of DNA while the other passes through the central cavity as the enzyme moves. Because these enzymes can bind and translocate on duplex DNA without unwinding (Kaplan et al. 2003), two additional models have been proposed. In the ploughshare model, duplex DNA enters the central cavity and exits as unwound DNA by virtue of a domain/protein that acts as a ploughshare or pin, which disrupts the hydrogen bonds of DNA as it is pumped through the enzyme (reviewed in Takahashi et al. 2005). A second model, the DNA-pumping model, was proposed on the basis of the double-hexameric forms of SV40, Mcm2–7, and other replicative helicases (Mastrangelo et al. 1989; Remus et al. 2009). This model proposes that the two helicases pump duplex DNA toward one another, resulting in torsional strain that forces the two strands apart, at which point they exit the central channel as two ssDNA loops. Current evidence supports the steric exclusion model (Jezewska et al. 1998b; Kaplan 2000; Galletto et al. 2004; Fu et al. 2011). Of interest, the *E. coli* enzyme moves in the 5′-to-3′ direction relative to the engaged ssDNA that passes through the central cavity, whereas archaeal and eukaryotic enzymes move in the 3′-to-5′ direction.

HELICASE LOADING IN *E. coli*

The *E. coli* replication origin (*oriC*) has two essential functions. One is to serve as a site where

Table 1. Replicative DNA helicases of free-living organisms are hexameric

Domain	Model organism(s)	Helicase ^a	Direction of movement
Bacteria	<i>Escherichia coli</i>	DnaB	5′ → 3′
Archaea	<i>Sulfolobus solfataricus</i>	MCM	3′ → 5′
Eukarya	<i>Saccharomyces cerevisiae</i>	Mcm2–7	3′ → 5′
	<i>Drosophila melanogaster</i>		
	<i>Xenopus laevis</i>		

^aDnaB and *S. solfataricus* MCM are homohexamers, whereas eukaryotic Mcm2–7 is composed of six nonidentical subunits.

the replication fork machinery assembles. DNA sequence motifs in this chromosomal locus and their roles are described in detail elsewhere (Leonard and Méchali 2013). The second is to act as a site where the process of DNA replication is controlled. As described in Skarstad and Katayama (2013), several separate mechanisms modulate the frequency of initiation of DNA replication. At this stage, a critical event is the loading of the replicative helicase named DnaB (Bell and Botchan 2013), which must be in complex with its partner, DnaC. A summary of the functions of DnaA, DnaB, and DnaC serves as the foundation for a description of the helicase loading process.

DnaA IS THE REPLICATION INITIATOR IN BACTERIA

E. coli DnaA performs a central role in the initiation of DNA replication at *oriC*, recognizing specific DNA sequences named the DnaA box, I-site, τ -site, and C-site within *oriC* (Leonard and Méchali 2013). Studies reveal that this protein, which is highly conserved among bacteria, can be divided into four domains designated 1 to 4 from amino to carboxyl terminus (<http://www.molgen.mpg.de/~messer/>) (reviewed in Mott and Berger 2007; Ozaki and Katayama 2009; Leonard and Grimwade 2011). The interactions of these domains with specific proteins, DNA sequences, and nucleotides are summarized below.

The amino-terminal domain 1 (amino acids 1–90 of *E. coli* DnaA) interacts with a variety of proteins, including DnaB (Sutton et al. 1998), and DnaA itself in the assembly of the DnaA oligomer at *oriC* (Simmons et al. 2003; Felczak and Kaguni 2004; Abe et al. 2007). These interactions are critical for helicase loading and are described in more detail below. Other proteins that interact with domain 1 are HU (Chodavarapu et al. 2008a), DiaA (Keyamura et al. 2007, 2009), Dps (Chodavarapu et al. 2008b), ribosomal protein L2 (Chodavarapu et al. 2011), and RNA polymerase (Flåtten et al. 2009). The interaction with RNA polymerase correlates with the role of DnaA as a transcriptional activator in stabilizing the binding

of RNA polymerase at the *gidA* and λP_R promoters (Szalewska-Pałasz et al. 1998; Flåtten et al. 2009), and possibly at the *glpD* and *fliC* promoters (Mizushima et al. 1994; Messer and Weigel 1997). With HU or DiaA, their respective binding stabilizes DnaA oligomerized at *oriC* to stimulate initiation in vitro (Keyamura et al. 2007, 2009; Chodavarapu et al. 2008a). By comparison, L2 inhibits initiation by impairing the formation of a DnaA oligomer at *oriC* (Chodavarapu et al. 2011). Like L2, Dps bound to domain 1 interferes with strand opening of *oriC* in vitro, which correlates with less frequent initiation in vivo when the Dps level is elevated (Chodavarapu et al. 2008b). These observations suggest that domain 1 acts as a sensor, responding to proteins that modulate the initiation process by affecting the assembly of DnaA at *oriC*.

Among DnaAs of bacteria (<http://www.molgen.mpg.de/~messer/>), domain 2 (residues 90–130 of *E. coli* DnaA) is not conserved in amino acid sequence or length. For *E. coli* DnaA, the removal of residues 96–120 in domain 2 leads to reduced activity in DNA replication, but deletion of other consecutive sequences from this region has no apparent effect on activity (Nozaki and Ogawa 2008; Molt et al. 2009). These observations suggest that domain 2 acts as a flexible link between domains 1 and 3.

The third domain of DnaA (residues 130–347 of *E. coli* DnaA) is responsible for ATP binding and hydrolysis (Duderstadt and Berger 2008). Like other members of the AAA⁺ family of ATPases, this region of DnaA carries the Walker A (P-loop) and B boxes that are involved in ATP binding. The sensor I, II (box VII), and box VIII motifs are believed to coordinate ATP hydrolysis with a change in conformation (reviewed in Erzberger and Berger 2006). The adenine nucleotide bound to a shortened variant of *Aquifex aeolicus* DnaA lacking domains 1 and 2 has a striking effect on its X-ray crystal structure (Erzberger et al. 2002, 2006). With ADP, six monomers of this DnaA assemble as a closed ring. With ATP instead, an open ring with a right-handed pitch forms by virtue of a distended conformation of domain 3.

Acidic phospholipids in a fluid bilayer bind to a segment spanning the junction between

S.P. Bell and J.M. Kaguni

domains 3 and 4 (reviewed in Boeneman and Crooke 2005). Intriguingly, this interaction promotes nucleotide release from DnaA (Sekimizu and Kornberg 1988). As the concentration of ATP is higher than that of ADP in vivo, this effect may favor the ATP-bound state of DnaA and is the basis for speculation that this interaction may regulate the activity of DnaA. Although fluorescence microscopy indicates that DnaA associates with the inner membrane in vivo (Boeneman et al. 2009) as well as with the *oriC* region (Nozaki et al. 2009), it is not known if interactions with the inner membrane promote nucleotide exchange in living cells.

Domain 4 (residues 347–467) binds to the DnaA box. Nuclear magnetic resonance and X-ray crystallographic analysis of this domain reveal a helix-turn-helix amino acid motif and a basic loop (Erzberger et al. 2002; Fujikawa et al. 2003). Combined with the biochemical characterization of mutant DnaAs bearing specific amino acid substitutions that impair the recognition of the DnaA box, we have an atomic understanding of how DnaA binds to the DnaA box (Sutton and Kaguni 1997; Blaesing et al. 2000).

DnaB IS THE REPLICATIVE HELICASE, AND INTERACTS WITH DnaC TO FORM THE DnaB–DnaC COMPLEX

In *E. coli*, DnaB is the replicative helicase whose native structure is a toroid of six identical subunits oriented in the same direction (Reha-Krantz and Hurwitz 1978; Arai et al. 1981; Donate et al. 2000; Bailey et al. 2007). Whereas Mg^{2+} ion stabilizes this native structure, removal of the metal ion by dialysis or chelation causes the DnaB hexamer to dissociate into trimers and monomers (Bujalowski et al. 1994). X-ray crystallography of the homologous proteins from *Geobacillus kaustophilus* and *Geobacillus stearothermophilus* provides a detailed view of this helicase and the relative arrangement of its smaller amino-terminal and larger carboxy-terminal domains (Bailey et al. 2007; Lo et al. 2009).

Studies of *E. coli* DnaB and others in the DnaB helicase family indicate that the enzyme unwinds DNA by translocating in the 5′-to-3′

direction on the ssDNA to which it is bound (Venkatesan et al. 1982; Matson et al. 1983; LeBowitz and McMacken 1986; Lee et al. 1989; Richardson and Nossal 1989). Fluorescent energy transfer experiments suggest that 20 ± 3 nucleotides of ssDNA pass through the central cavity of DnaB during translocation (Jezewska et al. 1996; Kaplan 2000). DnaB is uniquely oriented on a forked DNA molecule with a 5′ ssDNA tail (Jezewska et al. 1998a,b). On the basis of this orientation, the smaller amino-terminal domain is upstream of the carboxy-terminal domain, which is nearer the apex of the replication fork (Fig. 1, step 3).

To participate in helicase loading at *oriC*, DnaB must be in a complex with its partner, DnaC. Early studies suggested that ATP bound to DnaC is necessary for DnaC to form an isolable complex with DnaB and to protect DnaC from inactivation by *N*-ethylmaleimide (Wickner and Hurwitz 1975; Kobori and Kornberg 1982). More recent studies indicate that ATP is not needed for DnaC to interact with DnaB (Davey et al. 2002; Galletto et al. 2003; Mott et al. 2008). Both cryoelectron microscopy of the DnaB–DnaC complex in comparison with DnaB and fluorescence energy transfer experiments indicate that DnaC is bound to the larger (carboxy-terminal) end of the DnaB toroid (Bárcena et al. 2001; Galletto et al. 2003). Relative to the ensuing direction of helicase movement, DnaC in the DnaB–DnaC complex is positioned proximal to the replication fork junction (Fig. 1).

Because amino acid substitutions within residues 10–44 near the amino terminus of DnaC impair the interaction between DnaC and DnaB, this region of DnaC apparently binds directly to DnaB to form the DnaB–DnaC complex (Ludlam et al. 2001). Initial studies showed that six DnaC monomers bind per DnaB hexamer (Wickner and Hurwitz 1975; Kobori and Kornberg 1982; Lanka and Schuster 1983). Subsequent sedimentation velocity analysis to determine the affinity of DnaC for DnaB in combination with estimates of the in vivo levels of DnaC and DnaB suggest that the cellular DnaB–DnaC complex is heterogeneous in composition, with a fraction containing less than the

Helicase Loading at Chromosomal Origins

Figure 1. Helicase loading at the *E. coli* replication origin (*oriC*) is a stepwise process. Near the top of the figure, DNA sequence elements in *E. coli oriC* are shown (see Leonard and Méchali 2013 for details). Binding sites for Fis and IHF are shown, which are believed to alter the architecture of *oriC*, and their roles are described in detail in Leonard and Méchali (2013). The four domains of DnaA shown in green are represented by the overlaid numbers. In step 1, DnaA complexed to ATP binds to the DnaA boxes, I-, τ -, and C-sites to form a DnaA oligomer. Following the unwinding of the region of *oriC* containing the 13-mers, DnaA then loads a DnaB–DnaC complex on each of the separated strands (step 2). In step 3, primase (DnaG) interacts with the amino-terminal region of DnaB. Primer formation by primase (steps 4 and 5) leads to the dissociation of DnaC from the carboxy-terminal domain of DnaB, which is necessary to activate DnaB as a DNA helicase (see Bell and Botchan 2013). Although not shown, the primers displayed in black are then extended by the cellular replicase, DNA polymerase III holoenzyme, for the continuous synthesis of each leading strand at each replication fork (see Johansson and Dixon 2013). A dimer of this DNA polymerase is believed to be at each replication fork. DnaB is proposed to interact with one unit of a dimer of DNA polymerase III holoenzyme, coordinating the unwinding of the parental DNA with DNA synthesis. As this DNA helicase translocates in the 5'-to-3' direction to support replication fork movement, primase occasionally interacts with DnaB to synthesize additional primers. These primers are used by the unit of the DNA polymerase dimer that synthesizes Okazaki fragments.

S.P. Bell and J.M. Kaguni

maximum of six DnaC monomers per DnaB hexamer (Galletto et al. 2003).

DnaC CONTROLS THE ACTIVITY OF DnaB

DnaC, like DnaA, is a member of the AAA⁺ family of ATPases (Makarova and Koonin 2013). Despite the presence in DnaC of motifs shared by AAA⁺ family members and the high affinity of other AAA⁺ proteins for ATP, DnaC binds weakly to this nucleotide; the K_d is about 8 μ M (Davey et al. 2002; Galletto et al. 2003; Biswas et al. 2004). By itself, DnaC is also a feeble ATPase. The influence of ATP on DnaC function has been an enigma, but new evidence suggests that its ATPase activity is activated at a specific step of the initiation process (see below). Stimulated by ATP, DnaC also interacts weakly with ssDNA (Learn et al. 1997; Davey et al. 2002; Mott et al. 2008), supporting the idea that this activity may be involved in helicase loading. Because DnaC complexed to DnaB inhibits its ATPase and helicase activities (Wahle et al. 1989a,b; Allen and Kornberg 1991; Davey et al. 2002; Mott et al. 2008), DnaC must dissociate from DnaB for helicase activation to occur.

DnaA UNWINDS A REGION NEAR THE LEFT BORDER OF *oriC*

With a negatively supercoiled plasmid carrying *oriC*, DnaA bound to ATP unwinds an AT-rich region containing the M and R 13-mers as judged by potassium permanganate sensitivity (Fig. 1) (Sekimizu et al. 1987; Bramhill and Kornberg 1988; Gille and Messer 1991). Considering that isolated plasmids have a negative superhelix density ranging from -0.05 to -0.08 , as much as 330 to 530 nucleotides can be single-stranded if these DNAs are supercoiled solely as a result of unwinding of the duplex DNA instead of the writhing of the DNA upon itself (Cozzarelli 1980; Bliska and Cozzarelli 1987). These observations suggest that DnaA confines some of the single-stranded character of the plasmid to this AT-rich region of *oriC*. To provide a biochemical mechanism of unwinding, recent studies on the assembly of a DnaA oligomer as a right-handed filament suggest that positively

charged and hydrophobic amino acids that line the interior of the DnaA filament interact with the negatively charged ssDNA (Erzberger et al. 2006; Ozaki et al. 2008; Duderstadt et al. 2010, 2011; Ozaki and Katayama 2011). Consistent with the ATP requirement for DnaA-dependent origin unwinding, the formation of this DnaA filament requires the ATP-bound form of the enzyme.

HELICASE LOADING AT *oriC* BY DnaA

After unwinding of *oriC*, DnaA loads the DnaB–DnaC complex onto this unwound DNA, which apparently becomes enlarged to include the left 13-mer and additional sequences beyond it (Fig. 1, step 2) (Davey et al. 2002). Footprinting studies suggest that the DnaB–DnaC complex protects a region on the top strand near the left border of *oriC*. On the bottom strand the area protected is to the left of DnaA box R1 that overlaps the right 13-mer. Quantitative analysis of this protein complex assembled at *oriC* supports the conclusion from footprinting studies that two DnaB–DnaC complexes are bound (Fang et al. 1999; Carr and Kaguni 2001).

Helicase loading appears to involve two regions of DnaA. One region is within domain 3, based on studies of a monoclonal antibody that interferes with the interaction between DnaA and DnaB measured in solid-phase binding assays (Marszalek and Kaguni 1994; Sutton et al. 1998). This antibody recognizes an epitope within residues 111–148 of DnaA. Deletion analysis localized the interacting region to amino acids 135–148 (Seitz et al. 2000). A region within domain 1 is also needed because an alanine substitution for phenylalanine 46 abrogates this interaction (Keyamura et al. 2009). On the basis that DnaC from *A. aeolicus* can interact with DnaA, and like DnaA forms a helical filament as determined by X-ray crystallography, a specific proposal is that DnaA oligomerized at *oriC* loads one DnaB–DnaC complex on the top strand via the interaction between DnaA molecules at the left end of the DnaA filament and DnaC in the DnaB–DnaC complex (Mott et al. 2008). The second DnaB–DnaC complex loads on the bottom strand

through the interaction between the left end of the DnaA oligomer and DnaB complexed to DnaC. During loading, it is interesting to consider that one of the interfaces between DnaB protomers must open in this ring-shaped protein in order for the helicase to encircle the ssDNA. The role of DnaA or DnaC in this ring opening remains unclear.

After DnaA loads one DnaB–DnaC complex on each of the separated strands of *oriC*, DnaC must dissociate for DnaB to be active as a helicase. Of interest, recent studies suggest that three DnaC monomers are bound to each DnaB hexamer (Makowska-Grzyska and Kaguni 2010), whereas earlier experiments suggested that the DnaB–DnaC complex contains six DnaC monomers per DnaB (Wickner and Hurwitz 1975; Kobori and Kornberg 1982; Lanka and Schuster 1983; Wahle et al. 1989a). Perhaps the different forms of the DnaB–DnaC complex function at the initiation stage of DNA replication and in restarting stalled or collapsed replication forks.

ATP or ATP γ S bound to DnaC supports the DnaA-dependent loading of the DnaB–DnaC complex at *oriC* (Davey et al. 2002; Makowska-Grzyska and Kaguni 2010). That DnaC remains bound to DnaB in the presence of either ATP or the analog suggests that the act of helicase loading does not stimulate the hydrolysis of ATP bound to DnaC or its release from DnaB (Makowska-Grzyska and Kaguni 2010). In the presence of ATP and other ribonucleotides, the inclusion of primase (DnaG), which interacts with the amino-terminal domain of DnaB as it forms primers for DNA replication, induces the release of DnaC from the carboxy-terminal (large) domain of DnaB (Fig. 1, steps 3–5). Apparently the interaction of primase with DnaB while it synthesizes a primer alters the conformation of DnaB that causes DnaC to dissociate. Other results with mutant forms of DnaC that are speculated to be defective in ATP hydrolysis suggest that this interaction of primase with DnaB stimulates the hydrolysis of ATP bound by DnaC. This step of helicase activation is comparable in function to the activation of the Mcm2–7 complex at a eukaryotic replication origin (Labib 2010).

HELICASE LOADING IN EUKARYOTIC CELLS

Origin Licensing, Pre-RC Formation, and Helicase Loading

Early work on eukaryotic DNA replication revealed that the events of replication initiation were separated into separate cell-cycle stages. Mammalian cell fusion studies suggested the existence of positive factors in S-phase cells that cause G₁ nuclei to initiate DNA replication (Johnson and Rao 1970). Analyses of cycling *Xenopus* egg extracts suggested the existence of a “licensing factor” that gained access to the DNA only after nuclear envelope breakdown (during mitosis) and was consumed during S phase (Blow and Laskey 1988). Finally, in vivo footprinting studies at *S. cerevisiae* origins showed a distinct and more extensive protection pattern during G₁ (called the prereplicative complex or pre-RC) than in S, G₂, and M phase (Diffley et al. 1994).

We now know that replication origin licensing and pre-RC formation are both related to the helicase loading event that only occurs in G₁ cells (Bell and Dutta 2002; Sclafani and Holzen 2007). The licensing factor that is excluded from the nucleus is probably not identical among different cell types (Arias and Walter 2007), but origin licensing is clearly equivalent to helicase loading. The pre-RC footprint that is associated with *S. cerevisiae* origins during G₁ is due to binding of helicase loading factors to the origin DNA (Perkins and Diffley 1998; Speck et al. 2005). Much of this footprint is attributed to origin recognition complex (ORC) and Cdc6, but it additionally reflects helicase loading. For this reason the term “pre-RC formation” has become synonymous with helicase loading. We will refer to the G₁ events at origins as helicase loading and the events that occur in S phase as helicase activation and replisome assembly because these terms most accurately reflect the events that are occurring at the origin.

Mcm2–7 HELICASE

The eukaryotic replicative DNA helicase is the Mcm2–7 complex. Each Mcm2–7 complex contains one copy of the six essential and related

S.P. Bell and J.M. Kaguni

Mcm2, Mcm3, Mcm4, Mcm5, Mcm6, and Mcm7 proteins (reviewed in Bochman and Schwacha 2009). Although high-resolution structural data are not available, electron microscopy (EM) and subunit interaction studies indicate that the Mcm2–7 proteins interact in a defined order (Davey et al. 2003; Yu et al. 2004; Bochman et al. 2008) to form either a ring (Remus et al. 2009) or a gapped ring (Costa et al. 2011) with a positively charged central channel (Fig. 2). The carboxy-terminal half of each Mcm2–7 subunit contains a conserved AAA⁺ domain that includes insertions specific to the Mcm2–7 proteins (Iyer et al. 2004) that are predicted to form β -hairpins that may interact with the ssDNA during unwinding (Bochman and Schwacha 2009). As with other AAA⁺ multimers, the interface between subunits in the Mcm2–7 complex forms an ATPase active site. The amino-terminal half of each subunit contains an OB-fold-related motif that is found in many ssDNA-binding proteins, and several of the subunits also contain a Zn-finger motif. Although the AAA⁺ domain is highly conserved among Mcm2–7 subunits, the characteristic amino- and carboxy-terminal extensions allow homologs for each of the six Mcm2–7 proteins to be readily

identified in all eukaryotes studied. As described above, as a 3' $\rightarrow$ 5' helicase the Mcm2–7 helicase moves on the leading-strand DNA template at a replication fork.

EUKARYOTIC HELICASE LOADING PROTEINS

Eukaryotic helicase loading is directed by three proteins: ORC, Cdc6, and Cdt1. ORC is a heterohexamer with five of the six subunits (Orc1, -2, -3, -4, and -5) showing homology with AAA⁺ proteins, although only two of these subunits (Orc1 and Orc5) are known to bind ATP (Klemm et al. 1997). Four ORC subunits (Orc1, -2, -4, and -5) also contain potential winged-helix domains at their carboxyl termini. The smallest ORC subunit, Orc6, is related to the RNA Pol II auxiliary factor TFIIB in metazoans (Liu et al. 2011), although fungal Orc6 lacks this similarity. Like Mcm2–7 and ORC, Cdc6 is an AAA⁺ protein in the same initiator clade as the AAA⁺ ORC subunits, DnaA and DnaC (Iyer et al. 2004). In addition, the carboxy-terminal portion of Cdc6 folds into a winged-helix domain (Liu et al. 2000). Both Orc6 and Cdt1 are poorly conserved across eukaryotes, but all Cdt1

Figure 2. Structure of the Mcm2–7 complex. (A) The Mcm2–7 proteins assemble into a hexameric toroid. Each Mcm2–7 complex includes one copy of the Mcm2–7 proteins that are arranged in the indicated order around a central channel. (B) Directionality of Mcm2–7 movement. The Mcm2–7 proteins move in a 3' $\rightarrow$ 5' direction along ssDNA. By analogy with the archaea homologs, the carboxy-terminal AAA⁺ motif is proximal to the replication fork.

proteins carry a pair of winged-helix domains in their carboxy-terminal half (Lee et al. 2004; Khayrutdinov et al. 2009; Jee et al. 2010) and the most carboxy-terminal winged helix binds to Mcm2–7 (Yanagi et al. 2002; Teer and Dutta 2008; You and Masai 2008; Jee et al. 2010). Interestingly, structural studies suggest that the winged-helix domains associated with ORC, Cdc6, and Cdt1 are related (Khayrutdinov et al. 2009; Jee et al. 2010). Mcm8 and Mcm9 are Mcm2–7-related proteins found in metazoan cells, but are not found to associate with any of the Mcm2–7 proteins either in or out of the Mcm2–7 complex (Maiorano et al. 2005; Lutzmann and Méchali 2008). Although these proteins have been implicated in helicase loading (Volkening and Hoffmann 2005; Lutzmann and Méchali 2008), this view remains controversial (Blanton et al. 2005; Hartford et al. 2011) and a direct mechanistic involvement in helicase loading has not been shown.

INITIAL RECRUITMENT OF THE HELICASE TO THE ORIGIN DNA

Before helicase loading, ORC, Cdc6, Cdt1, and Mcm2–7 are recruited to the origin DNA. This complex is short-lived and is only detected in vitro when Cdc6 ATP hydrolysis is inhibited (Randell et al. 2006). In vivo, chromatin immunoprecipitation assays only detect ORC and Mcm2–7 at specific origins (Aparicio et al. 1997; Tanaka et al. 1997), presumably because of the rapid release of Cdc6 and Cdt1 after helicase loading.

ORIGIN RECOGNITION

The first step of helicase loading is the binding of ORC to origin DNA. In the yeast *S. cerevisiae*, ORC recognizes a conserved sequence within the origin of replication and this event is ATP-dependent (Bell and Stillman 1992). This sequence is not sufficient to direct ORC binding in vivo. An important additional determinant is an asymmetric nucleosome-free region surrounding the consensus sequence (Eaton et al. 2010). *S. pombe* ORC shows limited sequence specificity that is mediated by the AT-rich DNA-

binding specificity of an AT-hook domain within the Orc4 subunit (Chuang and Kelly 1999; Kong and DePamphilis 2001; Lee et al. 2001).

The determinants for ORC binding in metazoan cells are less clear. ORC from metazoan cells does not exhibit sequence-specific DNA binding in vitro (Vashee et al. 2003; Remus et al. 2004), but ORC is localized to specific sites along metazoan chromosomes, indicating the existence of other determinants for ORC localization (Karnani et al. 2010; MacAlpine et al. 2010). Genome-wide studies of metazoan ORC binding support a number of determinants including nucleosome-depleted regions, promoter proximity, and dynamic nucleosomes (reviewed in Ding and MacAlpine 2011). In addition, DNA-binding studies show that ORC prefers to bind negatively supercoiled DNA (Remus et al. 2004; Houchens et al. 2008), which may be related to the wrapping of DNA around ORC (Clarey et al. 2006, 2008). Perhaps the most intriguing potential mechanism of ORC localization is the recognition of specifically modified nucleosomes. Human ORC binds a protein called ORCA or LRWD1 that interacts with methylated nucleosomes that are associated with heterochromatin, suggesting an alternative method to localize ORC to compacted chromatin (Bartke et al. 2010; Shen et al. 2010; Vermeulen et al. 2010). In addition, the amino terminus of the Orc1 protein contains a conserved bromo-adjacent homology (BAH) domain. In metazoan cells this domain interacts with histone H4 tails methylated at lysine 20, suggesting another mechanism for ORC to be localized along the chromatin (Kuo et al. 2012).

INTERACTION OF Cdc6, Cdt1, AND Mcm2–7

As cells enter G₁ phase, ORC recruits Cdc6, Cdt1, and Mcm2–7 to the origin DNA. Biochemical studies support a model in which ORC first interacts with Cdc6 and this complex then recruits Cdt1 and Mcm2–7. Loss of Cdt1 does not prevent Cdc6 chromatin association in vivo or origin recruitment in vitro (Maiorano et al. 2000; Nishitani et al. 2000; Tsuyama et al. 2005; Randell et al. 2006; Remus et al. 2009). In contrast,

S.P. Bell and J.M. Kaguni

elimination of Cdc6 interferes with Cdt1 origin binding in vitro (Randell et al. 2006; Remus et al. 2009). Studies in *Xenopus* extracts have shown that Cdt1 associates with chromatin in the absence of Cdc6 (Gillespie et al. 2001; Tsuyama et al. 2005); however, only Cdt1 associated in the presence of Cdc6 is able to contribute to Mcm2–7 loading (Tsuyama et al. 2005). Consistent with a robust interaction between Cdc6 and ORC, a complex between the proteins has been structurally characterized (Speck et al. 2005; Sun et al. 2012) and origin-bound ORC stimulates Cdc6 ATP hydrolysis (Randell et al. 2006).

Mcm2–7 and Cdt1 are recruited to the origin as a complex. The carboxy-terminal winged-helix domain of Cdt1 binds Mcm2–7 (Yanagi et al. 2002; Ferenbach et al. 2005; Teer and Dutta 2008; You and Masai 2008; Zhang et al. 2010; Takara and Bell 2011). Cdt1 mutants lacking this region prevent or dominantly inhibit Mcm2–7 loading (Ferenbach et al. 2005; Zhang et al. 2010; Takara and Bell 2011). In *S. cerevisiae* cells, Cdt1 and Mcm2–7 are imported to the nucleus as a complex (Tanaka and Diffley 2002), and formation of this complex is necessary for the recruitment of either protein to the origin (Takara and Bell 2011). The primary Cdt1-binding site on Mcm2–7 is the carboxy-terminal region of Mcm6, and structural studies suggest that this region also folds into a winged-helix domain (Yanagi et al. 2002; Wei et al. 2010).

Characterization of the helicase loading intermediate composed of ORC, Cdc6, Cdt1, and Mcm2–7 suggests how a single asymmetric ORC protein can direct the assembly of a bi-directional replication fork. Binding of Cdt1/Mcm2–7 to the origin requires the smallest ORC subunit, Orc6, which includes two binding sites for Cdt1 (Chen et al. 2007). Analysis of the protein complex formed at the origin before helicase loading (when Cdc6 ATP hydrolysis is inhibited) indicates that multiple Cdt1 proteins and their associated Mcm2–7 complexes are initially recruited to each ORC–Cdc6 complex (Takara and Bell 2011). It is most likely that this complex contains two Cdt1 molecules, one bound to each of the two Cdt1-binding sites on Orc6. Importantly, interfering with the for-

mation of this multi-Cdt1 intermediate prevents subsequent helicase loading, and it is likely that this intermediate facilitates the simultaneous loading of an Mcm2–7 double hexamer (see below).

How Cdc6 contributes to Cdt1 and Mcm2–7 recruitment is less well understood. It is possible that there are direct interactions between Cdc6 and Cdt1 and/or Mcm2–7. Alternatively, Cdc6 could alter the conformation of ORC to allow Cdt1 and Mcm2–7 to bind. The latter hypothesis would explain why ORC does not bind either Cdt1 or Mcm2–7 in the absence of Cdc6 (Takara and Bell 2011) but isolated Orc6 can bind Cdt1 (Chen et al. 2007). Consistent with this model, recent EM studies of ORC bound to DNA in the presence and absence of Cdc6 suggested that Orc6 becomes exposed upon Cdc6 binding (Sun et al. 2012).

LOADING OF RECRUITED HELICASES

After recruitment of the helicase loading factors and Mcm2–7 to the origin, loading of the recruited Mcm2–7 requires ATP hydrolysis and involves significant changes of both the proteins associated with the origin and the structure of the Mcm2–7 ring. Loaded Mcm2–7 complexes form double-hexameric pairs that encircle dsDNA. Thus, helicase loading necessarily requires the establishment of strong interactions between the amino termini of two Mcm2–7 complexes and opening and closing of the Mcm2–7 ring to allow DNA access to the Mcm2–7 central channel. Finally, to ensure the proper regulation of replication initiation, as initially loaded, the Mcm2–7 complexes are inactive for unwinding the parental duplex.

STRUCTURE OF LOADED Mcm2–7

The reconstitution of Mcm2–7 loading using four purified proteins has led to important advances in the understanding of the architecture of loaded Mcm2–7. EM of loaded Mcm2–7 shows head-to-head double hexamers, with the amino termini of the Mcm2–7 subunits mediating hexamer interactions (Evrin et al. 2009; Remus et al. 2009). This structure is

similar to the double hexamers observed for the homohexameric archaea Mcm complexes (Brewster and Chen 2010). Unlike the archaea homologs, Mcm2–7 double hexamers are only detected after loading. Mcm2–7 double hexamers survive treatment with DNase and gel filtration, indicating that once formed, these complexes are very stable (Evrin et al. 2009; Gambus et al. 2011). Based on structural studies of the archaea Mcm complex, these interactions likely involve the Zn-finger domains in the amino termini of Mcm2–7 subunits (Fletcher et al. 2003). The inter-Mcm2–7 contacts within the double hexamer are unknown but must be largely heterotypic owing to their head-to-head interaction. A lack of loaded single hexamers and the presence of multiple Cdt1/Mcm2–7 complexes before Mcm2–7 loading supports a model in which both hexamers are loaded in a concerted fashion (Remus et al. 2009; Takara and Bell 2011). Importantly, both EM and topological linkage studies indicate that loaded Mcm2–7 complexes encircle and slide nondirectionally on dsDNA (Evrin et al. 2009; Remus et al. 2009). Thus, loaded Mcm2–7 is topologically linked to the DNA but is neither active as a helicase nor tightly engaged with the DNA.

ROLE OF ATP DURING HELICASE LOADING

ATP binding and hydrolysis play critical roles during helicase loading. As described above, ATP binding but not hydrolysis is required for the initial recruitment of the helicase to the origin (Gillespie et al. 2001; Randell et al. 2006). ATP hydrolysis by Cdc6 and ORC plays distinct and ordered roles in helicase loading. Cdc6 is activated to bind and hydrolyze ATP when it associates with origin-bound ORC, and this hydrolysis event is required to observe Mcm2–7 loading (Perkins and Diffley 1998; Randell et al. 2006). ORC ATP hydrolysis is also essential but functions after Cdc6 hydrolysis and appears to regulate repeated Mcm2–7 loading (Bowers et al. 2004; Randell et al. 2006). One likely role for ORC ATP hydrolysis is to cause the release of the other helicase loading factors from the origin (Tsakraklides and Bell 2010). Evidence from studies of *S. cerevisiae* suggests that a sec-

ond consequence of ATP hydrolysis is to disable ORC DNA binding until it is able to bind another molecule of ATP. An important unanswered question is how many Mcm2–7 hexamers are loaded per round of ATP hydrolysis by Cdc6 and ORC. Although current evidence suggests two hexamers are loaded coordinately, it remains possible that Cdc6 or ORC could require multiple rounds of ATP hydrolysis to accomplish the loading of an Mcm2–7 double hexamer. Overall, the coordinated and ordered set of ATP hydrolysis reactions ensures that Cdc6 only acts on Cdt1/Mcm2–7 when bound to ORC at the origin and a new round of loading cannot occur until the previous round is complete.

Mcm2–7 GATE FUNCTION AND IMPLICATIONS FOR LOADING

The Mcm2–7 ring structure must be opened during loading to provide access to the central DNA-binding channel. Binding of purified Mcm2–7 to ssDNA circles suggests that the interface between Mcm2 and Mcm5 acts as a “gate” to the central channel (Bochman and Schwacha 2008). EM studies of purified *Drosophila* Mcm2–7 support this conclusion (Costa et al. 2011). Although ATP binding is believed to close the Mcm2/5 gate, it is likely that some combination of Cdt1 binding and Cdc6 ATP hydrolysis modulates Mcm2–7 ring opening during helicase loading. Because Cdt1 is released from Mcm2–7 after loading (Randell et al. 2006), it is tempting to speculate that Cdt1 binding opens the Mcm2/5 gate and Cdc6 ATP hydrolysis separates Cdt1 from Mcm2–7, leaving behind a closed form of the Mcm2–7 ring. The role of ATP binding and hydrolysis by Mcm2–7 during helicase loading remains uncertain despite studies to address this issue. Mutations in the Mcm6 and Mcm7 ATP-binding sites show normal chromatin association but are defective for replication initiation (Ying and Gautier 2005). It is unclear, however, whether the observed chromatin association reflects loading or just Mcm2–7 recruitment. In addition, the distinct functions of six Mcm2–7 ATPase active sites (reviewed in Bochman and

S.P. Bell and J.M. Kaguni

Schwacha 2009) suggest that ATP bound to other sites could be required for helicase loading.

REGULATION AND DYNAMICS OF HELICASE LOADING

Helicase loading is tightly restricted to G₁ phase to ensure that no origin of replication can initiate DNA replication more than once per cell cycle (reviewed in Arias and Walter 2007; Siddiqui et al. 2013; Zielke et al. 2013). This regulation is mediated by the inhibition of helicase loading outside of G₁ phase. The mechanisms of inhibition are diverse and vary among organisms. In budding yeast, inhibition is primarily mediated by cyclin-dependent kinase (CDK) phosphorylation of Cdc6, which leads to its degradation (Drury et al. 2000), Mcm2–7 nuclear export (Labib et al. 1999; Nguyen et al. 2000), and the inhibition of Cdt1/Mcm2–7 binding to ORC (Chen and Bell 2011). In metazoan organisms, Cdt1 is a primary target for inhibition. Cdt1 is inhibited by geminin binding outside of G₁ (McGarry and Kirschner 1998; Wohlschlegel et al. 2000) and degraded in a proliferating cell nuclear antigen (PCNA)/DNA-dependent (Arias and Walter 2005) or CDK-dependent (Li et al. 2003; Liu et al. 2004) fashion in S phase. In addition, in many metazoans, CDK activity inhibits ORC DNA binding during G₂/M phase (reviewed in DePamphilis 2005). Although it is tempting to think that the multiple mechanisms that inhibit helicase loading are redundant, analysis of mutants that are defective for a subset of mechanisms shows partial rereplication (Green et al. 2006). Importantly, even limited rereplication is lethal to most dividing cells and can induce gene amplification associated with cancer (Green et al. 2010).

The role of the helicase loading proteins after loading is unclear. In vitro studies show that after loading, Mcm2–7 no longer requires ORC, Cdc6, or Cdt1 to associate with origin DNA (Donovan et al. 1997; Randell et al. 2006). Consistent with this observation, in vitro studies show that the helicase loading proteins are released from the origin upon Mcm2–7 loading (Randell et al. 2006; Tsakraklides and Bell 2010). In contrast to these findings, in vivo in-

activation of *S. cerevisiae* ORC or Cdc6 in late G₁ (after Mcm2–7 loading) results in loss of Mcm2–7 origin association (Aparicio et al. 1997; Semple et al. 2006; Chen et al. 2007). This suggests either that these proteins are required to maintain Mcm2–7 association with the origin DNA or that there is an activity that removes loaded Mcm2–7 from the DNA in G₁ cells and the helicase loading proteins are required to restore Mcm2–7 to the origin. In either case, this reliance on helicase loading proteins changes upon entry into S phase, as Mcm2–7 is tightly associated with origins at this stage (unless initiation has occurred) even though Cdc6 is eliminated from S phase cells by degradation. It is noteworthy that studies in mammalian G₁ cells suggest that there is little exchange between chromatin-associated and free Mcm2–7 (Kuijpers et al. 2011), arguing for a model in which helicase loading proteins stabilize loaded Mcm2–7 rather than a model in which Mcm2–7 release is balanced with new loading.

A MODEL FOR HELICASE LOADING

Based on current knowledge, we propose the following model for the events of eukaryotic helicase loading (Fig. 3). ORC recruitment to origin DNA is directed by a combination of DNA affinity (sequence-specific in budding yeast), local chromatin structure, and interaction with other proteins (e.g., ORCA). Upon entry into G₁ phase, ORC recruits Cdc6. In *S. cerevisiae* cells, we propose that this interaction reveals Cdt1-binding sites on Orc6 that are bound by two Cdt1/Mcm2–7 complexes. The distinct structure of Orc6 in metazoans raises the possibility that the initial recruitment is mediated by different interactions in these organisms. The connection of these events to Mcm2–7 ring opening is unclear; however, it seems most likely that the Mcm2/5 gate is opened at some point during these events. Alternatively, the ring could be opened before recruitment (e.g., after Cdt1 binds to Mcm2–7), although this would risk Mcm2–7 interacting with nonorigin DNA. Assuming two Mcm2–7 complexes are loaded coordinately, we suggest that the Mcm2/5 gates of the two helicases are aligned to form a single continuous

Figure 3. Model for eukaryotic replicative DNA helicase loading. After localization to the origin DNA, ATP-bound ORC recruits Cdc6 bound to ATP. The resulting ORC–Cdc6 complex then recruits two Cdt1/Mcm2–7 complexes via interactions between Cdt1 and Orc6. Although this illustration suggests that the Mcm2–7 complexes have initiated interactions at their amino termini at this stage, it is also possible that these interactions only occur during or after helicase loading. The interactions between ORC, Cdc6, Cdt1, and Mcm2–7 are proposed to result in the opening of the Mcm2–7 ring at the Mcm2/5 gate. Cdc6 ATP hydrolysis results in the loading of an Mcm2–7 double hexamer around double-stranded origin DNA and the release of Cdt1. Whether the DNA enters the Mcm2–7 central channel before (upon initial ring opening) or after (as illustrated) Cdc6 ATP hydrolysis is unknown. ORC ATP hydrolysis is proposed to lead to the release of Cdc6–ADP and loaded Mcm2–7 from ORC. ORC ADP/ATP exchange leads to resetting of the loading machinery, allowing a new round of helicase loading to initiate.

opening/gate. By analogy to the function of the sliding clamp loader (Kelch et al. 2011), we speculate that the Mcm2–7 complexes can encircle the DNA at this stage but remain open. The next stage of loading is triggered by Cdc6 ATP hydrolysis. This event leads to the release of Cdc6 and Cdt1 from the helicase, which is coupled with entrapment of the DNA within the Mcm2–7 central channel and closure of the Mcm2/5 gate. Finally, we propose that ATP hydrolysis by ORC leads to the release of the loaded helicases from ORC to prepare for a new round of loading after ADP → ATP exchange.

Once loaded, the Mcm2–7 complex awaits activation. Biochemical studies show that activation requires the association of two helicase activating proteins (Cdc45 and GINS) with Mcm2–7 (Ilves et al. 2010). Association of these proteins is triggered by the action of two kinases: S-phase CDK (S-CDK) and the Dbf4-dependent Cdc7 kinase (DDK). DDK phosphorylates Mcm2–7 (Sheu and Stillman 2006, 2010; Randell et al. 2010). As a result of Mcm2–7 phosphorylation, Cdc45 associates with Mcm2–7 (Heller et al. 2011; Tanaka et al. 2011). Interestingly, this event occurs at a subset of origins before entry into S phase (Aparicio et al. 1999; Kanemaki and Labib 2006; Heller et al. 2011; Tanaka et al. 2011). S-CDK targets two additional proteins, Sld2 and Sld3, causing them to associate with Dpb11 and eventually resulting in the recruitment of the GINS proteins as well as DNA Pol ϵ (Tanaka et al. 2007; Zegerman and Diffley 2007; Muramatsu et al. 2010). The subsequent activation of DNA unwinding is required for the recruitment of the lagging-strand DNA polymerases (DNA Pol α -primase and DNA Pol δ) (Mimura et al. 2000; Heller et al. 2011).

COMPARISON OF HELICASES AND HELICASE LOADING IN *E. coli* AND EUKARYOTIC CELLS

There are many similarities between helicase loading at *E. coli* and eukaryotic chromosomal origins, suggesting that the fundamental mechanism of these events is related. Both *E. coli* DnaB and eukaryotic Mcm2–7 are composed

S.P. Bell and J.M. Kaguni

of six subunits that assemble into ring-shaped molecules. Second, both events rely on origin recognition proteins (DnaA and ORC) that contact the origin DNA over an extended region. Third, loading of each DNA helicase requires at least one additional helicase loading factor that binds to the helicase. In *E. coli*, DnaC is required for loading and also restrains DnaB helicase activity. In eukaryotes, both Cdc6 and Cdt1 share properties with DnaC. Like DnaC, Cdc6 is an AAA⁺ protein and is a weak ATPase on its own, whereas Cdt1 shares DnaC's association with its cognate helicase. Finally, both helicases must be activated after loading, either by release of an inhibitor (DnaC in *E. coli*) or by recruiting helicase activators (Cdc45 and GINS in eukaryotic cells).

There are also a number of important differences. First, although similar in overall structure, DnaB and the Mcm2–7 complexes have opposite polarities: DnaB moves on the lagging-strand template, whereas Mcm2–7 moves on the leading-strand template. DnaB is a homohexamer that has a RecA-like ATPase domain (Bailey et al. 2007), whereas Mcm2–7 is a heterohexamer that has AAA⁺-like ATPase domains (Iyer et al. 2004). Second, although DnaA origin binding leads to local DNA melting, ORC DNA binding does not. Third, two separate single hexamers of DnaB are loaded around ssDNA after loading, whereas a head-to-head double hexamer of Mcm2–7 is loaded around dsDNA.

These differences are most likely related to the rate of helicase activation after loading. Loading of DnaB around ssDNA at *oriC* is consistent with the rapid activation of DnaB after loading. In contrast, the need to restrain helicase activation until S phase in eukaryotic cells suggests the need for a loaded helicase that is less easily activated. Recent findings indicate that Mcm2–7 surrounds ssDNA and acts as a single hexamer at the replication fork (Yardimci et al. 2010; Fu et al. 2011). This indicates that Mcm2–7 activation requires two major changes in the loaded double hexamer. First, each Mcm2–7 complex must dissociate from its double-hexamer partner. Second, the Mcm2–7 ring originally bound to duplex DNA must open to expel

the lagging-strand ssDNA template before DNA unwinding. Although we have an initial understanding of how DnaC expulsion leads to activation of DnaB (Makowska-Grzyska and Kaguni 2010), how the helicase activating proteins Cdc45 and GINS and the proteins that direct them to the origin (Sld2, Sld3, and Dpb11) catalyze these events remains unknown.

Although 25 years of research has provided an increasingly detailed understanding of the events of helicase loading, fundamental questions remain. How is the opening and closing of the DNA helicase ring controlled? How is the origin DNA properly positioned to ensure that the helicase encircles the appropriate form of the DNA at the correct position? How are these events regulated by the AAA⁺ ATPases that are conserved among the different realms of life? These and other questions propel studies of this central event in replication initiation.

ACKNOWLEDGMENTS

We thank members of our labs for helpful discussions. This work is supported by the Michigan Agricultural Station and by grant RO1 GM090063 from the National Institutes of Health to J.M.K. and by the Howard Hughes Medical Institute and grant RO1 GM052339 from the National Institutes of Health to S.P.B.

REFERENCES

*Reference is also in this collection.

- Abe Y, Jo T, Matsuda Y, Matsunaga C, Katayama T, Ueda T. 2007. Structure and function of DnaA N-terminal domains: Specific sites and mechanisms in inter-DnaA interaction and in DnaB helicase loading on *oriC*. *J Biol Chem* **282**: 17816–17827.
- Allen GC, Kornberg A. 1991. Fine balance in the regulation of DnaB helicase by DnaC protein in replication in *Escherichia coli*. *J Biol Chem* **266**: 22096–22101.
- Aparicio OM, Weinstein DM, Bell SP. 1997. Components and dynamics of DNA replication complexes in *S. cerevisiae*: Redistribution of MCM proteins and Cdc45p during S phase. *Cell* **91**: 59–69.
- Aparicio OM, Stout AM, Bell SP. 1999. Differential assembly of Cdc45p and DNA polymerases at early and late origins of DNA replication. *Proc Natl Acad Sci* **96**: 9130–9135.
- Arai K, Yasuda S, Kornberg A. 1981. Mechanism of *dnaB* protein action. I. Crystallization and properties of *dnaB*

- protein, an essential replication protein in *Escherichia coli*. *J Biol Chem* **256**: 5247–5252.
- Arias EE, Walter JC. 2005. PCNA functions as a molecular platform to trigger Cdt1 destruction and prevent re-replication. *Nat Cell Biol* **8**: 84–90.
- Arias EE, Walter JC. 2007. Strength in numbers: Preventing rereplication via multiple mechanisms in eukaryotic cells. *Genes Dev* **21**: 497–518.
- Bailey S, Eliason WK, Steitz TA. 2007. Structure of hexameric DnaB helicase and its complex with a domain of DnaG primase. *Science* **318**: 459–463.
- Bárcena M, Ruiz T, Donate LE, Brown SE, Dixon NE, Radermacher M, Carazo JM. 2001. The DnaB-DnaC complex: A structure based on dimers assembled around an occluded channel. *EMBO J* **20**: 1462–1468.
- Bartke T, Vermeulen M, Xhemalce B, Robson SC, Mann M, Kouzarides T. 2010. Nucleosome-interacting proteins regulated by DNA and histone methylation. *Cell* **143**: 470–484.
- * Bell SD, Botchan MR. 2013. The minichromosome maintenance replicative helicase. *Cold Spring Harb Perspect Biol* doi: 10.1101/cshperspect.a012807.
- Bell SP, Dutta A. 2002. DNA replication in eukaryotic cells. *Annu Rev Biochem* **71**: 333–374.
- Bell SP, Stillman B. 1992. ATP-dependent recognition of eukaryotic origins of DNA replication by a multiprotein complex. *Nature* **357**: 128–134.
- Biswas SB, Flowers S, Biswas-Fiss EE. 2004. Quantitative analysis of nucleotide modulation of DNA binding by DnaC protein of *Escherichia coli*. *Biochem J* **379**: 553–562.
- Blaesing F, Weigel C, Welzeck M, Messer W. 2000. Analysis of the DNA-binding domain of *Escherichia coli* DnaA protein. *Mol Microbiol* **36**: 557–569.
- Blanton HL, Radford SJ, McMahan S, Kearney HM, Ibrahim JG, Sekelsky J. 2005. REC, *Drosophila* MCM8, drives formation of meiotic crossovers. *PLoS Genet* **1**: e40.
- Bliska JB, Cozzarelli NR. 1987. Use of site-specific recombination as a probe of DNA structure and metabolism in vivo. *J Mol Biol* **194**: 205–218.
- Blow JJ, Laskey RA. 1988. A role for the nuclear envelope in controlling DNA replication within the cell cycle. *Nature* **332**: 546–548.
- Bochman ML, Schwacha A. 2008. The Mcm2-7 complex has in vitro helicase activity. *Mol Cell* **31**: 287–293.
- Bochman ML, Schwacha A. 2009. The Mcm complex: Unwinding the mechanism of a replicative helicase. *Microbiol Mol Biol Rev* **73**: 652–683.
- Bochman ML, Bell SP, Schwacha A. 2008. Subunit organization of Mcm2-7 and the unequal role of active sites in ATP hydrolysis and viability. *Mol Cell Biol* **28**: 5865–5873.
- Boeneman K, Crooke E. 2005. Chromosomal replication and the cell membrane. *Curr Opin Microbiol* **8**: 143–148.
- Boeneman K, Fossum S, Yang Y, Fingland N, Skarstad K, Crooke E. 2009. *Escherichia coli* DnaA forms helical structures along the longitudinal cell axis distinct from MreB filaments. *Mol Microbiol* **72**: 645–657.
- Bowers JL, Randell JCW, Chen S, Bell SP. 2004. ATP hydrolysis by ORC catalyzes reiterative Mcm2-7 assembly at a defined origin of replication. *Mol Cell* **16**: 967–978.
- Bramhill D, Kornberg A. 1988. Duplex opening by dnaA protein at novel sequences in initiation of replication at the origin of the *E. coli* chromosome. *Cell* **52**: 743–755.
- Brewster AS, Chen XS. 2010. Insights into the MCM functional mechanism: Lessons learned from the archaeal MCM complex. *Crit Rev Biochem Mol Biol* **45**: 243–256.
- Bujalowski W, Klonowska MM, Jezewska MJ. 1994. Oligomeric structure of *Escherichia coli* primary replicative helicase DnaB protein. *J Biol Chem* **269**: 31350–31358.
- Calzada A, Hodgson B, Kanemaki M, Bueno A, Labib K. 2005. Molecular anatomy and regulation of a stable replisome at a paused eukaryotic DNA replication fork. *Genes Dev* **19**: 1905–1919.
- Carr KM, Kaguni JM. 2001. Stoichiometry of DnaA and DnaB protein in initiation at the *Escherichia coli* chromosomal origin. *J Biol Chem* **276**: 44919–44925.
- Chen S, Bell SP. 2011. CDK prevents Mcm2-7 helicase loading by inhibiting Cdt1 interaction with Orc6. *Genes Dev* **25**: 363–372.
- Chen S, de Vries MA, Bell SP. 2007. Orc6 is required for dynamic recruitment of Cdt1 during repeated Mcm2-7 loading. *Genes Dev* **21**: 2897–2907.
- Chodavarapu S, Felczak MM, Yaniv JR, Kaguni JM. 2008a. *Escherichia coli* DnaA interacts with HU in initiation at the *E. coli* replication origin. *Mol Microbiol* **67**: 781–792.
- Chodavarapu S, Gomez R, Vicente M, Kaguni JM. 2008b. *Escherichia coli* Dps interacts with DnaA protein to impede initiation: A model of adaptive mutation. *Mol Microbiol* **67**: 1331–1346.
- Chodavarapu S, Felczak MM, Kaguni JM. 2011. Two forms of ribosomal protein L2 of *Escherichia coli* that inhibit DnaA in DNA replication. *Nucleic Acids Res* **39**: 4180–4191.
- Chuang RY, Kelly TJ. 1999. The fission yeast homologue of Orc4p binds to replication origin DNA via multiple AT-hooks. *Proc Natl Acad Sci* **96**: 2656–2661.
- Clarey MG, Erzberger JP, Grob P, Leschziner AE, Berger JM, Nogales E, Botchan M. 2006. Nucleotide-dependent conformational changes in the DnaA-like core of the origin recognition complex. *Nat Struct Mol Biol* **13**: 684–690.
- Clarey MG, Botchan M, Nogales E. 2008. Single particle EM studies of the *Drosophila melanogaster* origin recognition complex and evidence for DNA wrapping. *J Struct Biol* **164**: 241–249.
- Costa A, Ilves I, Tamberg N, Petojevic T, Nogales E, Botchan MR, Berger JM. 2011. The structural basis for MCM2-7 helicase activation by GINS and Cdc45. *Nat Struct Mol Biol* **18**: 471–477.
- Cozzarelli NR. 1980. DNA gyrase and the supercoiling of DNA. *Science* **207**: 953–960.
- Davey MJ, Fang L, McInerney P, Georgescu RE, O'Donnell M. 2002. The DnaC helicase loader is a dual ATP/ADP switch protein. *EMBO J* **21**: 3148–3159.
- Davey MJ, Indiani C, O'Donnell M. 2003. Reconstitution of the Mcm2-7p heterohexamer, subunit arrangement, and ATP site architecture. *J Biol Chem* **278**: 4491–4499.
- DePamphilis ML. 2005. Cell cycle dependent regulation of the origin recognition complex. *Cell Cycle* **4**: 70–79.

S.P. Bell and J.M. Kaguni

- Diffley JE, Cocker JH, Dowell SJ, Rowley A. 1994. Two steps in the assembly of complexes at yeast replication origins in vivo. *Cell* **78**: 303–316.
- Ding Q, MacAlpine DM. 2011. Defining the replication program through the chromatin landscape. *Crit Rev Biochem Mol Biol* **46**: 165–179.
- Donate LE, Llorca O, Bárcena M, Brown SE, Dixon NE, Carazo JM. 2000. pH-controlled quaternary states of hexameric DnaB helicase. *J Mol Biol* **303**: 383–393.
- Donovan S, Harwood J, Drury LS, Diffley JE. 1997. Cdc6p-dependent loading of Mcm proteins onto pre-replicative chromatin in budding yeast. *Proc Natl Acad Sci* **94**: 5611–5616.
- Drury LS, Perkins G, Diffley JE. 2000. The cyclin-dependent kinase Cdc28p regulates distinct modes of Cdc6p proteolysis during the budding yeast cell cycle. *Curr Biol* **10**: 231–240.
- Duderstadt KE, Berger JM. 2008. AAA⁺ ATPases in the initiation of DNA replication. *Crit Rev Biochem Mol Biol* **43**: 163–187.
- Duderstadt KE, Mott ML, Crisona NJ, Chuang K, Yang H, Berger JM. 2010. Origin remodeling and opening in bacteria rely on distinct assembly states of the DnaA initiator. *J Biol Chem* **285**: 28229–28239.
- Duderstadt KE, Chuang K, Berger JM. 2011. DNA stretching by bacterial initiators promotes replication origin opening. *Nature* **478**: 209–213.
- Eaton ML, Galani K, Kang S, Bell SP, MacAlpine DM. 2010. Conserved nucleosome positioning defines replication origins. *Genes Dev* **24**: 748–753.
- Erzberger JP, Berger JM. 2006. Evolutionary relationships and structural mechanisms of AAA⁺ proteins. *Annu Rev Biophys Biomol Struct* **35**: 93–114.
- Erzberger JP, Pirruccello MM, Berger JM. 2002. The structure of bacterial DnaA: Implications for general mechanisms underlying DNA replication initiation. *EMBO J* **21**: 4763–4773.
- Erzberger JP, Mott ML, Berger JM. 2006. Structural basis for ATP-dependent DnaA assembly and replication-origin remodeling. *Nat Struct Mol Biol* **13**: 676–683.
- Evrin C, Clarke P, Zech J, Lurz R, Sun J, Uhle S, Li H, Stillman B, Speck C. 2009. A double-hexameric MCM2-7 complex is loaded onto origin DNA during licensing of eukaryotic DNA replication. *Proc Natl Acad Sci* **106**: 20240–20245.
- Fang L, Davey MJ, O'Donnell M. 1999. Replisome assembly at *oriC*, the replication origin of *E. coli*, reveals an explanation for initiation sites outside an origin. *Mol Cell* **4**: 541–553.
- Felczak MM, Kaguni JM. 2004. The box VII motif of *Escherichia coli* DnaA protein is required for DnaA oligomerization at the *E. coli* replication origin. *J Biol Chem* **279**: 51156–51162.
- Ferenbach A, Li A, Brito-Martins M, Blow JJ. 2005. Functional domains of the *Xenopus* replication licensing factor Cdt1. *Nucleic Acids Res* **33**: 316–324.
- Flåtten I, Morigen, Skarstad K. 2009. DnaA protein interacts with RNA polymerase and partially protects it from the effect of rifampicin. *Mol Microbiol* **71**: 1018–1030.
- Fletcher RJ, Bishop BE, Leon RP, Sclafani RA, Ogata CM, Chen XS. 2003. The structure and function of MCM from archaeal *M. thermoautotrophicum*. *Nat Struct Biol* **10**: 160–167.
- Fu YV, Yardimci H, Long DT, Guainazzi A, Bermudez VP, Hurwitz J, van Oijen A, Schärer OD, Walter JC. 2011. Selective bypass of a lagging strand roadblock by the eukaryotic replicative DNA helicase. *Cell* **146**: 931–941.
- Fujikawa N, Kurumizaka H, Nureki O, Terada T, Shirouzu M, Katayama T, Yokoyama S. 2003. Structural basis of replication origin recognition by the DnaA protein. *Nucleic Acids Res* **31**: 2077–2086.
- Galletto R, Jezewska MJ, Bujalowski W. 2003. Interactions of the *Escherichia coli* DnaB helicase hexamer with the replication factor the DnaC protein. Effect of nucleotide cofactors and the ssDNA on protein–protein interactions and the topology of the complex. *J Mol Biol* **329**: 441–465.
- Galletto R, Jezewska MJ, Bujalowski W. 2004. Unzipping mechanism of the double-stranded DNA unwinding by a hexameric helicase: The effect of the 3' arm and the stability of the dsDNA on the unwinding activity of the *Escherichia coli* DnaB helicase. *J Mol Biol* **343**: 101–114.
- Gambus A, Khoudoli GA, Jones RC, Blow JJ. 2011. MCM2-7 form double hexamers at licensed origins in *Xenopus* egg extract. *J Biol Chem* **286**: 11855–11864.
- Gille H, Messer W. 1991. Localized DNA melting and structural perturbations in the origin of replication, *oriC*, of *Escherichia coli* in vitro and in vivo. *EMBO J* **10**: 1579–1584.
- Gillespie PJ, Li A, Blow JJ. 2001. Reconstitution of licensed replication origins on *Xenopus* sperm nuclei using purified proteins. *BMC Biochem* **2**: 15.
- Green BM, Morreale RJ, Ozaydin B, Derisi JL, Li JJ. 2006. Genome-wide mapping of DNA synthesis in *Saccharomyces cerevisiae* reveals that mechanisms preventing reinitiation of DNA replication are not redundant. *Mol Biol Cell* **17**: 2401–2414.
- Green BM, Finn KJ, Li JJ. 2010. Loss of DNA replication control is a potent inducer of gene amplification. *Science* **329**: 943–946.
- Hartford SA, Luo Y, Southard TL, Min IM, Lis JT, Schimenti JC. 2011. Minichromosome maintenance helicase paralog MCM9 is dispensable for DNA replication but functions in germ-line stem cells and tumor suppression. *Proc Natl Acad Sci* **108**: 17702–17707.
- Heller RC, Kang S, Lam WM, Chen S, Chan CS, Bell SP. 2011. Eukaryotic origin-dependent DNA replication in vitro reveals sequential action of DDK and S-CDK kinases. *Cell* **146**: 80–91.
- Houchens CR, Lu W, Chuang R-Y, Frattini MG, Fuller A, Simanek P, Kelly TJ. 2008. Multiple mechanisms contribute to *Schizosaccharomyces pombe* origin recognition complex-DNA interactions. *J Biol Chem* **283**: 30216–30224.
- Ivles I, Petojevic T, Pesavento JJ, Botchan MR. 2010. Activation of the MCM2-7 helicase by association with Cdc45 and GINS proteins. *Mol Cell* **37**: 247–258.
- Iyer LM, Leipe DD, Koonin EV, Aravind L. 2004. Evolutionary history and higher order classification of AAA⁺ ATPases. *J Struct Biol* **146**: 11–31.
- Jee J, Mizuno T, Kamada K, Tochio H, Chiba Y, Yanagi K-I, Yasuda G, Hiroaki H, Hanaoka F, Shirakawa M. 2010.

- Structure and mutagenesis studies of the C-terminal region of licensing factor Cdt1 enable the identification of key residues for binding to replicative helicase Mcm proteins. *J Biol Chem* **285**: 15931–15940.
- Jezewska MJ, Kim US, Bujalowski W. 1996. Binding of *Escherichia coli* primary replicative helicase DnaB protein to single-stranded DNA. Long-range allosteric conformational changes within the protein hexamer. *Biochemistry* **35**: 2129–2145.
- Jezewska MJ, Rajendran S, Bujalowska D, Bujalowski W. 1998a. Does single-stranded DNA pass through the inner channel of the protein hexamer in the complex with the *Escherichia coli* DnaB helicase? Fluorescence energy transfer studies. *J Biol Chem* **273**: 10515–10529.
- Jezewska MJ, Rajendran S, Bujalowski W. 1998b. Complex of *Escherichia coli* primary replicative helicase DnaB protein with a replication fork: Recognition and structure. *Biochemistry* **37**: 3116–3136.
- * Johansson E, Dixon N. 2013. Replicative DNA polymerases. *Cold Spring Harb Perspect Biol* doi: 10.1101/cshperspect.a012799.
- Johnson RT, Rao PN. 1970. Mammalian cell fusion: Induction of premature chromosome condensation in interphase nuclei. *Nature* **226**: 717–722.
- Kaguni JM. 2011. Replication initiation at the *Escherichia coli* chromosomal origin. *Curr Opin Chem Biol* **15**: 606–613.
- Kanemaki M, Labib K. 2006. Distinct roles for Sld3 and GINS during establishment and progression of eukaryotic DNA replication forks. *EMBO J* **25**: 1753–1763.
- Kaplan DL. 2000. The 3'-tail of a forked-duplex sterically determines whether one or two DNA strands pass through the central channel of a replication-fork helicase. *J Mol Biol* **301**: 285–299.
- Kaplan DL, Davey MJ, O'Donnell M. 2003. Mcm4, 6, 7 uses a "pump in ring" mechanism to unwind DNA by steric exclusion and actively translocate along a duplex. *J Biol Chem* **278**: 49171–49182.
- Karnani N, Taylor CM, Malhotra A, Dutta A. 2010. Genomic study of replication initiation in human chromosomes reveals the influence of transcription regulation and chromatin structure on origin selection. *Mol Biol Cell* **21**: 393–404.
- Kelch BA, Makino DL, O'Donnell M, Kuriyan J. 2011. How a DNA polymerase clamp loader opens a sliding clamp. *Science* **334**: 1675–1680.
- Keyamura K, Fujikawa N, Ishida T, Ozaki S, Su'etsugu M, Fujimitsu K, Kagawa W, Yokoyama S, Kurumizaka H, Katayama T. 2007. The interaction of DiaA and DnaA regulates the replication cycle in *E. coli* by directly promoting ATP DnaA-specific initiation complexes. *Genes Dev* **21**: 2083–2099.
- Keyamura K, Abe Y, Higashi M, Ueda T, Katayama T. 2009. DiaA dynamics are coupled with changes in initial origin complexes leading to helicase loading. *J Biol Chem* **284**: 25038–25050.
- Khayrutdinov BI, Bae WJ, Yun YM, Lee JH, Tsuyama T, Kim JJ, Hwang E, Ryu K-S, Cheong H-K, Cheong C, et al. 2009. Structure of the Cdt1 C-terminal domain: Conservation of the winged helix fold in replication licensing factors. *Protein Sci* **18**: 2252–2264.
- Klemm RD, Austin RJ, Bell SP. 1997. Coordinate binding of ATP and origin DNA regulates the ATPase activity of the origin recognition complex. *Cell* **88**: 493–502.
- Kobori JA, Kornberg A. 1982. The *Escherichia coli* dnaC gene product. III. Properties of the dnaB–dnaC protein complex. *J Biol Chem* **257**: 13770–13775.
- Kong D, DePamphilis ML. 2001. Site-specific DNA binding of the *Schizosaccharomyces pombe* origin recognition complex is determined by the Orc4 subunit. *Mol Cell Biol* **21**: 8095–8103.
- Kuipers MA, Stasevich TJ, Sasaki T, Wilson KA, Hazelwood KL, McNally JG, Davidson MW, Gilbert DM. 2011. Highly stable loading of Mcm proteins onto chromatin in living cells requires replication to unload. *J Cell Biol* **192**: 29–41.
- Kuo AJ, Song J, Cheung P, Ishibe-Murakami S, Yamazoe S, Chen JK, Patel DJ, Gozani O. 2012. The BAH domain of ORC1 links H4K20me2 to DNA replication licensing and Meier–Gorlin syndrome. *Nature* **484**: 115–119.
- Labib K. 2010. How do Cdc7 and cyclin-dependent kinases trigger the initiation of chromosome replication in eukaryotic cells? *Genes Dev* **24**: 1208–1219.
- Labib K, Diffley JE, Kearsey SE. 1999. G1-phase and B-type cyclins exclude the DNA-replication factor Mcm4 from the nucleus. *Nat Cell Biol* **1**: 415–422.
- Lanka E, Schuster H. 1983. The dnaC protein of *Escherichia coli*. Purification, physical properties and interaction with dnaB protein. *Nucleic Acids Res* **11**: 987–997.
- Learn BA, Um SJ, Huang L, McMacken R. 1997. Cryptic single-stranded-DNA binding activities of the phage λ P and *Escherichia coli* DnaC replication initiation proteins facilitate the transfer of *E. coli* DnaB helicase onto DNA. *Proc Natl Acad Sci* **94**: 1154–1159.
- LeBowitz JH, McMacken R. 1986. The *Escherichia coli* dnaB replication protein is a DNA helicase. *J Biol Chem* **261**: 4738–4748.
- Lee EH, Kornberg A, Hidaka M, Kobayashi T, Horiuchi T. 1989. *Escherichia coli* replication termination protein impedes the action of helicases. *Proc Natl Acad Sci* **86**: 9104–9108.
- Lee JK, Moon KY, Jiang Y, Hurwitz J. 2001. The *Schizosaccharomyces pombe* origin recognition complex interacts with multiple AT-rich regions of the replication origin DNA by means of the AT-hook domains of the spOrc4 protein. *Proc Natl Acad Sci* **98**: 13589–13594.
- Lee C, Hong B, Choi JM, Kim Y, Watanabe S, Ishimi Y, Enomoto T, Tada S, Kim Y, Cho Y. 2004. Structural basis for inhibition of the replication licensing factor Cdt1 by geminin. *Nature* **430**: 913–917.
- Leonard AC, Grimwade JE. 2011. Regulation of DnaA assembly and activity: Taking directions from the genome. *Annu Rev Microbiol* **65**: 19–35.
- * Leonard AC, Méchali M. 2013. DNA replication origins. *Cold Spring Harb Perspect Biol* doi: 10.1101/cshperspect.a010116.
- Li X, Zhao Q, Liao R, Sun P, Wu X. 2003. The SCF^{Skp2} ubiquitin ligase complex interacts with the human replication licensing factor Cdt1 and regulates Cdt1 degradation. *J Biol Chem* **278**: 30854–30858.
- Liu J, Smith CL, DeRyckere D, DeAngelis K, Martin GS, Berger JM. 2000. Structure and function of Cdc6/

S.P. Bell and J.M. Kaguni

- Cdc18: Implications for origin recognition and checkpoint control. *Mol Cell* **6**: 637–648.
- Liu E, Li X, Yan F, Zhao Q, Wu X. 2004. Cyclin-dependent kinases phosphorylate human Cdt1 and induce its degradation. *J Biol Chem* **279**: 17283–17288.
- Liu S, Balasov M, Wang H, Wu L, Chesnokov IN, Liu Y. 2011. Structural analysis of human Orc6 protein reveals a homology with transcription factor TFIIB. *Proc Natl Acad Sci* **108**: 7373–7378.
- Lo Y-H, Tsai K-L, Sun Y-J, Chen W-T, Huang C-Y, Hsiao C-D. 2009. The crystal structure of a replicative hexameric helicase DnaC and its complex with single-stranded DNA. *Nucleic Acids Res* **37**: 804–814.
- Ludlam AV, McNatt MW, Carr KM, Kaguni JM. 2001. Essential amino acids of *Escherichia coli* DnaC protein in an N-terminal domain interact with DnaB helicase. *J Biol Chem* **276**: 27345–27353.
- Lutzmann M, Méchali M. 2008. MCM9 binds Cdt1 and is required for the assembly of prereplication complexes. *Mol Cell* **31**: 190–200.
- MacAlpine HK, Gordán R, Powell SK, Hartemink AJ, MacAlpine DM. 2010. *Drosophila* ORC localizes to open chromatin and marks sites of cohesin complex loading. *Genome Res* **20**: 201–211.
- Maierano D, Moreau J, Méchali M. 2000. XCDT1 is required for the assembly of pre-replicative complexes in *Xenopus laevis*. *Nature* **404**: 622–625.
- Maierano D, Cuvier O, Danis E, Méchali M. 2005. MCM8 is an MCM2-7-related protein that functions as a DNA helicase during replication elongation and not initiation. *Cell* **120**: 315–328.
- * Makarova KS, Koonin EV. 2013. Archaeology of eukaryotic DNA replication. *Cold Spring Harb Perspect Biol* doi: 10.1101/cshperspect.a012963.
- Makowska-Grzyska M, Kaguni JM. 2010. Primase directs the release of DnaC from DnaB. *Mol Cell* **37**: 90–101.
- Marszalek J, Kaguni JM. 1994. DnaA protein directs the binding of DnaB protein in initiation of DNA replication in *Escherichia coli*. *J Biol Chem* **269**: 4883–4890.
- Mastrangelo IA, Hough PV, Wall JS, Dodson M, Dean FB, Hurwitz J. 1989. ATP-dependent assembly of double hexamers of SV40 T antigen at the viral origin of DNA replication. *Nature* **338**: 658–662.
- Matson SW, Tabor S, Richardson CC. 1983. The gene 4 protein of bacteriophage T7. Characterization of helicase activity. *J Biol Chem* **258**: 14017–14024.
- McGarry TJ, Kirschner MW. 1998. Geminin, an inhibitor of DNA replication, is degraded during mitosis. *Cell* **93**: 1043–1053.
- Messer W, Weigel C. 1997. DnaA initiator—also a transcription factor. *Mol Microbiol* **24**: 1–6.
- Mimura S, Masuda T, Matsui T, Takisawa H. 2000. Central role for Cdc45 in establishing an initiation complex of DNA replication in *Xenopus* egg extracts. *Genes Cells* **5**: 439–452.
- Mizushima T, Tomura A, Shinpuku T, Miki T, Sekimizu K. 1994. Loss of flagellation in *dnaA* mutants of *Escherichia coli*. *J. Bacteriol.* **176**: 5544–5546.
- Molt KL, Sutera VA, Moore KK, Lovett ST. 2009. A role for nonessential domain II of initiator protein, DnaA, in replication control. *Genetics* **183**: 39–49.
- Mott ML, Berger JM. 2007. DNA replication initiation: Mechanisms and regulation in bacteria. *Nat Rev Microbiol* **5**: 343–354.
- Mott ML, Erzberger JP, Coons MM, Berger JM. 2008. Structural synergy and molecular crosstalk between bacterial helicase loaders and replication initiators. *Cell* **135**: 623–634.
- Muramatsu S, Hirai K, Tak Y-S, Kamimura Y, Araki H. 2010. CDK-dependent complex formation between replication proteins Dpb11, Sld2, Pol ϵ , and GINS in budding yeast. *Genes Dev* **24**: 602–612.
- Nguyen VQ, Co C, Irie K, Li JJ. 2000. Clb/Cdc28 kinases promote nuclear export of the replication initiator proteins Mcm2–7. *Curr Biol* **10**: 195–205.
- Nishitani H, Lygerou Z, Nishimoto T, Nurse P. 2000. The Cdt1 protein is required to license DNA for replication in fission yeast. *Nature* **404**: 625–628.
- Nozaki S, Ogawa T. 2008. Determination of the minimum domain II size of *Escherichia coli* DnaA protein essential for cell viability. *Microbiology* **154**: 3379–3384.
- Nozaki S, Niki H, Ogawa T. 2009. Replication initiator DnaA of *Escherichia coli* changes its assembly form on the replication origin during the cell cycle. *J. Bacteriol.* **191**: 4807–4814.
- O'Donnell M. 2006. Replisome architecture and dynamics in *Escherichia coli*. *J Biol Chem* **281**: 10653–10656.
- Ozaki S, Katayama T. 2009. DnaA structure, function, and dynamics in the initiation at the chromosomal origin. *Plasmid* **62**: 71–82.
- Ozaki S, Katayama T. 2011. Highly organized DnaA–oriC complexes recruit the single-stranded DNA for replication initiation. *Nucleic Acids Res* **40**: 1648–1665.
- Ozaki S, Kawakami H, Nakamura K, Fujikawa N, Kagawa W, Park S-Y, Yokoyama S, Kurumizaka H, Katayama T. 2008. A common mechanism for the ATP-DnaA-dependent formation of open complexes at the replication origin. *J Biol Chem* **283**: 8351–8362.
- Patel SS, Pandey M, Nandakumar D. 2011. Dynamic coupling between the motors of DNA replication: Hexameric helicase, DNA polymerase, and primase. *Curr Opin Chem Biol* **15**: 595–605.
- Perkins G, Diffley JE. 1998. Nucleotide-dependent prereplicative complex assembly by Cdc6p, a homolog of eukaryotic and prokaryotic clamp-loaders. *Mol Cell* **2**: 23–32.
- Randell JCW, Bowers JL, Rodriguez HK, Bell SP. 2006. Sequential ATP hydrolysis by Cdc6 and ORC directs loading of the Mcm2–7 helicase. *Mol Cell* **21**: 29–39.
- Randell JCW, Fan A, Chan C, Francis LI, Heller RC, Galani K, Bell SP. 2010. Mec1 is one of multiple kinases that prime the Mcm2–7 helicase for phosphorylation by Cdc7. *Mol Cell* **40**: 353–363.
- Reha-Krantz LJ, Hurwitz J. 1978. The *dnaB* gene product of *Escherichia coli*. II. Single stranded DNA-dependent ribonucleoside triphosphatase activity. *J Biol Chem* **253**: 4051–4057.
- Remus D, Diffley JFX. 2009. Eukaryotic DNA replication control: Lock and load, then fire. *Curr Opin Cell Biol* **21**: 771–777.
- Remus D, Beall EL, Botchan MR. 2004. DNA topology, not DNA sequence, is a critical determinant for *Drosophila* ORC–DNA binding. *EMBO J* **23**: 897–907.

- Remus D, Beuron F, Tolun G, Griffith JD, Morris EP, Diffley JFX. 2009. Concerted loading of Mcm2–7 double hexamers around DNA during DNA replication origin licensing. *Cell* **139**: 719–730.
- * Rhind N, Gilbert DM. 2013. DNA replication timing. *Cold Spring Harb Perspect Biol* doi: 10.1101/cshperspect.a010132.
- Richardson RW, Nossal NG. 1989. Characterization of the bacteriophage T4 gene 41 DNA helicase. *J Biol Chem* **264**: 4725–4731.
- Scalfani RA, Holzen TM. 2007. Cell cycle regulation of DNA replication. *Annu Rev Genet* **41**: 237–280.
- Seitz H, Weigel C, Messer W. 2000. The interaction domains of the DnaA and DnaB replication proteins of *Escherichia coli*. *Mol Microbiol* **37**: 1270–1279.
- Sekimizu K, Kornberg A. 1988. Cardiolipin activation of dnaA protein, the initiation protein of replication in *Escherichia coli*. *J Biol Chem* **263**: 7131–7135.
- Sekimizu K, Bramhill D, Kornberg A. 1987. ATP activates dnaA protein in initiating replication of plasmids bearing the origin of the *E. coli* chromosome. *Cell* **50**: 259–265.
- Seiple JW, Da-Silva LF, Jarvis EJ, Ah-Kee J, Al-Attar H, Kummer L, Heikkila JJ, Pasero P, Duncker BP. 2006. An essential role for Orc6 in DNA replication through maintenance of pre-replicative complexes. *EMBO J* **25**: 5150–5158.
- Shen Z, Sathyan KM, Geng Y, Zheng R, Chakraborty A, Freeman B, Wang F, Prasanth KV, Prasanth SG. 2010. A WD-repeat protein stabilizes ORC binding to chromatin. *Mol Cell* **40**: 99–111.
- Sheu Y-J, Stillman B. 2006. Cdc7-Dbf4 phosphorylates MCM proteins via a docking site-mediated mechanism to promote S phase progression. *Mol Cell* **24**: 101–113.
- Sheu Y-J, Stillman B. 2010. The Dbf4-Cdc7 kinase promotes S phase by alleviating an inhibitory activity in Mcm4. *Nature* **463**: 113–117.
- * Siddiqui K, On KE, Diffley JFX. 2013. Regulating DNA replication in eukarya. *Cold Spring Harb Perspect Biol* doi: 10.1101/cshperspect.a012930.
- Simmons LA, Felczak M, Kaguni JM. 2003. DnaA protein of *Escherichia coli*: Oligomerization at the *E. coli* chromosomal origin is required for initiation and involves specific N-terminal amino acids. *Mol Microbiol* **49**: 849–858.
- * Skarstad K, Katayama T. 2013. Regulating DNA replication in bacteria. *Cold Spring Harb Perspect Biol* doi: 10.1101/cshperspect.a012922.
- Speck C, Chen Z, Li H, Stillman B. 2005. ATPase-dependent cooperative binding of ORC and Cdc6 to origin DNA. *Nat Struct Mol Biol* **12**: 965–971.
- Sun J, Kawakami H, Zech J, Speck C, Stillman B, Li H. 2012. Cdc6-induced conformational changes in ORC bound to origin DNA revealed by cryo-electron microscopy. *Structure* **20**: 534–544.
- Sutton MD, Kaguni JM. 1997. Threonine 435 of *Escherichia coli* DnaA protein confers sequence-specific DNA binding activity. *J Biol Chem* **272**: 23017–23024.
- Sutton MD, Carr KM, Vicente M, Kaguni JM. 1998. *Escherichia coli* DnaA protein. The N-terminal domain and loading of DnaB helicase at the *E. coli* chromosomal origin. *J Biol Chem* **273**: 34255–34262.
- Szalewska-Palasz A, Wegrzyn A, Błaszczak A, Taylor K, Wegrzyn G. 1998. DnaA-stimulated transcriptional activation of *oriλ*: *Escherichia coli* RNA polymerase β subunit as a transcriptional activator contact site. *Proc Natl Acad Sci* **95**: 4241–4246.
- Takahashi TS, Wigley DB, Walter JC. 2005. Pumps, paradoxes and ploughshares: Mechanism of the MCM2–7 DNA helicase. *Trends Biochem Sci* **30**: 437–444.
- Takara TJ, Bell SP. 2011. Multiple Cdt1 molecules act at each origin to load replication-competent Mcm2–7 helicases. *EMBO J* **30**: 4885–4896.
- Tanaka S, Diffley JFX. 2002. Interdependent nuclear accumulation of budding yeast Cdt1 and Mcm2–7 during G1 phase. *Nat Cell Biol* **4**: 198–207.
- Tanaka T, Knapp D, Nasmyth K. 1997. Loading of an Mcm protein onto DNA replication origins is regulated by Cdc6p and CDKs. *Cell* **90**: 649–660.
- Tanaka S, Umemori T, Hirai K, Muramatsu S, Kamimura Y, Araki H. 2007. CDK-dependent phosphorylation of Sld2 and Sld3 initiates DNA replication in budding yeast. *Nature* **445**: 328–332.
- Tanaka S, Nakato R, Katou Y, Shirahige K, Araki H. 2011. Origin association of Sld3, Sld7, and Cdc45 proteins is a key step for determination of origin-firing timing. *Curr Biol* **21**: 2055–2063.
- Teer JK, Dutta A. 2008. Human Cdt1 lacking the evolutionarily conserved region that interacts with MCM2–7 is capable of inducing re-replication. *J Biol Chem* **283**: 6817–6825.
- Tsakraklides V, Bell SP. 2010. Dynamics of pre-replicative complex assembly. *J Biol Chem* **285**: 9437–9443.
- Tsuyama T, Tada S, Watanabe S, Seki M, Enomoto T. 2005. Licensing for DNA replication requires a strict sequential assembly of Cdc6 and Cdt1 onto chromatin in *Xenopus* egg extracts. *Nucleic Acids Res* **33**: 765–775.
- Vashee S, Cvetic C, Lu W, Simancek P, Kelly TJ, Walter JC. 2003. Sequence-independent DNA binding and replication initiation by the human origin recognition complex. *Genes Dev* **17**: 1894–1908.
- Venkatesan M, Silver LL, Nossal NG. 1982. Bacteriophage T4 gene 41 protein, required for the synthesis of RNA primers, is also a DNA helicase. *J Biol Chem* **257**: 12426–12434.
- Vermeulen M, Eberl HC, Matarese F, Marks H, Denissov S, Butter F, Lee KK, Olsen JV, Hyman AA, Stunnenberg HG, et al. 2010. Quantitative interaction proteomics and genome-wide profiling of epigenetic histone marks and their readers. *Cell* **142**: 967–980.
- Volkening M, Hoffmann I. 2005. Involvement of human MCM8 in prereplication complex assembly by recruiting hcdc6 to chromatin. *Mol Cell Biol* **25**: 1560–1568.
- Wahle E, Lasken RS, Kornberg A. 1989a. The dnaB-dnaC replication protein complex of *Escherichia coli*. I. Formation and properties. *J Biol Chem* **264**: 2463–2468.
- Wahle E, Lasken RS, Kornberg A. 1989b. The dnaB-dnaC replication protein complex of *Escherichia coli*. II. Role of the complex in mobilizing dnaB functions. *J Biol Chem* **264**: 2469–2475.
- Wei Z, Liu C, Wu X, Xu N, Zhou B, Liang C, Zhu G. 2010. Characterization and structure determination of the

S.P. Bell and J.M. Kaguni

- Cdt1 binding domain of human minichromosome maintenance (Mcm) 6. *J Biol Chem* **285**: 12469–12473.
- Wickner S, Hurwitz J. 1975. Interaction of *Escherichia coli* *dnaB* and *dnaC(D)* gene products *in vitro*. *Proc Natl Acad Sci* **72**: 921–925.
- Wohlschlegel JA, Dwyer BT, Dhar SK, Cvetcic C, Walter JC, Dutta A. 2000. Inhibition of eukaryotic DNA replication by geminin binding to Cdt1. *Science* **290**: 2309–2312.
- Yanagi K-I, Mizuno T, You Z, Hanaoka F. 2002. Mouse geminin inhibits not only Cdt1-MCM6 interactions but also a novel intrinsic Cdt1 DNA binding activity. *J Biol Chem* **277**: 40871–40880.
- Yardimci H, Loveland AB, Habuchi S, van Oijen AM, Walter JC. 2010. Uncoupling of sister replisomes during eukaryotic DNA replication. *Mol Cell* **40**: 834–840.
- Ying CY, Gautier J. 2005. The ATPase activity of MCM2–7 is dispensable for pre-RC assembly but is required for DNA unwinding. *EMBO J* **24**: 4334–4344.
- You Z, Masai H. 2008. Cdt1 forms a complex with the minichromosome maintenance protein (MCM) and activates its helicase activity. *J Biol Chem* **283**: 24469–24477.
- Yu Z, Feng D, Liang C. 2004. Pairwise interactions of the six human MCM protein subunits. *J Mol Biol* **340**: 1197–1206.
- Zegerman P, Diffley JFX. 2007. Phosphorylation of Sld2 and Sld3 by cyclin-dependent kinases promotes DNA replication in budding yeast. *Nature* **445**: 281–285.
- Zhang J, Yu L, Wu X, Zou L, Sou KKL, Wei Z, Cheng X, Zhu G, Liang C. 2010. The interacting domains of hCdt1 and hMcm6 involved in the chromatin loading of the MCM complex in human cells. *Cell Cycle* **9**: 4848–4857.
- * Zielke N, Edgar BA, DePamphilis ML. 2013. Endoreplication. *Cold Spring Harb Perspect Biol* **5**: a012948.

Cold Spring Harbor Perspectives in Biology

Helicase Loading at Chromosomal Origins of Replication

Stephen P. Bell and Jon M. Kaguni

Cold Spring Harb Perspect Biol 2013; doi: 10.1101/cshperspect.a010124 originally published online April 23, 2013

Subject Collection [DNA Replication](#)

Replication of Epstein–Barr Viral DNA

Wolfgang Hammerschmidt and Bill Sugden

Replication Proteins and Human Disease

Andrew P. Jackson, Ronald A. Laskey and Nicholas Coleman

Break-Induced DNA Replication

Ranjith P. Anand, Susan T. Lovett and James E. Haber

Regulating DNA Replication in Eukarya

Khalid Siddiqui, Kin Fan On and John F.X. Diffley

Archaeology of Eukaryotic DNA Replication

Kira S. Makarova and Eugene V. Koonin

Translesion DNA Polymerases

Myron F. Goodman and Roger Woodgate

Human Papillomavirus Infections: Warts or Cancer?

Louise T. Chow and Thomas R. Broker

Chromatin and DNA Replication

David M. MacAlpine and Geneviève Almouzni

Endoreplication

Norman Zielke, Bruce A. Edgar and Melvin L. DePamphilis

Replication-Fork Dynamics

Karl E. Duderstadt, Rodrigo Reyes-Lamothe, Antoine M. van Oijen, et al.

Helicase Activation and Establishment of Replication Forks at Chromosomal Origins of Replication

Seiji Tanaka and Hiroyuki Araki

Poxvirus DNA Replication

Bernard Moss

The Minichromosome Maintenance Replicative Helicase

Stephen D. Bell and Michael R. Botchan

DNA Replication Origins

Alan C. Leonard and Marcel Méchali

Principles and Concepts of DNA Replication in Bacteria, Archaea, and Eukarya

Michael O'Donnell, Lance Langston and Bruce Stillman

DNA Replication Timing

Nicholas Rhind and David M. Gilbert

For additional articles in this collection, see <http://cshperspectives.cshlp.org/cgi/collection/>

**All Modifications and
Oligo Types Synthesized**
Long Oligos • Fluorescent • Chimeric • DNA • RNA • Antisense

Oligo Modifications?
Your wish is our command.