The Meiotic Checkpoint Network: Step-by-Step through Meiotic Prophase

Vijayalakshmi V. Subramanian and Andreas Hochwagen

Department of Biology, New York University, New York, New York 10003

Correspondence: andi@nyu.edu

The generation of haploid gametes by meiosis is a highly conserved process for sexually reproducing organisms that, in almost all cases, involves the extensive breakage of chromosomes. These chromosome breaks occur during meiotic prophase and are essential for meiotic recombination as well as the subsequent segregation of homologous chromosomes. However, their formation and repair must be carefully monitored and choreographed with nuclear dynamics and the cell division program to avoid the creation of aberrant chromosomes and defective gametes. It is becoming increasingly clear that an intricate checkpoint-signaling network related to the canonical DNA damage response is deeply interwoven with the meiotic program and preserves order during meiotic prophase. This meiotic checkpoint network (MCN) creates a wide range of dependent relationships controlling chromosome movement, chromosome pairing, chromatin structure, and double-strand break (DSB) repair. In this review, we summarize our current understanding of the MCN. We discuss commonalities and differences in different experimental systems, with a particular emphasis on the emerging design principles that control and limit cross talk between signals to ultimately ensure the faithful inheritance of chromosomes by the next generation.

Meiosis is a specialized chromosome segregation process, wherein a diploid parent cell gives rise to haploid gametes (Kleckner 1996; Petronczki et al. 2003; Gerton and Hawley 2005). The reduction in ploidy is essential for gametogenesis in all sexually reproducing organisms and is achieved by a single round of DNA replication followed by two chromosome segregation events that uniquely segregate not only sister chromatids but also homologous chromosomes. Separation of homologous chromosomes occurs during the first meiotic division (meiosis I), followed by the separation of sister chromatids during meiosis II.

The mechanics of chromosome segregation require that chromosome pairs that are to be segregated must first be connected to each other to ensure their proper orientation on the spindle (Miller et al. 2013). Just like during mitosis, meiotic sister chromatids are held together by sister chromatid cohesion that is established when the diploid genome is duplicated during premeiotic S phase (Fig. 1A). However, no such a priori linkage exists for homologous chromosomes. Consequently, a major mechanistic challenge of meiosis is to identify homologous chromosome pairs and establish connections between them. Much of meiotic prophase, the extended

Editors: Stephen Kowalczykowski, Neil Hunter, and Wolf-Dietrich Heyer

Additional Perspectives on DNA Recombination available at www.cshperspectives.org

Copyright © 2014 Cold Spring Harbor Laboratory Press; all rights reserved; doi: 10.1101/cshperspect.a016675

Cite this article as *Cold Spring Harb Perspect Biol* 2014;6:a016675

V.V. Subramanian and A. Hochwagen

Figure 1. A crossover establishes a physical link between homologous chromosomes. (A) Schematic of a pair of homologous chromosomes (red and purple). The replicated sister chromatids are held together by cohesion (green rings). (B) A crossover between homologous chromosomes, in conjunction with cohesion distal to the crossover site, establishes a physical connection between them. (C) A crossover allows homologous chromosomes to orient properly on the meiotic spindle (gray lines).

G₂ phase preceding meiosis I, is dedicated to achieving this goal.

In most organisms, the connections between homologous chromosomes are established by crossover recombination (Fig. 1B). Crossovers exchange covalent links between sequences of homologous chromosomes, and in conjunction with crossover-distal sister chromatid cohesion, provide the physical connections necessary for homologous chromosome segregation during meiosis I (Fig. 1C) (van Heemst and Heyting 2000; Lee and Orr-Weaver 2001). Crossover recombination is initiated after premeiotic DNA replication with the programmed introduction of numerous DNA DSBs by the conserved SPO11 enzyme (Fig. 2A) (Keeney 2001). Removal of SPO11 and 5' resection of DSB ends produces 3' single-strand-

ed DNA (ssDNA) ends that are used by the strand-invasion proteins RAD51 and DMC1 to search for homologous repair templates (Neale and Keeney 2006). In meiosis, a distinct inter-homolog (IH) bias is generated to promote crossover recombination between homologous chromosomes rather than sister chromatids (Hollingsworth 2010; Lao and Hunter 2010). Moreover, a process known as crossover interference ensures an even distribution of crossovers before stable strand-invasion interactions with the homolog (Carpenter and Sandler 1974; Bishop and Zickler 2004; Berchowitz and Copenhagen 2010). Only stabilized strand-invasion intermediates are processed into double-Holliday junctions and ultimately resolved as crossovers, whereas the remaining intermediates are displaced from the homolog to be repaired as non-

Figure 2. Meiotic DNA replication and DSB repair events occur concomitantly with chromosome structural morphogeneses. Schematic of DNA metabolism (A), and chromosomal organization events (B) during meiotic prophase. The homologous chromosomes replicate during premeiotic S phase. At leptotema, the DSBs are initiated, whereas telomeres of the chromosomes become tethered to the nuclear envelope and the meiotic chromosomes assume a bouquet conformation (in most organisms). Synapsis (depicted by gray lines) between homologous chromosome pairs is thought to initiate at sites of crossover repair in zygonema. By pachynema the homologous chromosomes are fully synapsed and the crossover-designated repair is at the double-Holliday junction intermediate stage. The synaptonemal complex disassembles at diplonema to reveal the crossover sites between the homologous chromosomes.

V.V. Subramanian and A. Hochwagen

crossovers (Allers and Lichten 2001; Hunter and Kleckner 2001).

DSB formation and repair are facilitated by chromosome structure transitions that are easily observable by cytology and underlie the cytologically defined stages of meiotic prophase—leptonema, zygonema, pachynema, and diplonema (Fig. 2B) (Baarends and Grootegoed 2003; Storlazzi et al. 2003). Meiotic chromosome morphogenesis initiates concomitantly with DNA replication with the assembly of proteinaceous chromosome axes that give each chromosome a rod-like center with emanating chromatin loops (Klein et al. 1999; Blat et al. 2002; Panizza et al. 2011; Borde and de Massy 2013). The loop-axis organization is completed in leptonema and is important for DSB formation as well as for establishing IH bias (Blat et al. 2002; Storlazzi et al. 2003; Carballo et al. 2008; Kim et al. 2010; Hong et al. 2013). As cells progress through zygonema, homologous chromosomes pair, their axes align, and in many organisms, the chromosomes progressively synapse. Synapsis refers to the assembly of a tripartite proteinaceous scaffold called the synaptonemal complex (SC) that is formed by central transverse filaments laid down between the paired axes of the homologous chromosomes (Page and Hawley 2004; Fraune et al. 2012). DSBs that have been designated to become crossovers are suggested to be the sites of synapsis initiation, in addition to synapsis initiation at the centromeres in some organisms (Klein et al. 1999; Henderson and Keeney 2004; Tsubouchi and Roeder 2005; Obeso and Dawson 2010; Subramanian and Hochwagen 2011). The final stages of crossover recombination occur in the context of the SC. When all chromosomes achieve full-length synapsis, the cells are in pachynema. By the subsequent diplonema, the cells have completed repair and disassemble their SC as they prepare to segregate the homologous chromosomes.

Alongside these chromosomal transitions, nuclear organization is often found to undergo remarkable changes (Fig. 2B). The specific nuclear restructuring varies between organisms and can take the form of telomere clustering in the nuclear envelope (the “bouquet” stage observed in many organisms), the subnuclear

congression of chromosomes, as observed in *Caenorhabditis elegans* and *Drosophila melanogaster*, or a dramatic elongation of the nucleus, as seen in *Tetrahymena thermophila* (Scherthan 2001; Sheehan and Pawlowski 2009; Takeo et al. 2011; Tanneti et al. 2011; Loidl et al. 2012; Woglar and Jantsch 2013). In addition, chromosomes often undergo periods of extraordinary dynamicity, exemplified by the “horsetail movement” in *Schizosaccharomyces pombe* and the rapid pachytene movements in *Saccharomyces cerevisiae* or maize (Ding et al. 1998; Tomita and Cooper 2006; Koszul et al. 2009; Sheehan and Pawlowski 2009; Sonntag Brown et al. 2011; Lee et al. 2012). These processes typically occur in a stage-specific manner and, in most instances, are thought to either help chromosome pairing or resolve unproductive chromosomal interactions (Koszul and Kleckner 2009).

Work over the past several years has revealed that meiotic cells rely on an intricate network of signaling mechanisms to coordinate this complex program and create dependencies between different processes (Roeder and Bailis 2000; Hochwagen and Amon 2006; Longhese et al. 2009; MacQueen and Hochwagen 2011). These dependencies are necessary to establish the correct timing of meiotic prophase events and to avoid deleterious interactions between different processes. They also provide an opportunity to delay or even cull meiotic cells if meiotic processes go awry. Here, we attempt to summarize our current understanding of this network of dependencies. In an effort to simplify, we will refer to the checkpoint components by their human homologs wherever possible and indicate the organism-specific nomenclature in superscript when referencing the function in an organism-specific context.

A WEB OF DEPENDENCIES CREATES ORDER IN MEIOTIC PROPHASE

Throughout this review, we refer to the overall signaling network comprising these mechanisms as the meiotic checkpoint network (MCN). In line with the original definition of cell cycle checkpoints (Hartwell and Weinert 1989), we use the term “checkpoint mechanism”

to describe any signaling mechanism that creates a dependent relationship between metabolically independent meiotic processes (e.g., DSB formation and SC assembly). This broad definition is intended to emphasize that checkpoints are not primarily surveillance mechanisms that respond to abnormal events. Although meiotic DSBs are a form of genome damage, their formation is an inherent part of every meiotic prophase, and thus per se not abnormal. Consequently, we view the MCN not simply a damage response network, but as an integral coordinating mechanism that is central to the ordered execution of meiotic prophase.

Figure 3 provides a broad overview of our current understanding of the MCN. By far, the most dependencies arise from the formation of DSBs, presumably reflecting the inherent dangers associated with chromosome breakage. However, some processes are also linked to the completion of DNA replication or the proper pairing and synapsis of chromosomes. Remarkably, almost all currently known dependencies in meiotic prophase involve the activities of the conserved PI3-like kinases ATM and ATR. This means that the MCN must have mechanisms to differentiate between signals to elicit the appropriate responses. We discuss this important feature of the MCN in a later part of this review, but first focus on the general architecture of the MCN.

THE MAIN PLAYERS

The core signaling machinery of the MCN uses many of the players of the canonical DNA damage response (DDR) network (Table 1), including the conserved checkpoint sensor kinases ATM and ATR (MacQueen and Hochwagen 2011). ATM and ATR are evolutionarily related serine/threonine kinases that become activated by distinct forms of DNA damage as well as by asynapsis during meiosis (Carballo and Cha 2007; Burgoyne et al. 2009). ATM responds primarily to blunt and protein-conjugated DSB ends, whereas ATR is activated by RPA-coated ssDNA resulting from DSB processing, as well as ssDNA/dsDNA junctions (Harrison and Haber 2006; Lovejoy and Cortez 2009). Both kinases rely on the activity of cofactors for damage rec-

ognition. ATM detects blunt ends with the help of the MRN complex (MRE11-RAD50-NBS1) (Usui et al. 2001; Nakada et al. 2003; You et al. 2005). ATR detects ssDNA through its activator ATRIP, and ssDNA/dsDNA junctions through the PCNA-like 9-1-1 complex (RAD9-RAD1-HUS1) (Zou and Elledge 2003; Harrison and Haber 2006; Refolio et al. 2011). In addition, the cofactors BRCA1 and TOPBP1 promote ATR activity in response to unsynapsed meiotic chromatin (Refolio et al. 2011; Royo et al. 2013). ATM and ATR phosphorylate large and often overlapping sets of substrates on serine-glutamine (SQ) or threonine-glutamine (TQ) dipeptides. Many of the known effectors of the MCN are direct targets of ATM/ATR (Table 2), creating immediate links between signal and outcome. In addition, ATM/ATR activate the CHK1 and CHK2 effector kinases, which further relay checkpoint signals but typically control a more restricted set of processes.

CONTROL OF DSB FORMATION

Ongoing Replication Blocks DSB Formation

The first known checkpoint mechanism in meiotic prophase is the meiotic replication checkpoint. As in mitotic cells, a primary function of the meiotic replication checkpoint is the maintenance of replication potential, which occurs through ATR- and CHK2-dependent stabilization of replication forks (Branzei and Foiani 2010; Blitzblau and Hochwagen 2013). However, in addition, the meiotic replication checkpoint also prevents DSB formation as long as replication is ongoing (Figs. 3 and 4A) (Tonami et al. 2005; Ogino and Masai 2006; Blitzblau and Hochwagen 2013). The enforced temporal separation of replication and DSB formation is important because it ensures that crossovers only form between replicated chromosomes (see Fig. 1). Moreover, it prevents lethal conflicts between DSB formation and DNA replication (Blitzblau and Hochwagen 2013). The replication checkpoints of *S. pombe* and *S. cerevisiae* suppress DSBs through transcriptional repression of essential regulators of DSB formation, although the identity of the ultimate checkpoint

V.V. Subramanian and A. Hochwagen

A Normal meiotic progression **B** Pathological responses

Figure 3. Dependent relationships established by the MCN. The meiotic checkpoint network creates a web of dependencies to promote sequential progression of meiotic events (A), or prevent meiotic progression in the face of defective repair or synapsis (B). Dashed lines and arrows indicate a modulation in activity.

target differs between the two yeasts. In *S. pombe* expression of Mde2, an axis associated DSB regulator, is down-regulated by the checkpoint, whereas expression of SPO11 itself is under the control of the replication checkpoint in the budding yeast (Ogino and Masai 2006; Miyoshi et al. 2012; Blitzblau and Hochwagen 2013). In

addition, the replication checkpoint of *S. cerevisiae* also directly controls the chromosomal localization and activation of other components of the DSB machinery (MER2 and REC114). Signaling in this case occurs both through $\text{CHK2}^{\text{Rad53}}$ -dependent and independent mechanisms and involves regulation of the conserved

Table 1. MCN proteins and their homologs

Mammals	<i>S. cerevisiae</i>	<i>C. elegans</i>	<i>S. pombe</i>	<i>Drosophila</i>	<i>Arabidopsis</i>	Function
Signaling proteins						
ATR	Mec1	ATL-1	Rad3p	Mei-41	ATR	PI3 ^a kinase-like kinase
ATM	Tel1	ATM-1	Tellp	Atm	ATM	PI3 kinase-like kinase
RAD9A, RAD9B	Ddc1	HPR-9	Rad9p	Rad9A, Rad9B	–	PCNA ^b -like clamp (9-1-1 complex)
RAD1	Rad17	MRT-2	Rad1p	Rad1		PCNA-like clamp (9-1-1 complex)
HUS1 HUS1B	Mec3	HUS-1	Hus1p	Hus1	–	PCNA-like clamp (9-1-1 complex)
CHK1	(Chk1)	CHK-1	Chk1p	(Grp)	–	Protein kinase
CHK2	Rad53 Mek1	CHK-2	Cds1p Mek1p	Mnk	–	Protein kinase with FHA ^c domain
Several	Cdc5	PLK-2	Plo1p	Polo	–	Protein kinase
HORMAD1 HORMAD2	Hop1	HTP-1 HTP-2 HIM-3	Hop1	–	ASY1	Chromosomal HORMA-domain proteins
SYCP3	Red1?	HTP-3?	Rec10?	C(2)M?	ASY3?	Chromosome axis component
Several	Sir2	SIR-2	Sir2p	Sir2	SRT1 SRT2	NAD-dependent deacetylase
TRIP13	Pch2	PCH-2	–	Pch2	–	AAA ⁺ -ATPase ^d

Proteins shown to be involved in the MCN pathway are in bold. Proteins that were shown not to be part of the MCN function are in parentheses. Proteins that share functional homology but no obvious sequence homology are followed by “?”.

^aPhosphoinositide 3-kinase.

^bProliferating cell nuclear antigen.

^cFork-head associated domain.

^dATPases associated with diverse cellular ATPase.

cell cycle kinase DDK (Blitzblau and Hochwagen 2013). A role for CHK2 in coordinating premeiotic DNA replication and subsequent meiotic prophase entry was also suggested for *C. elegans* (MacQueen and Villeneuve 2001). A possibly even earlier meiotic role of ATR and CHK2 in DNA replication was recently suggested in *S. pombe*. In this organism, mutation of an ATR/ATM- and SPO11^{Rec12}-dependent phosphorylation site on CHK2^{Mek1} resulted in a delay in DNA replication (Tougan et al. 2010). Although this phosphorylation event may represent feedback regulation of replication by DSBs, this possibility requires further study as deletion of ATR^{Rad3} or CHK2^{Mek1} did not similarly affect S-phase progression (Ogino and Masai 2006).

DSB Levels—A Balancing Act

In addition to linking DSB formation to the sufficient completion of DNA replication, there

is increasing evidence that the MCN also feeds back to modulate DSB levels once DSB formation has initiated. In mouse and *Drosophila*, loss of ATM leads to increased abundance of DSB markers, whereas loss of ATR causes similar phenotypes in *A. thaliana*, suggesting that DSB-dependent activation of these kinases down-regulates further DSB formation (Joyce et al. 2011; Lange et al. 2011; Kurzbauer et al. 2012). Meiotic DSB levels must be tightly controlled, as an excessive DSB load can lead to severe problems in DNA repair (Johnson et al. 2007). Indeed, a number of meiotic phenotypes of *Atm*^{−/−} mice can be alleviated by reducing *SPO11* copy number (Bellani et al. 2005; Barchi et al. 2008). A particular function of ATM in this context may be to prevent repeated DSB formation at the same chromosomal locus (including the sister chromatid). Spatial proximity of DSBs could explain why *Atm*^{−/−} mice display a strong increase in the amount of postcleavage SPO11-

Table 2. Phosphorylation events involved in creating dependencies

Modifying enzyme(s)	Target	Phosphorylated residue(s)	Organism	Effect(s)	References
ATR ^{Mec1} /ATM ^{Tel1}	CtIP ^{Sae2}	(Ser73) (Thr90) Ser249 Ser279 Ser289	<i>S. cerevisiae</i>	Activation of DSB resection	Cartagena-Lirola et al. 2006; Terasawa et al. 2008
ATR ^{Mec1} /ATM ^{Tel1}	HORMAD ^{Hop1}	Ser298 (Ser311) Thr318	<i>S. cerevisiae</i>	Inhibition of intersister repair	Carballo et al. 2008
ATR	HORMAD1	Ser375	Mouse	MSCI, signaling of asynapsis?	Fukuda et al. 2012; Royo et al. 2013
ATR ATR ^{Mec1} /ATM ^{Tel1}	HORMAD2 RPA2 ^{Rfa2}	Ser271 Ser122	Mouse <i>S. cerevisiae</i>	MSCI? Altered crossover distribution in some intervals	Royo et al. 2013 Bartrand et al. 2006
ATR ^{Mec1} /ATM ^{Tel1}	RNF212 ^{Zip3}	Up to four residues	<i>S. cerevisiae</i>	Altered crossover levels in some intervals	Serrentino et al. 2013
ATR ^{Mec1} /ATM ^{Tel1}	SCP1 ^{Zip1}	Ser75	<i>S. cerevisiae</i>	Dissolution of centromere pairing	Falk et al. 2010
ATM	Histone H2AFX	Ser139	Mouse	Persistence of the bouquet	Fernandez-Capetillo et al. 2003
ATR	Histone H2AFX	Ser139	Mouse	MSCI	Royo et al. 2013
ATR ^{Mec1} /ATM ^{Tel1}	Rec114	Thr175 Ser187 (Ser256)	<i>S. cerevisiae</i>	Reduced DSB formation	Carballo et al. 2008
ATR ^{Mec1} /ATM ^{Tel1}	CHK2 ^{Mek1}	(Ser12) (Ser14) Thr15	<i>S. pombe</i>	Activation of CHK2 ^{Mek1} kinase activity	Tougan et al. 2010
CHK2 ^{Mek1}	Rad54	Thr132	<i>S. cerevisiae</i>	Inhibition of intersister repair	Niu et al. 2009
CHK2 ^{Mek1}	Rdh54	Thr6 Thr673	<i>S. pombe</i>	Inhibition of intersister repair?	Tougan et al. 2010
CHK2 ^{Mek1}	Mus81	Thr281 Thr422	<i>S. pombe</i>	Inhibition of intersister repair?	Tougan et al. 2010
CHK2 ^{Mek1}	Histone H3	Thr11	<i>S. cerevisiae</i>	Inhibition of intersister repair?	Govin et al. 2010
CHK2 ^{Mek1}	Cdc25	Up to nine residues	<i>S. pombe</i>	DSB-dependent nuclear exclusion of Cdc25; cell cycle delay	Perez-Hidalgo et al. 2008
CHK2	SUN-1	Ser8 Ser12 ^a Ser24 Ser35/Thr36 Ser43 Ser58 Ser62	<i>C. elegans</i>	Chromosome pairing and synapsis initiation	Penkner et al. 2009

^aSer12 phosphorylation is likely indirect because it also depends on PLK-2 (Labella et al. 2011).

Meiotic Checkpoint Network

Figure 4. The meiotic checkpoint network integrates signal to the appropriate response. (A) Stalled replication forks prevent DSB formation via several mechanisms in *S. cerevisiae*. Mec1 regulates transcription of *SPO11* and recruitment of Rec114 to the meiotic chromosomes, whereas the downstream Rad53 kinase controls phosphorylation of Mer2 by regulating the activity of DDK kinase. (B) MCN regulates resection. Both Tel1 and Mec1 kinases activate Sae2 for DSB end resection to generate 3' ssDNA overhangs. The MCN also prevents hyper-resection of break ends. (C) Mec1/Tel1 kinases promote IH bias via phosphorylation of Hop1, which in turn leads to recruitment, dimerization, and activation of Mek1 kinase. Regulation of Rad54 activity by Mek1 inhibits IS repair thus promoting IH bias. (D) In *Drosophila*, the MCN negatively regulates NHK-1 kinase. NHK-1 kinase controls condensation of the oocyte chromatin and also allows its release from the nuclear envelope on completion of DSB repair. (E) Unsynapsed chromatin in mouse recruits ATR via HORMAD1/2. ATR facilitates phosphorylation of H2AFX that spreads into the chromatin loops and recruits silencing factors. (F) MCN regulates exit from meiotic prophase by controlling the expression and localization of Ndt80 transcription factor as well as by inhibiting CDK kinase. Cdc5 kinase relieves inhibition of Ndt80 by the MCN in a feedforward loop to allow rapid exit from prophase.

V.V. Subramanian and A. Hochwagen

oligonucleotide complexes, but only a mild increase in the number of cytologically discernable RAD51 foci (Barchi et al. 2008; Lange et al. 2011). Analyses of recombinant chromatids from *S. cerevisiae* tetrads at specific DSB sites in mutants lacking ATM^{Tel1} or ATR^{Mec1} also support this model (Zhang et al. 2011). Although the target of ATM in this context remains to be identified, recent experiments in *S. cerevisiae* suggest the conserved SPO11 accessory factor REC114 as a promising candidate. REC114 is a substrate of ATM^{Tel1} and ATR^{Mec1} and mutations mimicking constitutive ATM/ATR-dependent phosphorylation cause a notable decrease in DSB levels (Carballo et al. 2013). CHK2-dependent regulation of the DSB regulators DSB-1 and DSB-2 may have the equivalent function in *C. elegans* (Rosu et al. 2013; Stamper et al. 2013).

A number of recent studies in *S. cerevisiae* indicate that defects in DSB repair further modulate DSB levels. The effects are rather complex as the MCN shows both DSB-promoting and DSB-suppressing effects depending on the amount of DSBs formed, the type of repair defect, and whether cells are able to prematurely exit prophase (Argunhan et al. 2013; Blitzblau and Hochwagen 2013; Carballo et al. 2013; Gray et al. 2013; Lao et al. 2013; Rockmill et al. 2013). Finally, work in yeast and mice also suggests a feedback between DSB formation and homolog interactions, as DSB formation continues on unsynapsed chromosomes (Kauppi et al. 2013a,b; Thacker et al. 2014).

CONTROL OF DSB REPAIR

Activation of DSB End Processing

DSB formation itself triggers a major activation of the MCN (Fig. 3). One of the first events following meiotic DSB formation is MRN/CtIP-initiated end resection, which promotes homologous recombination and also creates a barrier to error-prone end-joining mechanisms of repair (Joyce et al. 2012; Yin and Smolikove 2013). Resection is initiated by MRE11-dependent endonucleolytic incisions near DSBs, followed by bidirectional resection that requires both MRN and EXO1 (Zakharyevich et al.

2010; Garcia et al. 2011). In *S. cerevisiae*, the MRN^{Xrs2} complex detects unprocessed meiotic DSB ends and activates ATM^{Tel1} kinase, which in turn phosphorylates the MRN interacting protein CtIP^{Sae2} to initiate DSB resection (Fig. 4B) (Usui et al. 2001; Cartagena-Lirola et al. 2006; Terasawa et al. 2008). In a positive feedback loop, resected DNA ends lead to the activation of ATR^{Mec1} , which further contributes to CtIP^{Sae2} activation. However, this dependency is not strictly linear because ATR^{Mec1} also becomes activated independently of ATM^{Tel1} , and is itself sufficient to phosphorylate CtIP^{Sae2} and initiate resection (Cartagena-Lirola et al. 2008).

Meiotic resection initially is limited, but if DSB repair is blocked, meiotic cells enter a phase of DSB hyperresection. Intriguingly, ATR^{Mec1} and the 9-1-1 complex are also required to restrain hyperresection (Shinohara et al. 2003; Gray et al. 2013; Clerici et al. 2014). Given that a number of nucleases are involved in the resection process (Mimitou and Symington 2009; Zakharyevich et al. 2010; Garcia et al. 2011; Schaetzlein et al. 2013), an appealing model is that the MCN ensures appropriate resection rates by activating some nucleases, while (temporarily) inhibiting others (Segurado and Diffley 2008; Manfrini et al. 2010; Luo et al. 2013; Souquet et al. 2013). In *S. cerevisiae*, resection by BLM^{Sgs1}/DNA2, in particular, is likely only activated late in meiosis (Manfrini et al. 2010; Zakharyevich et al. 2010).

Suppression of Intersister Recombination

For meiotic DSBs to support crossover formation between homologous chromosomes, repair from the more readily available homologous sequences on the sister chromatid must be suppressed. Several mechanisms act in concert to achieve this goal, both by down-regulating sister-directed RAD51-recombinase activity and by promoting the homolog as the preferred repair template (Kim et al. 2010; Lao and Hunter 2010; Kurzbauer et al. 2012; Hong et al. 2013; Lao et al. 2013; Liu et al. 2014). Research in a number of organisms indicates a central role of the MCN in establishing meiotic homolog bias (Carballo et al. 2008; Latypov et al. 2010), al-

though the mechanistic details are best understood in *S. cerevisiae* (Fig. 4C). In this organism, ATM^{Tel1}/ATR^{Mec1} phosphorylate the HORMA-domain-containing chromosomal axis protein HORMAD^{Hop1}, the homolog of mammalian HORMAD1/2, on several clustered S/TQ sites (Table 1) (Carballo et al. 2008). This leads to the recruitment, dimerization, and activation of the CHK2-like effector kinase CHK2^{Mek1} (Niu et al. 2005, 2007; Carballo et al. 2008; Wu et al. 2010) whose binding, in turn, stabilizes the phosphorylation mark on HORMAD^{Hop1} (Chuang et al. 2012). Once activated, CHK2^{Mek1} kinase promotes IH bias possibly in part by phosphorylating and inhibiting RAD54, a SWI/SNF-family ATPase that stimulates RAD51-recombinase activity for repair from the sister chromatid (Niu et al. 2009). However, genetic experiments suggest that other (currently unknown) targets of CHK2^{Mek1} provide the primary mechanism to promote IH bias (Niu et al. 2009; Terentyev et al. 2010). Research in *S. cerevisiae* and *S. pombe* has identified several additional CHK2^{Mek1} targets, including a chromatin mark (histone H3 T11), the RAD54-related yeast protein Rdh54, and the resolvase MUS81 (Govin et al. 2010; Tougan et al. 2010). So far, only Rdh54 has been excluded as a likely functional target of the MCN (Niu et al. 2009). Notably, Rdh54 phosphorylation during vegetative growth is implicated in checkpoint adaptation (Ferrari et al. 2013).

Suppression of Ectopic Recombination

There is evidence that the MCN also protects genome stability by preventing nonallelic (ectopic) recombination. Mutants of the plant *Arabidopsis thaliana* lacking both ATM and ATR display DSB-dependent associations between nonhomologous chromosomes that persist into metaphase I, suggestive of ectopic crossover recombination (Culligan and Britt 2008). Moreover, increased ectopic recombination is also observed in *S. cerevisiae* cells lacking a functional 9-1-1 complex and mice lacking the 9-1-1 component HUS1 (Grushcow et al. 1999; Thompson and Stahl 1999; Shinohara et al. 2003; Lyndaker et al. 2013a; Shinohara and Shinohara 2013). Although the relevant check-

point targets remain unknown, the checkpoint network may restrain ectopic recombination by coordinating the two DSB ends (Shinohara and Shinohara 2013), a notion supported by the observation that RAD51 and DMC1 recombinases frequently appear in an abnormal side-by-side arrangement in 9-1-1 complex mutants of *S. cerevisiae* (Shinohara et al. 2003). The increased number of RAD51 and DMC1 foci in ATR mutants of *A. thaliana* may reflect a similar defect, although the geometry of recombinase loading appears to differ between the two organisms (Kurzbaue et al. 2012). It is possible that increased ectopic recombination is responsible for the decreased crossover levels observed in a number of checkpoint mutants (Shinohara et al. 2003). However, in many cases mutations in checkpoint factors also show a strong delay in DSB repair, which may argue for a more direct role of the meiotic checkpoint machinery in promoting meiotic recombination (Shimada et al. 2002; Shinohara et al. 2003; Peretz et al. 2009; Joyce and McKim 2010).

Obligate Crossover Formation and Crossover Interference

Finally, there is limited evidence that the MCN has a role in regulating crossover distribution. Several processes are at work to ensure that each homologous chromosome pair receives a crossover (the obligate crossover), and that neighboring crossovers do not occur too close to each other (crossover interference). In male mice, ATM activity is required for the obligate crossover in the small pseudoautosomal region of homology that allows pairing between X and Y chromosomes (Barchi et al. 2008). In addition, mice lacking ATM have increased autosomal crossover numbers accompanied by reduced crossover interference (Barchi et al. 2008). Similarly, in *S. cerevisiae*, ATM^{Tel1}/ATR^{Mec1}-dependent phosphorylation of the Rfa2 subunit of RPA, as well as of the SC component RNF212^{Zip3}, alters crossover distribution in some genetic intervals (Bartrand et al. 2006; Serrentino et al. 2013), although the generality of these effects remains to be determined. A crossover interference defect was also observed

V.V. Subramanian and A. Hochwagen

in *S. cerevisiae* mutants lacking the phosphatase PP4, which is responsible for the dephosphorylation of several ATR/ATM substrates (Falk et al. 2010). However, the mechanism by which the MCN influences crossover distribution has so far remained elusive.

NUCLEAR ORGANIZATION, PAIRING, AND SYNAPSIS

Checkpoint Control of Nuclear Restructuring

In addition to coordinating DSB repair, the MCN links meiotic nuclear dynamics to both DNA replication and DSB metabolism. In *S. cerevisiae*, one of the first instances of nuclear restructuring, the dispersal of mitotic telomere clusters, is linked to S-phase completion (Trelles-Sticken et al. 2005a), and at least under some circumstances requires ATR^{Mec1} activity (Trelles-Sticken et al. 2005b). A related connection between S-phase and nuclear restructuring may exist in *C. elegans*. Following S phase in this organism, chromosomes aggregate in a polarized nuclear crescent with their telomeres anchored in clusters in the nuclear envelope, reminiscent of the bouquet stage seen in many organisms. This nuclear reorganization requires CHK2, although not ATM/ATR (MacQueen and Villeneuve 2001; Penkner et al. 2009). Thus, how CHK2 is activated in this case is unclear. CHK2 activity is required for multiple events in this context, including the enrichment of ZIM proteins at chromosomal pairing centers and the phosphorylation of the chromosome anchor SUN-1 at the nuclear envelope (Phillips and Dernburg 2006; Penkner et al. 2009). Dephosphorylation of SUN-1, in turn, is required for the dissolution of the polarized crescent as cells enter pachynema (Penkner et al. 2009; Woglar et al. 2013). A checkpoint-dependent restructuring of the nucleus into a bouquet-like state is also observed in *Tetrahymena*, although this process requires both DSB formation and ATR (Loidl and Mochizuki 2009; Loidl et al. 2012).

One of the best-understood mechanisms of MCN-dependent nuclear reorganization in this

context is the release of chromosomes from the nuclear envelope in *Drosophila* (Fig. 4D). Chromosomal release occurs on completion of meiotic recombination and leads to the formation of a compact chromosome cluster called the karyosome. Before DSB repair, karyosome formation is prevented by the ATR^{Mei-41} and CHK2^{Mnk}-dependent inhibition of NHK-1 kinase (Ghabrial and Schupbach 1999; Abdu et al. 2002; Lancaster et al. 2010). One of the substrates of NHK-1 is the nuclear envelope protein BAF, which must be phosphorylated to release chromosomes into the nucleus (Lancaster et al. 2007). NHK-1 is also required for histone H2A Thr119 phosphorylation, SC disassembly, and condensin loading, which may further contribute to karyosome formation (Ivanovska et al. 2005; Lancaster et al. 2010).

Chromosome Pairing and the Bouquet

The bouquet stage coincides with the active pairing of homologous chromosomes, and a failure to properly pair is associated with delayed dispersal of the chromosomal bouquet in many organisms. For example, the presence of an extra chromosome extends the bouquet stage and alters repair dynamics in trisomic human oocytes (Roig et al. 2005; Robles et al. 2013). Similarly, telomere clusters persist in pairing-defective *spo11* mutants of *S. cerevisiae* and *Sordaria* (Trelles-Sticken et al. 1999; Storlazzi et al. 2003). DSB processing is also necessary for exit from the bouquet stage in *Tetrahymena* (Loidl et al. 2012). At least in some cases, these delays may depend on checkpoint regulation, as mouse mutants lacking ATM or its substrate histone H2AFX (formerly known as H2AX) fail to exit the bouquet stage (Fernandez-Capetillo et al. 2003; Liebe et al. 2006). Finally, in *S. cerevisiae*, the MCN also destabilizes the nonhomologous pairing of centromeres in response to DSBs through ATR^{Mec1}-dependent phosphorylation of the central SC component SCP1^{Zip1} (Falk et al. 2010).

Control of Synapsis Initiation

An interesting case of meiotic checkpoint control is the initiation of chromosome synapsis,

which is coupled to chromosome pairing or DSB formation in a variety of organisms. In *C. elegans*, synapsis initiation is blocked by MCN-mediated Ser12-phosphorylation of SUN-1, the nuclear envelope protein that establishes connections between chromosome ends and the cytoskeleton during meiosis. Ser12-phosphorylation of SUN-1 depends on CHK2 and the Polo-like kinase PLK2, but is independent of ATM/ATR (Penkner et al. 2009; Labella et al. 2011; Woglar et al. 2013). Erasure of Ser12 phosphorylation, and thus synapsis initiation, requires appropriate DSB repair (Woglar et al. 2013) as well as appropriate pairing interactions between chromosomes, which appear to be monitored by a force-dependent checkpoint mechanism (Penkner et al. 2009; Wynne et al. 2012; Rog and Dernburg 2013). A different mechanism appears to link synapsis initiation to the onset of DSB formation in *S. cerevisiae*. In the absence of DSBs, synapsis initiation at centromeres is actively blocked by a mechanism involving the putative SUMO-ligase RNF212^{Zip3} and the proline isomerase Fpr3 (MacQueen and Roeder 2009). How the DSB signal is transmitted to allow synapsis initiation at centromeres in this situation remains unknown, although RNF212^{Zip3} has recently emerged as a promising MCN substrate (Serrentino et al. 2013).

ASYNAPSIS AND TRANSCRIPTIONAL SILENCING

It is now well established that unsynapsed chromosomes or chromosome segments elicit the activation of several branches of the MCN. Checkpoint signaling is apparent in meiocytes with partial asynapsis, including cells carrying extra chromosomes or chromosome translocations (Mahadevaiah et al. 2008; Burgoyne et al. 2009; Garcia-Cruz et al. 2009; Kouznetsova et al. 2009), and also occurs transiently on late-synapsing chromosomal regions (Blanco-Rodriguez 2012). In *C. elegans*, asynapsis is associated with a delayed exit from the bouquet state (Carlton et al. 2006; Colaiacovo 2006), and can trigger apoptosis (Bhalla and Dernburg 2005).

In mammals, sites of asynapsis are associated with the phosphorylation of several axis

proteins including HORMAD1 and 2 (Fig. 4E) (Fukuda et al. 2012; Royo et al. 2013) and lead to the recruitment of BRCA1, ATRIP, TOPBP1, and ATR to the unsynapsed chromosome axes, followed by the ATR-dependent accumulation of γ -H2AFX (histone H2AFX phosphorylated on Ser139) (Perera et al. 2004; Turner et al. 2005; Burgoyne et al. 2009; Refolio et al. 2011). If asynapsis persists, γ -H2AFX and ATR spread over the entire chromatin with the help of the γ -H2AFX-binding factor MDC1 (Ichijima et al. 2011), and trigger the heterochromatinization and meiotic silencing of unsynapsed chromatin (MSUC). Transcriptional silencing as a consequence of asynapsis is also observed in a number of nonmammalian organisms, including *Neurospora* and *C. elegans* (Shiu et al. 2001; Bean et al. 2004; Checchi and Engebrecht 2011). Depending on which chromosomal regions are silenced, MSUC in mice frequently leads to the loss of spermatocytes, presumably as a result of the depletion of essential survival factors (Burgoyne et al. 2009; Manterola et al. 2009).

MSUC is closely related to the meiotic silencing of sex chromosomes within the sex body by meiotic sex chromosome inactivation (MSCI) (Fig. 5), a physiological process that responds to the unavoidable partial asynapsis of heteromorphic sex chromosomes, but does not lead to cell death (Turner et al. 2006). Sex body formation takes place in late zygonema and is associated with a second wave of γ -H2AFX formation. γ -H2AFX formation occurs in two waves in mouse meiocytes. The first wave coincides with the onset of recombination, is ATM-dependent, and forms foci that are thought to mark DSBs (Mahadevaiah et al. 2001; Barchi et al. 2005; Bellani et al. 2005). In contrast, the second wave of γ -H2AFX formation depends on ATR, marks the remaining unsynapsed chromosomes, and behaves like an MSUC response in that γ -H2AFX and ATR spread across the associated chromatin loops (Mahadevaiah et al. 2001; Turner et al. 2005; Royo et al. 2013). Curiously, although mediated by the DNA-damage sensor kinase ATR, the second wave is independent of SPO11 (Barchi et al. 2005; Bellani et al. 2005). We discuss potential alternative modes of ATR activation later in this review.

V.V. Subramanian and A. Hochwagen

Figure 5. Meiotic chromosome spread from mouse spermatocyte in pachynema depicting MSCI. The XY pair manifests as the sex body (white arrowhead) and is enriched for ATR (red). SCP3 (green) marks the axes of synapsed and unsynapsed chromosomes, DNA is in blue. (Image courtesy of Sarai Pacheco and Ignasi Roig.)

CELL CYCLE PROGRESSION, CELL DEATH, AND LINKS TO DEVELOPMENT

Control of Prophase Exit

Similar to the mitotic DNA damage response, activation of the MCN also triggers a delay in meiotic prophase, presumably to provide sufficient time for the completion of meiotic recombination. A checkpoint-dependent prophase delay or arrest in response to defects in synapsis or DNA repair is apparent in many organisms (Lydall et al. 1996; Shimada et al. 2002; Hochwagen and Amon 2006; Joyce and McKim 2010; Lyndaker et al. 2013a; Woglar et al. 2013), although mechanistic details on how the MCN influences the cell cycle machinery are so far largely restricted to *S. cerevisiae* and *S. pombe*. In both yeasts, the meiotic cell cycle delay is mediated by ATR-dependent activation and dimerization of $\text{CHK2}^{\text{Mek1}}$ (Lydall et al. 1996; Xu et al. 1997; Shimada et al. 2002; Wu et al. 2010). $\text{CHK2}^{\text{Mek1}}$ acts in part through inhibition of cyclin-dependent kinase (CDK). In *S. cerevisiae*, $\text{CHK-2}^{\text{Mek1}}$ kinase phosphorylates and activates

the CDK-inhibitory kinase $\text{WEE1}^{\text{Swe1}}$ (Fig. 4F) (Tung et al. 2000; Acosta et al. 2011), whereas in *S. pombe*, $\text{CHK2}^{\text{Mek1}}$ kinase promotes the nuclear exclusion (and thus inactivation) of the CDK-activating phosphatase CDC25 (Perez-Hidalgo et al. 2008). The outcome in both cases is persistent inhibitory tyrosine phosphorylation of CDK. In parallel, the MCN of *S. cerevisiae* also triggers the nuclear export of the Ndt80 (Hepworth et al. 1998; Wang et al. 2011), a key transcription factor that activates the transcription of a large set of genes including B-type cyclins as well as the prophase-exit promoting kinase PLK^{Cdc5} (Chu and Herskowitz 1998; Sourirajan and Lichten 2008). Tyrosine phosphorylation and repression of cyclins both keep CDK inactive and thus prevent precocious prophase exit while the MCN is active. Recent research and modeling has furthermore shown that Ndt80 and PLK^{Cdc5} are embedded in an intricate system of feedback and feedforward loops that creates a bistable switch for rapid exit from meiotic prophase once the checkpoint network is inactivated (Acosta et al. 2011; Okaz et al. 2012).

Persistent Defects and the Induction of Cell Death

If defects in repair or synapsis persist, the checkpoint networks of various organisms adopt different terminal strategies. *S. cerevisiae* cells enter a prolonged prophase arrest that can be exited by aborting meiosis should environmental conditions become favorable for mitotic growth (Simchen 2009). Alternatively, *S. cerevisiae* can adapt to the damage by inactivating the MCN, and attempt meiosis (Bailis et al. 2000; Hochwagen et al. 2005; Iacovella et al. 2010). In contrast, meiocytes in metazoans are frequently culled by checkpoint-dependent induction of the apoptotic cell death program (Gartner et al. 2000; Bhalla and Dernburg 2005; Di Giacomo et al. 2005), a process that also functions as a screening mechanism against germ cell precursors with chromosomal abnormalities (Ahmed et al. 2013; Stevens et al. 2013; Titen et al. 2014). As in the mitotic DNA damage response, the decision to enter the apoptotic program in response to repair defects requires CHK2 -dependent ac-

tivation of the p53 family of proteins, and is generally restricted to specific stages in meiotic prophase (Derry et al. 2001; Barchi et al. 2005; Suh et al. 2006; Rutkowski et al. 2011; Bolcun-Filas et al. 2014; Kim and Suh 2014). The role of the MCN in this decision has been difficult to define because, in most cases, loss of MCN factors itself causes DSB repair defects, which in turn trigger germ cell death (Barchi et al. 2008; Burgoyne et al. 2009). Conversely, even a relatively downstream factor in the MCN, such as p53, not only regulates apoptosis but also leads to reduced crossover formation in *Drosophila* (Lu et al. 2010). However, disruption of the 9-1-1 complex component HUS1 was recently shown to bypass both pachynema arrest and apoptosis in mouse spermatocytes (Lyndaker et al. 2013a), and disruption of CHK2 has similar effects in mouse oocytes (Bolcun-Filas et al. 2014), supporting a role of the MCN in this decision.

Interestingly, in both mouse and *C. elegans*, the meiotic cell death response shows a profound sexual dimorphism. Mouse spermatocytes experiencing defects in DSB repair or synapsis typically undergo cell death in pachynema, usually in conjunction with defective sex body formation and the resulting aberrant gene expression. In contrast, defective oocytes often proceed through the meiotic divisions (Nagaoka et al. 2011, 2012). Although many are later removed by atresia, surviving oocytes have a substantially higher rate of chromosome abnormalities compared with mature sperm. The reason for this inefficiency in removing aberrant oocytes is unclear. A different sexual dimorphism is observed in worms. *C. elegans* hermaphrodites show a robust apoptotic response to persistent meiotic defects (Gartner et al. 2000; Bhalla and Dernburg 2005). In contrast, *C. elegans* males only initiate the early stages of the apoptotic program but prevent caspase activation (Jaramillo-Lambert et al. 2010). This signaling modification may be linked to the constitutively asynaptic single X chromosome in male worms. Unexpectedly, despite the apoptotic culling mechanism only being active in hermaphrodites, male worms with synapsis defects produce fewer aberrant gametes, indicating the

existence of apoptosis-independent proofing mechanisms in the male (Jaramillo-Lambert et al. 2010).

Links to Development

An alternative to triggering cell death in the face of persistent defects is to prevent the formation of mature gametes. Accordingly, in several organisms, the MCN creates dependencies between DSB repair and subsequent developmental events. One well-studied example occurs in *Drosophila*, in which DSB repair is linked to the developmental patterning of the oocyte. In this organism, persistent DSBs lead to ATR^{mei-41} and CHK2^{Mnk}-dependent modification of Vasa, an RNA helicase required for the translation of *gurken* mRNA and the dorsoventral patterning of the eggshell (Ghabrial et al. 1998; Staeva-Vieira et al. 2003). As a result, the MCN can block oocyte development. Analogously, the presence of persistent DSBs or defective synapsis also leads to an MCN-dependent block of the developmental program for spore formation in several fungi (Tung et al. 2000; Anderson et al. 2012; Guo and King 2013), ultimately preventing chromosomal defects from being passed on to the next generation.

ARCHITECTURAL FEATURES OF THE MCN

From this overview of the different checkpoint branches in meiotic prophase, it is apparent that the MCN is highly interconnected but uses a surprisingly small number of signaling molecules. This raises a number of questions, including how some of the signals are generated in the first place, how an appropriately modulated response is elicited, and how individual dependencies are separated and integrated.

The Context Matters

Increasing evidence suggests that the specialized architecture of meiotic chromosomes plays a fundamental role in shaping the response of the MCN. In many organisms, disruption of components of the meiotic chromosome axes, including SYCP3 and cohesins, leads to a defect in MCN signaling (Wang and Hoog 2006; Kouz-

V.V. Subramanian and A. Hochwagen

netsova et al. 2009; Callender and Hollingsworth 2010; Lightfoot et al. 2011), although in some cases the loss of signal has been attributed to reduced DSB formation (Callender and Hollingsworth 2010). Perhaps not surprisingly, MCN roles have also been reported for several histone methyltransferases (San-Segundo and Roeder 2000; Checchi and Engebrecht 2011; Lamelza and Bhalla 2012; Ontoso et al. 2013a,b), which presumably promote chromosome axis assembly or contribute to the structural environment of MCN signaling. Further supporting an instructive role of the chromatin environment, differential chromatin marks on sex chromosomes and autosomes are associated with the differential response to asynapsis in *C. elegans* (Checchi and Engebrecht 2011; Lamelza and Bhalla 2012).

The chromosome axes likely have multiple functions in the context of the MCN. They form the platform for the binding and phosphorylation of HORMAD proteins, which form a key part of the chromosomal environment for the MCN in many organisms (Xu et al. 1997; Martinez-Perez and Villeneuve 2005; Carballo et al. 2008; Lin et al. 2010; Shin et al. 2010, 2013; Daniel et al. 2011; Kogo et al. 2012a,b; Wojtasz et al. 2012; Cheng et al. 2013). In addition, chromosome axis components also interact directly with components of MCN. For example, the *S. cerevisiae* axis protein Red1 physically associates with the 9-1-1 complex, an interaction required for MCN activity (Eichinger and Jentsch 2010). Cohesin is similarly required for the recruitment of the 9-1-1 complex in *C. elegans* (Lightfoot et al. 2011). Furthermore, because the activation of CHK2 kinases by ATM/ATR typically requires the presence of adaptor proteins, it has been suggested that chromosome axis proteins may provide such an adaptor function for the activation of the MCN (Niu et al. 2005; Carballo et al. 2008; Hunter 2008; Eichinger and Jentsch 2010; Tougan et al. 2010).

There is also increasing evidence that the sequential dynamic changes of meiotic chromosome axes play a significant role in the activation and modulation of the MCN. Most of this evidence stems from the functional analysis

of TRIP13^{Pch2}, a widely conserved AAA⁺-ATPase. TRIP13^{Pch2} modulates meiotic chromosome structure in a variety of contexts, in many cases by controlling the chromosomal depletion or phosphorylation of HORMAD proteins (San-Segundo and Roeder 1999; Borner et al. 2008; Wojtasz et al. 2009; Roig et al. 2010; Vader et al. 2011; Miao et al. 2013; Chen et al. 2014; Lo et al. 2014). Mutants lacking TRIP13^{Pch2} share a number of phenotypic features with mutants lacking ATM or ATR, consistent with the model that TRIP13^{Pch2} is required for full activation of the MCN (San-Segundo and Roeder 1999; Borner et al. 2008; Joshi et al. 2009; Joyce and McKim 2009, 2010; Wojtasz et al. 2009; Zanders and Alani 2009; Roig et al. 2010; Zanders et al. 2011; Farmer et al. 2012). These effects are likely to a large extent a secondary consequence of disrupting HORMAD function, although in *S. cerevisiae*, TRIP13^{Pch2} also directly modulates ATM^{Tel1} by interacting with the MRN^{Xrs2} complex (Ho and Burgess 2011).

Signal Generation

Chromosomal architecture may also lie at the root of one of the more perplexing aspects of meiotic checkpoint regulation, the ability of the MCN to respond to synapsis defects independently of SPO11-induced DSBs (Barchi et al. 2005; Bellani et al. 2005; Bhalla and Dernburg 2005; Barbosa et al. 2007; Joyce and McKim 2009; Lu et al. 2010). Available evidence suggests that features of the meiosis-specific chromosome structure itself may be able to activate the MCN. In most cases, checkpoint activity in response to asynapsis requires the activity of TRIP13^{Pch2} on HORMAD proteins (Bhalla and Dernburg 2005; Joyce and McKim 2009, 2010; Kogo et al. 2012a; Wojtasz et al. 2012), and in several cases it also requires the activity of the histone deacetylase Sir2 (San-Segundo and Roeder 1999; Joyce and McKim 2010; Pek et al. 2012).

What DSB-independent feature of chromosome structure ultimately initiates the MCN signal remains unclear. The small ubiquitin-like protein modifier SUMO may be involved in signal formation, as it is one of the earliest

marks distinguishing the unsynapsed sex chromosomes for MSCI (Vigodner 2009), and is involved in MCN activation in *S. cerevisiae* (Eichinger and Jentsch 2010). However, SUMO accumulation on sex chromosomes depends on ATR (Royo et al. 2013), indicating that another aspect of asynapsis serves as a signal in this case. It is possible that the presence of HORMADs on unsynapsed chromosomes is itself the signal activating the MCN. In mouse, HORMAD1 has multiple roles in meiotic prophase (Shin et al. 2010, 2013; Daniel et al. 2011), whereas HORMAD2 is selectively required for SPO11-independent spreading of γ -H2AFX and MSUC/MSCI (Wojtasz et al. 2012). Because HORMAD2 directly binds to HORMAD1 (Wojtasz et al. 2012), an intriguing possibility is that HORMAD1/2 colocalization creates a SPO11-independent signal to activate the MCN. A direct checkpoint-activating function of HORMADs is also suggested by the observation that *C. elegans* mutants precociously expressing the HORMAD protein HTP-3 arrest at meiotic entry in an ATM^{ATL-1}-dependent manner without apparent DNA damage (Burger et al. 2013). Alternatively, there may be a SPO11-independent source of DSB formation and MCN activation, as indicated by the recent observation of SPO11-independent DNA repair foci on unsynapsed chromosomal regions of mouse meiocytes as well as SPO11-independent crossovers in *Coprinus cinereus* (Carofiglio et al. 2013; Crown et al. 2013).

Response Modulation

Another interesting feature of the MCN is the relative insensitivity to damage that is observed in several branches of this network. For example, whereas mitotic *S. cerevisiae* cells arrest in response to a single induced DSB (Lee et al. 2000), the same lesion elicits no comparable response in meiotic prophase (Malkova et al. 1996). A similar insensitivity to damage is also observed in mouse oocytes (Marangos and Carroll 2012). Intriguingly, the canonical *S. cerevisiae* CHK2^{Rad53} checkpoint kinase, which would launch the arrest response in mitotic cells, is prevented from accessing meiotic chro-

mosomes in most circumstances (Cartagena-Lirola et al. 2008) and is maintained in an inactive state by protein phosphatase 4 (Falk et al. 2010). Indeed, overexpression of the CHK2^{Rad53} kinase delays meiotic progression (Usui and Kanehara 2013). Conversely, there also exist titration effects, whereby too many aberrant structures impair the normal checkpoint response. In particular, the MSUC response in mouse breaks down in the presence of too many unsynapsed chromosomes (Mahadevaiah et al. 2008; Kouznetsova et al. 2009). This may indicate a signaling limit for the MSUC response, and may also provide a safeguard against initiating MSUC in the early stages of meiotic prophase when most chromosomes are unsynapsed.

Signal Integration and Separation

The complexity of the meiotic checkpoint network raises the question how signals are integrated or separated. ATM and ATR achieve signal integration simply by phosphorylating many of the same target sites. Thus, the presence of blunt DNA ends and ssDNA can both elicit a delay in meiotic progression (Hochwagen and Amon 2006; Wu and Burgess 2006), and both kinases can stimulate DNA resection and regulate repair partner choice (Cartagena-Lirola et al. 2006; Carballo et al. 2008).

In other cases, it likely is necessary to avoid cross talk between signals. For example, ATR^{Mec1} regulates DSB formation differently in response to stalled replication during premeiotic S phase compared with leptotema when DSB formation has initiated (Argunhan et al. 2013; Blitzblau and Hochwagen 2013; Carballo et al. 2013; Cheng et al. 2013; Gray et al. 2013). Similarly, axis proteins (e.g., HORMAD1) assemble onto chromosomes concurrently with DNA replication but only become ATR substrates on DSB formation (Carballo et al. 2008; Wojtasz et al. 2009; Blitzblau et al. 2012; Cheng et al. 2013). One way to achieve signal separation is through alternative signaling complexes, such as the use of different CHK2 kinases (Blitzblau and Hochwagen 2013), or different signaling platforms, as has recently been suggested for the alternative

V.V. Subramanian and A. Hochwagen

9-1-1 complexes active during mouse meiotic prophase (Lyndaker et al. 2013a,b; Vasileva et al. 2013). The decision for which signaling complex will ultimately be activated is likely driven by the distinct spectrum of MCN interactions that are possible at replication forks, DSBs, or unsynapsed regions, and will therefore again be strongly context dependent.

Finally, there is now substantial evidence that the signaling environment of the MCN evolves over the course of meiotic prophase. In *S. cerevisiae* and mouse there are clear differences in the timing by which specific MCN-mediated phosphorylation events appear and disappear (Barchi et al. 2005; Bellani et al. 2005; Fukuda et al. 2012; Cheng et al. 2013). Of particular interest here is the entry into pachynema, which is associated with an apparent switch in the signaling and response properties of the MCN. For example, the phosphorylated forms of HORMADs and CHK2^{Mek1} specifically disappear in pachynema (Cartagena-Lirola et al. 2008; Fukuda et al. 2012; Cheng et al. 2013). In *C. elegans*, exogenous DSBs can trigger nuclear reorganization and persistent SUN-1 phosphorylation in leptonema/zygonema but not in pachynema (Woglar and Jantsch 2013). Moreover, repair pathway choice for exogenous DSBs also changes at later stages in meiotic prophase (Rosu et al. 2011; Libuda et al. 2013). Temporal evolution of MCN activity may in some cases be the result of stage-specific activation of phosphatases that remove MCN-dependent signals (Bailis et al. 2000; Hochwagen and Amon 2006; Falk et al. 2010; Cheng et al. 2013). In addition, in the spatially structured gonads of metazoans, temporal differentiation of the checkpoint response can also be imparted by external signals. For example, in *C. elegans* hermaphrodites, checkpoint-induced apoptosis is restricted by Ras/MAP kinase signaling to late pachynema, perhaps to avoid inappropriate cell death induction at earlier stages when SPO11-induced DSBs are prevalent (Rutkowski et al. 2011). The MCN therefore integrates both spatial and temporal information to yield a highly context-dependent coordination hub for the step-by-step progression through meiotic prophase.

CONCLUDING REMARKS

Our understanding of the MCN has progressed by leaps and bounds over the past several years, although with new insights, new experimental challenges have emerged. The increasingly evident interconnectedness of the MCN means that signaling outputs must be understood as the summation of signaling branches that modulate and feed back on each other. Dissecting this network will require more precisely regulatable genetic tools, new modeling approaches, as well as a better description of meiotic chromosome structure. Encouragingly, with the accelerating rate of discovery of direct MCN targets in multiple organisms as well as the first applications of systems-level analyses, the study of the MCN has clearly reached a new stage, and a comprehensive understanding of the MCN is starting to be within reach. Ultimately, of course, a major goal of this research is to use the emerging knowledge of the MCN for a better understanding of human chromosome inheritance and fertility. Because the MCN modulates the timing and activity of meiotic processes, partial loss-of-function mutations of MCN components are expected to have a major impact on gamete quality. The increasing abundance of whole-genome patient data holds big promise in this respect. In the coming years, informed by the research conducted in model organisms, we expect that these data will provide major insights into the high incidence of spontaneous abortions and chromosomal birth defects in humans.

ACKNOWLEDGMENTS

This work is supported by National Institutes of Health Grant GM088248 to A.H.

REFERENCES

- Abdu U, Brodsky M, Schupbach T. 2002. Activation of a meiotic checkpoint during *Drosophila* oogenesis regulates the translation of Gurken through Chk2/Mnk. *Curr Biol* 12: 1645–1651.
- Acosta I, Ontoso D, San-Segundo PA. 2011. The budding yeast polo-like kinase Cdc5 regulates the Ndt80 branch of the meiotic recombination checkpoint pathway. *Mol Biol Cell* 22: 3478–3490.

- Ahmed EA, Sfeir A, Takai H, Scherthan H. 2013. Ku70 and non-homologous end joining protect testicular cells from DNA damage. *J Cell Sci* **126**: 3095–3104.
- Allers T, Lichten M. 2001. Differential timing and control of noncrossover and crossover recombination during meiosis. *Cell* **106**: 47–57.
- Anderson E, Burns C, Zolan ME. 2012. Global gene expression in *Coprinopsis cinerea* meiotic mutants reflects checkpoint arrest. *G3 (Bethesda)* **2**: 1213–1221.
- Argunhan B, Farmer S, Leung WK, Terentyev Y, Humphries N, Tsubouchi T, Toyozumi H, Tsubouchi H. 2013. Direct and indirect control of the initiation of meiotic recombination by DNA damage checkpoint mechanisms in budding yeast. *PLoS ONE* **8**: e65875.
- Baarends WM, Grootegoed JA. 2003. Chromatin dynamics in the male meiotic prophase. *Cytogenet Genome Res* **103**: 225–234.
- Bailis JM, Smith AV, Roeder GS. 2000. Bypass of a meiotic checkpoint by overproduction of meiotic chromosomal proteins. *Mol Cell Biol* **20**: 4838–4848.
- Barbosa V, Kimm N, Lehmann R. 2007. A maternal screen for genes regulating *Drosophila* oocyte polarity uncovers new steps in meiotic progression. *Genetics* **176**: 1967–1977.
- Barchi M, Mahadevaiah S, Di Giacomo M, Baudat F, de Rooij DG, Burgoyne PS, Jasin M, Keeney S. 2005. Surveillance of different recombination defects in mouse spermatocytes yields distinct responses despite elimination at an identical developmental stage. *Mol Cell Biol* **25**: 7203–7215.
- Barchi M, Roig I, Di Giacomo M, de Rooij DG, Keeney S, Jasin M. 2008. ATM promotes the obligate XY crossover and both crossover control and chromosome axis integrity on autosomes. *PLoS Genet* **4**: e1000076.
- Bartrand AJ, Iyasu D, Marincio SM, Brush GS. 2006. Evidence of meiotic crossover control in *Saccharomyces cerevisiae* through Mec1-mediated phosphorylation of replication protein A. *Genetics* **172**: 27–39.
- Bean CJ, Schaner CE, Kelly WG. 2004. Meiotic pairing and imprinted X chromatin assembly in *Caenorhabditis elegans*. *Nat Genet* **36**: 100–105.
- Bellani MA, Romanienko PJ, Cairatti DA, Camerini-Otero RD. 2005. SPO11 is required for sex-body formation, and Spo11 heterozygosity rescues the prophase arrest of *Atm*^{-/-} spermatocytes. *J Cell Sci* **118**: 3233–3245.
- Berchowitz LE, Copenhaver GP. 2010. Genetic interference: Don't stand so close to me. *Curr Genomics* **11**: 91–102.
- Bhalla N, Dernburg AF. 2005. A conserved checkpoint monitors meiotic chromosome synapsis in *Caenorhabditis elegans*. *Science* **310**: 1683–1686.
- Bishop DK, Zickler D. 2004. Early decision; meiotic crossover interference prior to stable strand exchange and synapsis. *Cell* **117**: 9–15.
- Blanco-Rodriguez J. 2012. Programmed phosphorylation of histone H2AX precedes a phase of DNA double-strand break-independent synapsis in mouse meiosis. *Reproduction* **144**: 699–712.
- Blat Y, Protacio RU, Hunter N, Kleckner N. 2002. Physical and functional interactions among basic chromosome organizational features govern early steps of meiotic chromosome formation. *Cell* **111**: 791–802.
- Blitzblau HG, Hochwagen A. 2013. ATR/Mec1 prevents lethal meiotic recombination initiation on partially replicated chromosomes in budding yeast. *eLife* **2**: e00844.
- Blitzblau HG, Chan CS, Hochwagen A, Bell SP. 2012. Separation of DNA replication from the assembly of break-competent meiotic chromosomes. *PLoS Genet* **8**: e1002643.
- Bolcun-Filas E, Rinaldi VD, White ME, Schimenti JC. 2014. Reversal of female infertility by *Chk2* ablation reveals the oocyte DNA damage checkpoint pathway. *Science* **343**: 533–536.
- Borde V, de Massy B. 2013. Programmed induction of DNA double strand breaks during meiosis: Setting up communication between DNA and the chromosome structure. *Curr Opin Genet Dev* **23**: 147–155.
- Borner GV, Barot A, Kleckner N. 2008. Yeast Pch2 promotes domainal axis organization, timely recombination progression, and arrest of defective recombinosomes during meiosis. *Proc Natl Acad Sci* **105**: 3327–3332.
- Branzei D, Foiani M. 2010. Maintaining genome stability at the replication fork. *Nat Rev Mol Cell Biol* **11**: 208–219.
- Burger J, Merlet J, Tavernier N, Richaudeau B, Arnold A, Ciosk R, Bowerman B, Pintard L. 2013. CRL2^{LRR-1} E3-ligase regulates proliferation and progression through meiosis in the *Caenorhabditis elegans* germline. *PLoS Genet* **9**: e1003375.
- Burgoyne PS, Mahadevaiah SK, Turner JM. 2009. The consequences of asynapsis for mammalian meiosis. *Nat Rev Genet* **10**: 207–216.
- Callender TL, Hollingsworth NM. 2010. Mek1 suppression of meiotic double-strand break repair is specific to sister chromatids, chromosome autonomous and independent of Rec8 cohesin complexes. *Genetics* **185**: 771–782.
- Carballo JA, Cha RS. 2007. Meiotic roles of Mec1, a budding yeast homolog of mammalian ATR/ATM. *Chromosome Res* **15**: 539–550.
- Carballo JA, Johnson AL, Sedgwick SG, Cha RS. 2008. Phosphorylation of the axial element protein Hop1 by Mec1/Tel1 ensures meiotic interhomolog recombination. *Cell* **132**: 758–770.
- Carballo JA, Panizza S, Serrentino ME, Johnson AL, Geymonat M, Borde V, Klein F, Cha RS. 2013. Budding yeast ATM/ATR control meiotic double-strand break (DSB) levels by down-regulating Rec114, an essential component of the DSB-machinery. *PLoS Genet* **9**: e1003545.
- Carlton PM, Farruggio AP, Dernburg AF. 2006. A link between meiotic prophase progression and crossover control. *PLoS Genet* **2**: e12.
- Carofiglio F, Inagaki A, de Vries S, Wassenaar E, Schoenmakers S, Vermeulen C, van Cappellen WA, Sleddens-Linkels E, Grootegoed JA, Te Riele HP, et al. 2013. SPO11-independent DNA repair foci and their role in meiotic silencing. *PLoS Genet* **9**: e1003538.
- Carpenter AT, Sandler L. 1974. On recombination-defective meiotic mutants in *Drosophila melanogaster*. *Genetics* **76**: 453–475.
- Cartagena-Lirola H, Guerini I, Viscardi V, Lucchini G, Longhese MP. 2006. Budding yeast Sae2 is an in vivo target of the Mec1 and Tel1 checkpoint kinases during meiosis. *Cell Cycle* **5**: 1549–1559.

V.V. Subramanian and A. Hochwagen

- Cartagena-Lirola H, Guerini I, Manfrini N, Lucchini G, Longhese MP. 2008. Role of the *Saccharomyces cerevisiae* Rad53 checkpoint kinase in signaling double-strand breaks during the meiotic cell cycle. *Mol Cell Biol* **28**: 4480–4493.
- Checchi PM, Engebrecht J. 2011. *Caenorhabditis elegans* histone methyltransferase MET-2 shields the male X chromosome from checkpoint machinery and mediates meiotic sex chromosome inactivation. *PLoS Genet* **7**: e1002267.
- Chen C, Jomaa A, Ortega J, Alani EE. 2014. Pch2 is a hexameric ring ATPase that remodels the chromosome axis protein Hop1. *Proc Natl Acad Sci* **111**: E44–E53.
- Cheng YH, Chuang CN, Shen HJ, Lin FM, Wang TF. 2013. Three distinct modes of Mec1/ATR and Tel1/ATM activation illustrate differential checkpoint targeting during budding yeast early meiosis. *Mol Cell Biol* **33**: 3365–3376.
- Chu S, Herskowitz I. 1998. Gametogenesis in yeast is regulated by a transcriptional cascade dependent on Ndt80. *Mol Cell* **1**: 685–696.
- Chuang CN, Cheng YH, Wang TF. 2012. Mek1 stabilizes Hop1-Thr318 phosphorylation to promote interhomolog recombination and checkpoint responses during yeast meiosis. *Nucleic Acids Res* **40**: 11416–11427.
- Clerici M, Trovesi C, Galbiati A, Lucchini G, Longhese MP. 2014. Mec1/ATR regulates the generation of single-stranded DNA that attenuates Tel1/ATM signaling at DNA ends. *EMBO J* **33**: 198–216.
- Colaiacono MP. 2006. The many facets of SC function during *C. elegans* meiosis. *Chromosoma* **115**: 195–211.
- Crown KN, Savvitsky OP, Malik SB, Logsdon J, Williams RS, Tainer JA, Zolan ME. 2013. A mutation in the FHA domain of *Coprinus cinereus* Nbs1 leads to Spo11-independent meiotic recombination and chromosome segregation. *G3 (Bethesda)* **3**: 1927–1943.
- Culligan KM, Britt AB. 2008. Both ATM and ATR promote the efficient and accurate processing of programmed meiotic double-strand breaks. *Plant J* **55**: 629–638.
- Daniel K, Lange J, Hached K, Fu J, Anastasiadis K, Roig I, Cooke HJ, Stewart AE, Wassmann K, Jasin M, et al. 2011. Meiotic homologue alignment and its quality surveillance are controlled by mouse HORMAD1. *Nat Cell Biol* **13**: 599–610.
- Derry WB, Putzke AP, Rothman JH. 2001. *Caenorhabditis elegans* p53: Role in apoptosis, meiosis, and stress resistance. *Science* **294**: 591–595.
- Di Giacomo M, Barchi M, Baudat F, Edelmann W, Keeney S, Jasin M. 2005. Distinct DNA-damage-dependent and -independent responses drive the loss of oocytes in recombination-defective mouse mutants. *Proc Natl Acad Sci* **102**: 737–742.
- Ding DQ, Chikashige Y, Haraguchi T, Hiraoka Y. 1998. Oscillatory nuclear movement in fission yeast meiotic prophase is driven by astral microtubules, as revealed by continuous observation of chromosomes and microtubules in living cells. *J Cell Sci* **111**: 701–712.
- Eichinger CS, Jentsch S. 2010. Synaptonemal complex formation and meiotic checkpoint signaling are linked to the lateral element protein Red1. *Proc Natl Acad Sci* **107**: 11370–11375.
- Falk JE, Chan AC, Hoffmann E, Hochwagen A. 2010. A Mec1- and PP4-dependent checkpoint couples centromere pairing to meiotic recombination. *Dev Cell* **19**: 599–611.
- Farmer S, Hong EJ, Leung WK, Argunhan B, Terentyev Y, Humphries N, Toyozumi H, Tsubouchi H. 2012. Budding yeast Pch2, a widely conserved meiotic protein, is involved in the initiation of meiotic recombination. *PLoS ONE* **7**: e39724.
- Fernandez-Capetillo O, Liebe B, Scherthan H, Nussenzweig A. 2003. H2AX regulates meiotic telomere clustering. *J Cell Biol* **163**: 15–20.
- Ferrari M, Nachimuthu BT, Donnianni RA, Klein H, Pelligioli A. 2013. Tid1/Rdh54 translocase is phosphorylated through a Mec1- and Rad53-dependent manner in the presence of DSB lesions in budding yeast. *DNA Repair* **12**: 347–355.
- Fraune J, Schramm S, Alsheimer M, Benavente R. 2012. The mammalian synaptonemal complex: Protein components, assembly and role in meiotic recombination. *Exp Cell Res* **318**: 1340–1346.
- Fukuda T, Pratto F, Schimenti JC, Turner JM, Camerini-Otero RD, Hoog C. 2012. Phosphorylation of chromosome core components may serve as axis marks for the status of chromosomal events during mammalian meiosis. *PLoS Genet* **8**: e1002485.
- Garcia V, Phelps SE, Gray S, Neale MJ. 2011. Bidirectional resection of DNA double-strand breaks by Mre11 and Exo1. *Nature* **479**: 241–244.
- Garcia-Cruz R, Roig I, Robles P, Scherthan H, Garcia Caldes M. 2009. ATR, BRCA1 and γ H2AX localize to unsynapsed chromosomes at the pachytene stage in human oocytes. *Reprod Biomed Online* **18**: 37–44.
- Gartner A, Milstein S, Ahmed S, Hodgkin J, Hengartner MO. 2000. A conserved checkpoint pathway mediates DNA damage-induced apoptosis and cell cycle arrest in *C. elegans*. *Mol Cell* **5**: 435–443.
- Gerton JL, Hawley RS. 2005. Homologous chromosome interactions in meiosis: Diversity amidst conservation. *Nat Rev Genet* **6**: 477–487.
- Ghabrial A, Schupbach T. 1999. Activation of a meiotic checkpoint regulates translation of Gurken during *Drosophila* oogenesis. *Nat Cell Biol* **1**: 354–357.
- Ghabrial A, Ray RP, Schupbach T. 1998. *okra* and *spindle-B* encode components of the RAD52 DNA repair pathway and affect meiosis and patterning in *Drosophila* oogenesis. *Genes Dev* **12**: 2711–2723.
- Govin J, Dorsey J, Gaucher J, Rousseaux S, Khochbin S, Berger SL. 2010. Systematic screen reveals new functional dynamics of histones H3 and H4 during gametogenesis. *Genes Dev* **24**: 1772–1786.
- Gray S, Allison RM, Garcia V, Goldman AS, Neale MJ. 2013. Positive regulation of meiotic DNA double-strand break formation by activation of the DNA damage checkpoint kinase Mec1(ATR). *Open Biol* **3**: 130019.
- Grushcow JM, Holzen TM, Park KJ, Weinert T, Lichten M, Bishop DK. 1999. *Saccharomyces cerevisiae* checkpoint genes *MEC1*, *RAD17* and *RAD24* are required for normal meiotic recombination partner choice. *Genetics* **153**: 607–620.

- Guo H, King MC. 2013. A quality control mechanism linking meiotic success to release of ascospores. *PLoS ONE* **8**: e82758.
- Harrison JC, Haber JE. 2006. Surviving the breakup: The DNA damage checkpoint. *Annu Rev Genet* **40**: 209–235.
- Hartwell LH, Weinert TA. 1989. Checkpoints: Controls that ensure the order of cell cycle events. *Science* **246**: 629–634.
- Henderson KA, Keeney S. 2004. Tying synaptonemal complex initiation to the formation and programmed repair of DNA double-strand breaks. *Proc Natl Acad Sci* **101**: 4519–4524.
- Hepworth SR, Friesen H, Segall J. 1998. *NDT80* and the meiotic recombination checkpoint regulate expression of middle sporulation-specific genes in *Saccharomyces cerevisiae*. *Mol Cell Biol* **18**: 5750–5761.
- Ho HC, Burgess SM. 2011. Pch2 acts through Xrs2 and Tel1/ATM to modulate interhomolog bias and checkpoint function during meiosis. *PLoS Genet* **7**: e1002351.
- Hochwagen A, Amon A. 2006. Checking your breaks: Surveillance mechanisms of meiotic recombination. *Curr Biol* **16**: R217–R228.
- Hochwagen A, Tham WH, Brar GA, Amon A. 2005. The FK506 binding protein Fpr3 counteracts protein phosphatase 1 to maintain meiotic recombination checkpoint activity. *Cell* **122**: 861–873.
- Hollingsworth NM. 2010. Phosphorylation and the creation of interhomolog bias during meiosis in yeast. *Cell Cycle* **9**: 436–437.
- Hong S, Sung Y, Yu M, Lee M, Kleckner N, Kim KP. 2013. The logic and mechanism of homologous recombination partner choice. *Mol Cell* **51**: 440–453.
- Hunter N. 2008. Hop1 and the meiotic DNA-damage response. *Cell* **132**: 731–732.
- Hunter N, Kleckner N. 2001. The single-end invasion: An asymmetric intermediate at the double-strand break to double-Holliday junction transition of meiotic recombination. *Cell* **106**: 59–70.
- Iacovella MG, Daly CN, Kelly JS, Michielsen AJ, Clyne RK. 2010. Analysis of Polo-like kinase Cdc5 in the meiosis recombination checkpoint. *Cell Cycle* **9**: 1182–1193.
- Ichijima Y, Ichijima M, Lou Z, Nussenzweig A, Camerini-Otero RD, Chen J, Andreassen PR, Namekawa SH. 2011. MDC1 directs chromosome-wide silencing of the sex chromosomes in male germ cells. *Genes Dev* **25**: 959–971.
- Ivanovska I, Khandan T, Ito T, Orr-Weaver TL. 2005. A histone code in meiosis: The histone kinase, NHK-1, is required for proper chromosomal architecture in *Drosophila* oocytes. *Genes Dev* **19**: 2571–2582.
- Jaramillo-Lambert A, Harigaya Y, Vitt J, Villeneuve A, Engebrecht J. 2010. Meiotic errors activate checkpoints that improve gamete quality without triggering apoptosis in male germ cells. *Curr Biol* **20**: 2078–2089.
- Johnson R, Borde V, Neale MJ, Bishop-Bailey A, North M, Harris S, Nicolas A, Goldman AS. 2007. Excess single-stranded DNA inhibits meiotic double-strand break repair. *PLoS Genet* **3**: e223.
- Joshi N, Barot A, Jamison C, Borner GV. 2009. Pch2 links chromosome axis remodeling at future crossover sites and crossover distribution during yeast meiosis. *PLoS Genet* **5**: e1000557.
- Joyce EF, McKim KS. 2009. *Drosophila* PCH2 is required for a pachytene checkpoint that monitors double-strand-break-independent events leading to meiotic crossover formation. *Genetics* **181**: 39–51.
- Joyce EF, McKim KS. 2010. Chromosome axis defects induce a checkpoint-mediated delay and interchromosomal effect on crossing over during *Drosophila* meiosis. *PLoS Genet* **6**.
- Joyce EF, Pedersen M, Tiong S, White-Brown SK, Paul A, Campbell SD, McKim KS. 2011. *Drosophila* ATM and ATR have distinct activities in the regulation of meiotic DNA damage and repair. *J Cell Biol* **195**: 359–367.
- Joyce EF, Paul A, Chen KE, Tanneti N, McKim KS. 2012. Multiple barriers to nonhomologous DNA end joining during meiosis in *Drosophila*. *Genetics* **191**: 739–746.
- Kauppi L, Barchi M, Lange J, Baudat F, Jasin M, Keeney S. 2013a. Numerical constraints and feedback control of double-strand breaks in mouse meiosis. *Genes Dev* **27**: 873–886.
- Kauppi L, Jasin M, Keeney S. 2013b. How much is enough? Control of DNA double-strand break numbers in mouse meiosis. *Cell Cycle* **12**: 2719–2720.
- Keeney S. 2001. Mechanism and control of meiotic recombination initiation. *Curr Top Dev Biol* **52**: 1–53.
- Kim DA, Suh EK. 2014. Defying DNA double strand break-induced death during prophase I meiosis by temporal TAp63 α phosphorylation regulation in developing mouse oocytes. *Mol Cell Biol*.
- Kim KP, Weiner BM, Zhang L, Jordan A, Dekker J, Kleckner N. 2010. Sister cohesion and structural axis components mediate homolog bias of meiotic recombination. *Cell* **143**: 924–937.
- Kleckner N. 1996. Meiosis: How could it work? *Proc Natl Acad Sci* **93**: 8167–8174.
- Klein F, Mahr P, Galova M, Buonomo SB, Michaelis C, Nairz K, Nasmyth K. 1999. A central role for cohesins in sister chromatid cohesion, formation of axial elements, and recombination during yeast meiosis. *Cell* **98**: 91–103.
- Kogo H, Tsutsumi M, Inagaki H, Ohye T, Kiyonari H, Kurahashi H. 2012a. HORMAD2 is essential for synapsis surveillance during meiotic prophase via the recruitment of ATR activity. *Genes Cells* **17**: 897–912.
- Kogo H, Tsutsumi M, Ohye T, Inagaki H, Abe T, Kurahashi H. 2012b. HORMAD1-dependent checkpoint/surveillance mechanism eliminates asynaptic oocytes. *Genes Cells* **17**: 439–454.
- Koszul R, Kleckner N. 2009. Dynamic chromosome movements during meiosis: Away to eliminate unwanted connections? *Trends Cell Biol* **19**: 716–724.
- Koszul R, Kameoka S, Weiner BM. 2009. Real-time imaging of meiotic chromosomes in *Saccharomyces cerevisiae*. *Methods Mol Biol* **558**: 81–89.
- Kouznetsova A, Wang H, Bellani M, Camerini-Otero RD, Jessberger R, Hoog C. 2009. BRCA1-mediated chromatin silencing is limited to oocytes with a small number of asynapsed chromosomes. *J Cell Sci* **122**: 2446–2452.
- Kurzbaue MT, Uanschou C, Chen D, Schlegelhofer P. 2012. The recombinases DMC1 and RAD51 are functionally

V.V. Subramanian and A. Hochwagen

- and spatially separated during meiosis in *Arabidopsis*. *Plant Cell* **24**: 2058–2070.
- Labella S, Woglar A, Jantsch V, Zetka M. 2011. Polo kinases establish links between meiotic chromosomes and cytoskeletal forces essential for homolog pairing. *Dev Cell* **21**: 948–958.
- Lamelza P, Bhalla N. 2012. Histone methyltransferases MES-4 and MET-1 promote meiotic checkpoint activation in *Caenorhabditis elegans*. *PLoS Genet* **8**: e1003089.
- Lancaster OM, Cullen CF, Ohkura H. 2007. NHK-1 phosphorylates BAF to allow karyosome formation in the *Drosophila* oocyte nucleus. *J Cell Biol* **179**: 817–824.
- Lancaster OM, Breuer M, Cullen CF, Ito T, Ohkura H. 2010. The meiotic recombination checkpoint suppresses NHK-1 kinase to prevent reorganization of the oocyte nucleus in *Drosophila*. *PLoS Genet* **6**: e1001179.
- Lange J, Pan J, Cole F, Thelen MP, Jasini M, Keeney S. 2011. ATM controls meiotic double-strand-break formation. *Nature* **479**: 237–240.
- Lao JP, Hunter N. 2010. Trying to avoid your sister. *PLoS Biol* **8**: e1000519.
- Lao JP, Cloud V, Huang CC, Grubb J, Thacker D, Lee CY, Dresser ME, Hunter N, Bishop DK. 2013. Meiotic cross-over control by concerted action of Rad51-Dmc1 in homolog template bias and robust homeostatic regulation. *PLoS Genet* **9**: e1003978.
- Latypov V, Rothenberg M, Lorenz A, Octobre G, Csutak O, Lehmann E, Loidl J, Kohli J. 2010. Roles of Hop1 and Mek1 in meiotic chromosome pairing and recombination partner choice in *Schizosaccharomyces pombe*. *Mol Cell Biol* **30**: 1570–1581.
- Lee JY, Orr-Weaver TL. 2001. The molecular basis of sister-chromatid cohesion. *Annu Rev Cell Dev Biol* **17**: 753–777.
- Lee SE, Pelliccioli A, Demeter J, Vaze MP, Gasch AP, Malkova A, Brown PO, Botstein D, Stearns T, Foiani M, et al. 2000. Arrest, adaptation, and recovery following a chromosome double-strand break in *Saccharomyces cerevisiae*. *Cold Spring Harb Symp Quant Biol* **65**: 303–314.
- Lee CY, Conrad MN, Dresser ME. 2012. Meiotic chromosome pairing is promoted by telomere-led chromosome movements independent of bouquet formation. *PLoS Genet* **8**: e1002730.
- Libuda DE, Uzawa S, Meyer BJ, Villeneuve AM. 2013. Meiotic chromosome structures constrain and respond to designation of crossover sites. *Nature* **502**: 703–706.
- Liebe B, Petukhova G, Barchi M, Bellani M, Braselmann H, Nakano T, Pandita TK, Jasini M, Fornace A, Meistrich ML, et al. 2006. Mutations that affect meiosis in male mice influence the dynamics of the mid-preleptotene and bouquet stages. *Exp Cell Res* **312**: 3768–3781.
- Lightfoot J, Testori S, Barroso C, Martinez-Perez E. 2011. Loading of meiotic cohesin by SCC-2 is required for early processing of DSBs and for the DNA damage checkpoint. *Curr Biol* **21**: 1421–1430.
- Lin FM, Lai YJ, Shen HJ, Cheng YH, Wang TF. 2010. Yeast axial-element protein, Red1, binds SUMO chains to promote meiotic interhomologue recombination and chromosome synapsis. *EMBO J* **29**: 586–596.
- Liu Y, Gaines WA, Callender T, Busygina V, Oke A, Sung P, Fung JC, Hollingsworth NM. 2014. Down-regulation of Rad51 activity during meiosis in yeast prevents competition with Dmc1 for repair of double-strand breaks. *PLoS Genet* **10**: e1004005.
- Lo YH, Chuang CN, Wang TF. 2014. Pch2 prevents Mec1/Tel1-mediated Hop1 phosphorylation occurring independently of Red1 in budding yeast meiosis. *PLoS ONE* **9**: e85687.
- Loidl J, Mochizuki K. 2009. *Tetrahymena* meiotic nuclear reorganization is induced by a checkpoint kinase-dependent response to DNA damage. *Mol Biol Cell* **20**: 2428–2437.
- Loidl J, Lukaszewicz A, Howard-Till RA, Koestler T. 2012. The *Tetrahymena* meiotic chromosome bouquet is organized by centromeres and promotes interhomolog recombination. *J Cell Sci* **125**: 5873–5880.
- Longhese MP, Bonetti D, Guerini I, Manfrini N, Clerici M. 2009. DNA double-strand breaks in meiosis: Checking their formation, processing and repair. *DNA Repair* **8**: 1127–1138.
- Lovejoy CA, Cortez D. 2009. Common mechanisms of PIKK regulation. *DNA Repair* **8**: 1004–1008.
- Lu WJ, Chappo J, Roig I, Abrams JM. 2010. Meiotic recombination provokes functional activation of the p53 regulatory network. *Science* **328**: 1278–1281.
- Luo M, Yang F, Leu NA, Landaiche J, Handel MA, Benavente R, La Salle S, Wang PJ. 2013. MEIOB exhibits single-stranded DNA-binding and exonuclease activities and is essential for meiotic recombination. *Nat Commun* **4**: 2788.
- Lydall D, Nikolsky Y, Bishop DK, Weinert T. 1996. A meiotic recombination checkpoint controlled by mitotic checkpoint genes. *Nature* **383**: 840–843.
- Lyndaker AM, Lim PX, Mleczko JM, Diggins CE, Holloway JK, Holmes RJ, Kan R, Schlafer DH, Freire R, Cohen PE, et al. 2013a. Conditional inactivation of the DNA damage response gene *Hus1* in mouse testis reveals separable roles for components of the RAD9-RAD1-HUS1 complex in meiotic chromosome maintenance. *PLoS Genet* **9**: e1003320.
- Lyndaker AM, Vasileva A, Wolgemuth DJ, Weiss RS, Lieberman HB. 2013b. Clamping down on mammalian meiosis. *Cell Cycle* **12**: 3135–3145.
- MacQueen AJ, Hochwagen A. 2011. Checkpoint mechanisms: The puppet masters of meiotic prophase. *Trends Cell Biol* **21**: 393–400.
- MacQueen AJ, Roeder GS. 2009. Fpr3 and Zip3 ensure that initiation of meiotic recombination precedes chromosome synapsis in budding yeast. *Curr Biol* **19**: 1519–1526.
- MacQueen AJ, Villeneuve AM. 2001. Nuclear reorganization and homologous chromosome pairing during meiotic prophase require *C. elegans* *chk-2*. *Genes Dev* **15**: 1674–1687.
- Mahadevaiah SK, Turner JM, Baudat F, Rogakou EP, de Boer P, Blanco-Rodriguez J, Jasini M, Keeney S, Bonner WM, Burgoyne PS. 2001. Recombinational DNA double-strand breaks in mice precede synapsis. *Nat Genet* **27**: 271–276.
- Mahadevaiah SK, Bourc'his D, de Rooij DG, Bestor TH, Turner JM, Burgoyne PS. 2008. Extensive meiotic asynapsis in mice antagonizes meiotic silencing of unsyn-

- napsed chromatin and consequently disrupts meiotic sex chromosome inactivation. *J Cell Biol* **182**: 263–276.
- Malkova A, Ross L, Dawson D, Hoekstra MF, Haber JE. 1996. Meiotic recombination initiated by a double-strand break in *rad50Δ* yeast cells otherwise unable to initiate meiotic recombination. *Genetics* **143**: 741–754.
- Manfrini N, Guerini I, Citterio A, Lucchini G, Longhese MP. 2010. Processing of meiotic DNA double strand breaks requires cyclin-dependent kinase and multiple nucleases. *J Biol Chem* **285**: 11628–11637.
- Manterola M, Page J, Vasco C, Berrios S, Parra MT, Viera A, Rufas JS, Zuccotti M, Garagna S, Fernandez-Donoso R. 2009. A high incidence of meiotic silencing of unsynapsed chromatin is not associated with substantial pachytene loss in heterozygous male mice carrying multiple simple robertsonian translocations. *PLoS Genet* **5**: e1000625.
- Marangos P, Carroll J. 2012. Oocytes progress beyond prophase in the presence of DNA damage. *Curr Biol* **22**: 989–994.
- Martinez-Perez E, Villeneuve AM. 2005. HTP-1-dependent constraints coordinate homolog pairing and synapsis and promote chiasma formation during *C. elegans* meiosis. *Genes Dev* **19**: 2727–2743.
- Miao C, Tang D, Zhang H, Wang M, Li Y, Tang S, Yu H, Gu M, Cheng Z. 2013. Central region component1, a novel synaptonemal complex component, is essential for meiotic recombination initiation in rice. *Plant Cell* **25**: 2998–3009.
- Miller MP, Amon A, Unal E. 2013. Meiosis I: When chromosomes undergo extreme makeover. *Curr Opin Cell Biol* **25**: 687–696.
- Mimitou EP, Symington LS. 2009. DNA end resection: Many nucleases make light work. *DNA Repair* **8**: 983–995.
- Miyoshi T, Ito M, Kugou K, Yamada S, Furuichi M, Oda A, Yamada T, Hirota K, Masai H, Ohta K. 2012. A central coupler for recombination initiation linking chromosome architecture to S phase checkpoint. *Mol Cell* **47**: 722–733.
- Nagaoka SI, Hodges CA, Albertini DE, Hunt PA. 2011. Oocyte-specific differences in cell-cycle control create an innate susceptibility to meiotic errors. *Curr Biol* **21**: 651–657.
- Nagaoka SI, Hassold TJ, Hunt PA. 2012. Human aneuploidy: Mechanisms and new insights into an age-old problem. *Nat Rev Genet* **13**: 493–504.
- Nakada D, Matsumoto K, Sugimoto K. 2003. ATM-related Tel1 associates with double-strand breaks through an Xrs2-dependent mechanism. *Genes Dev* **17**: 1957–1962.
- Neale MJ, Keeney S. 2006. Clarifying the mechanics of DNA strand exchange in meiotic recombination. *Nature* **442**: 153–158.
- Niu H, Wan L, Baumgartner B, Schaefer D, Loidl J, Hollingsworth NM. 2005. Partner choice during meiosis is regulated by Hop1-promoted dimerization of Mek1. *Mol Biol Cell* **16**: 5804–5818.
- Niu H, Li X, Job E, Park C, Moazed D, Gygi SP, Hollingsworth NM. 2007. Mek1 kinase is regulated to suppress double-strand break repair between sister chromatids during budding yeast meiosis. *Mol Cell Biol* **27**: 5456–5467.
- Niu H, Wan L, Busygina V, Kwon Y, Allen JA, Li X, Kunz RC, Kubota K, Wang B, Sung P, et al. 2009. Regulation of meiotic recombination via Mek1-mediated Rad54 phosphorylation. *Mol Cell* **36**: 393–404.
- Obeso D, Dawson DS. 2010. Temporal characterization of homology-independent centromere coupling in meiotic prophase. *PLoS ONE* **5**: e10336.
- Ogino K, Masai H. 2006. Rad3-Cds1 mediates coupling of initiation of meiotic recombination with DNA replication. *Mei4*-dependent transcription as a potential target of meiotic checkpoint. *J Biol Chem* **281**: 1338–1344.
- Okaz E, Arguello-Miranda O, Bogdanova A, Vinod PK, Lipp JJ, Markova Z, Zagoriy I, Novak B, Zachariae W. 2012. Meiotic prophase requires proteolysis of M phase regulators mediated by the meiosis-specific APC/C^{Ama1}. *Cell* **151**: 603–618.
- Ontoso D, Acosta I, van Leeuwen F, Freire R, San-Segundo PA. 2013a. Dot1-dependent histone H3K79 methylation promotes activation of the Mek1 meiotic checkpoint effector kinase by regulating the Hop1 adaptor. *PLoS Genet* **9**: e1003262.
- Ontoso D, Kauppi L, Keeney S, San-Segundo PA. 2013b. Dynamics of DOT1L localization and H3K79 methylation during meiotic prophase I in mouse spermatocytes. *Chromosoma* **123**: 147–164.
- Page SL, Hawley RS. 2004. The genetics and molecular biology of the synaptonemal complex. *Annu Rev Cell Dev Biol* **20**: 525–558.
- Panizza S, Mendoza MA, Berlinger M, Huang L, Nicolas A, Shirahige K, Klein F. 2011. Spo11-accessory proteins link double-strand break sites to the chromosome axis in early meiotic recombination. *Cell* **146**: 372–383.
- Pek JW, Ng BF, Kai T. 2012. Polo-mediated phosphorylation of Maelstrom regulates oocyte determination during oogenesis in *Drosophila*. *Development* **139**: 4505–4513.
- Penkner AM, Fridkin A, Gloggnitzer J, Baudrimont A, Machacek T, Woglar A, Csaszar E, Pasierbek P, Ammerer G, Gruenbaum Y, et al. 2009. Meiotic chromosome homology search involves modifications of the nuclear envelope protein Matefin/SUN-1. *Cell* **139**: 920–933.
- Perera D, Perez-Hidalgo L, Moens PB, Reini K, Lakin N, Syvaaja JE, San-Segundo PA, Freire R. 2004. TopBP1 and ATR colocalization at meiotic chromosomes: Role of TopBP1/Cut5 in the meiotic recombination checkpoint. *Mol Biol Cell* **15**: 1568–1579.
- Peretz G, Arie LG, Bakhrat A, Abdu U. 2009. The *Drosophila* *hus1* gene is required for homologous recombination repair during meiosis. *Mech Dev* **126**: 677–686.
- Perez-Hidalgo L, Moreno S, San-Segundo PA. 2008. The fission yeast meiotic checkpoint kinase Mek1 regulates nuclear localization of Cdc25 by phosphorylation. *Cell Cycle* **7**: 3720–3730.
- Petronczki M, Siomos MF, Nasmyth K. 2003. Un ménage à quatre: The molecular biology of chromosome segregation in meiosis. *Cell* **112**: 423–440.
- Phillips CM, Dernburg AE. 2006. A family of zinc-finger proteins is required for chromosome-specific pairing and synapsis during meiosis in *C. elegans*. *Dev Cell* **11**: 817–829.

V.V. Subramanian and A. Hochwagen

- Refolio E, Caverio S, Marcon E, Freire R, San-Segundo PA. 2011. The Ddc2/ATRIP checkpoint protein monitors meiotic recombination intermediates. *J Cell Sci* **124**: 2488–2500.
- Robles P, Roig I, Garcia R, Brieno-Enriquez M, Martin M, Cabero L, Toran N, Garcia Caldes M. 2013. Presence of an extra chromosome alters meiotic double-stranded break repair dynamics and MLH1 foci distribution in human oocytes. *Chromosoma* **122**: 93–102.
- Rockmill B, Lefrançois P, Voelkel-Meiman K, Oke A, Roeder GS, Fung JC. 2013. High throughput sequencing reveals alterations in the recombination signatures with diminishing Spo11 activity. *PLoS Genet* **9**: e1003932.
- Roeder GS, Bailis JM. 2000. The pachytene checkpoint. *Trend Genet* **16**: 395–403.
- Rog O, Dernburg AF. 2013. Chromosome pairing and synapsis during *Caenorhabditis elegans* meiosis. *Curr Opin Cell Biol* **25**: 349–356.
- Roig I, Robles P, Garcia R, Martinez-Flores I, Cabero L, Egozcue J, Liebe B, Scherthan H, Garcia M. 2005. Chromosome 18 pairing behavior in human trisomic oocytes. Presence of an extra chromosome extends bouquet stage. *Reproduction* **129**: 565–575.
- Roig I, Dowdle JA, Toth A, de Rooij DG, Jasin M, Keeney S. 2010. Mouse TRIP13/PCH2 is required for recombination and normal higher-order chromosome structure during meiosis. *PLoS Genet* **6**: e1001062.
- Rosu S, Libuda DE, Villeneuve AM. 2011. Robust crossover assurance and regulated interhomolog access maintain meiotic crossover number. *Science* **334**: 1286–1289.
- Rosu S, Zawadzki KA, Stamper EL, Libuda DE, Reese AL, Dernburg AF, Villeneuve AM. 2013. The *C. elegans* DSB-2 protein reveals a regulatory network that controls competence for meiotic DSB formation and promotes crossover assurance. *PLoS Genet* **9**: e1003674.
- Royo H, Prosser H, Ruzankina Y, Mahadevaiah SK, Cloutier JM, Baumann M, Fukuda T, Hoog C, Toth A, de Rooij DG, et al. 2013. ATR acts stage specifically to regulate multiple aspects of mammalian meiotic silencing. *Genes Dev* **27**: 1484–1494.
- Rutkowski R, Dickinson R, Stewart G, Craig A, Schimpl M, Keyse SM, Gartner A. 2011. Regulation of *Caenorhabditis elegans* p53/CEP-1-dependent germ cell apoptosis by Ras/MAPK signaling. *PLoS Genet* **7**: e1002238.
- San-Segundo PA, Roeder GS. 1999. Pch2 links chromatin silencing to meiotic checkpoint control. *Cell* **97**: 313–324.
- San-Segundo PA, Roeder GS. 2000. Role for the silencing protein Dot1 in meiotic checkpoint control. *Mol Biol Cell* **11**: 3601–3615.
- Schaetzlein S, Chahwan R, Avdievich E, Roa S, Wei K, Eoff RL, Sellers RS, Clark AB, Kunkel TA, Scharff MD, et al. 2013. Mammalian *Exo1* encodes both structural and catalytic functions that play distinct roles in essential biological processes. *Proc Natl Acad Sci* **110**: E2470–E2479.
- Scherthan H. 2001. A bouquet makes ends meet. *Nat Rev Mol Cell Biol* **2**: 621–627.
- Segurado M, Diffley JF. 2008. Separate roles for the DNA damage checkpoint protein kinases in stabilizing DNA replication forks. *Genes Dev* **22**: 1816–1827.
- Serrentino ME, Chaplais E, Sommermeyer V, Borde V. 2013. Differential association of the conserved SUMO ligase Zip3 with meiotic double-strand break sites reveals regional variations in the outcome of meiotic recombination. *PLoS Genet* **9**: e1003416.
- Sheehan MJ, Pawlowski WP. 2009. Live imaging of rapid chromosome movements in meiotic prophase I in maize. *Proc Natl Acad Sci* **106**: 20989–20994.
- Shimada M, Nabeshima K, Tougan T, Nojima H. 2002. The meiotic recombination checkpoint is regulated by checkpoint *rad*⁺ genes in fission yeast. *EMBO J* **21**: 2807–2818.
- Shin YH, Choi Y, Erdin SU, Yatsenko SA, Kloc M, Yang F, Wang PJ, Meistrich ML, Rajkovic A. 2010. *Hormad1* mutation disrupts synaptonemal complex formation, recombination, and chromosome segregation in mammalian meiosis. *PLoS Genet* **6**: e1001190.
- Shin YH, McGuire MM, Rajkovic A. 2013. Mouse HORMAD1 is a meiosis I checkpoint protein that modulates DNA double-strand break repair during female meiosis. *Biol Reprod* **89**: 29.
- Shinohara M, Shinohara A. 2013. Multiple pathways suppress non-allelic homologous recombination during meiosis in *Saccharomyces cerevisiae*. *PLoS ONE* **8**: e63144.
- Shinohara M, Sakai K, Ogawa T, Shinohara A. 2003. The mitotic DNA damage checkpoint proteins Rad17 and Rad24 are required for repair of double-strand breaks during meiosis in yeast. *Genetics* **164**: 855–865.
- Shiu PK, Raju NB, Zickler D, Metzberg RL. 2001. Meiotic silencing by unpaired DNA. *Cell* **107**: 905–916.
- Simchen G. 2009. Commitment to meiosis: What determines the mode of division in budding yeast? *Bioessays* **31**: 169–177.
- Sonntag Brown M, Zanders S, Alani E. 2011. Sustained and rapid chromosome movements are critical for chromosome pairing and meiotic progression in budding yeast. *Genetics* **188**: 21–32.
- Souquet B, Abby E, Herve R, Finsterbusch F, Tourpin S, Le Bouffant R, Duquenne C, Messiaen S, Martini E, Bernardino-Sgherri J, et al. 2013. MEIOB targets single-strand DNA and is necessary for meiotic recombination. *PLoS Genet* **9**: e1003784.
- Sourirajan A, Lichten M. 2008. Polo-like kinase Cdc5 drives exit from pachytene during budding yeast meiosis. *Genes Dev* **22**: 2627–2632.
- Staeva-Vieira E, Yoo S, Lehmann R. 2003. An essential role of *DmRad51 SpnA* in DNA repair and meiotic checkpoint control. *EMBO J* **22**: 5863–5874.
- Stamper EL, Rodenbusch SE, Rosu S, Ahringer J, Villeneuve AM, Dernburg AF. 2013. Identification of DSB-1, a protein required for initiation of meiotic recombination in *Caenorhabditis elegans*, illuminates a crossover assurance checkpoint. *PLoS Genet* **9**: e1003679.
- Stevens D, Oegema K, Desai A. 2013. Meiotic double-strand breaks uncover and protect against mitotic errors in the *C. elegans* Germline. *Curr Biol* **23**: 2400–2406.
- Storlazzi A, Tesse S, Gargano S, James F, Kleckner N, Zickler D. 2003. Meiotic double-strand breaks at the interface of chromosome movement, chromosome remodeling, and reductional division. *Genes Dev* **17**: 2675–2687.
- Subramanian VV, Hochwagen A. 2011. Centromere clustering: Where synapsis begins. *Curr Biol* **21**: R920–R922.

- Suh EK, Yang A, Kettenbach A, Bamberger C, Michaelis AH, Zhu Z, Elvin JA, Bronson RT, Crum CP, McKeon F. 2006. p53 protects the female germ line during meiotic arrest. *Nature* **444**: 624–628.
- Takeo S, Lake CM, Morais-de-Sa E, Sunkel CE, Hawley RS. 2011. Synaptonemal complex-dependent centromeric clustering and the initiation of synapsis in *Drosophila* oocytes. *Curr Biol* **21**: 1845–1851.
- Tanneti NS, Landy K, Joyce EF, McKim KS. 2011. A pathway for synapsis initiation during zygotene in *Drosophila* oocytes. *Curr Biol* **21**: 1852–1857.
- Terasawa M, Ogawa T, Tsukamoto Y, Ogawa H. 2008. Sae2p phosphorylation is crucial for cooperation with Mre11p for resection of DNA double-strand break ends during meiotic recombination in *Saccharomyces cerevisiae*. *Genes Genet Syst* **83**: 209–217.
- Terentiev Y, Johnson R, Neale MJ, Khisroon M, Bishop-Bailey A, Goldman AS. 2010. Evidence that *MEK1* positively promotes interhomologue double-strand break repair. *Nucleic Acids Res* **38**: 4349–4360.
- Thacker D, Mohibullah N, Zhu X, Keeney S. 2014. Homologue engagement controls meiotic DNA break number and distribution. *Nature* **510**: 241–246.
- Thompson DA, Stahl FW. 1999. Genetic control of recombination partner preference in yeast meiosis. Isolation and characterization of mutants elevated for meiotic unequal sister-chromatid recombination. *Genetics* **153**: 621–641.
- Titen SW, Lin HC, Bhandari J, Golic KG. 2014. Chk2 and p53 regulate the transmission of healed chromosomes in the *Drosophila* male germline. *PLoS Genet* **10**: e1004130.
- Tomita K, Cooper JP. 2006. The meiotic chromosomal bouquet: SUN collects flowers. *Cell* **125**: 19–21.
- Tonami Y, Murakami H, Shirahige K, Nakanishi M. 2005. A checkpoint control linking meiotic S phase and recombination initiation in fission yeast. *Proc Natl Acad Sci* **102**: 5797–5801.
- Tougan T, Kasama T, Ohtaka A, Okuzaki D, Saito TT, Russell P, Nojima H. 2010. The Mek1 phosphorylation cascade plays a role in meiotic recombination of *Schizosaccharomyces pombe*. *Cell Cycle* **9**: 4688–4702.
- Trelles-Sticken E, Loidl J, Scherthan H. 1999. Bouquet formation in budding yeast: Initiation of recombination is not required for meiotic telomere clustering. *J Cell Sci* **112**: 651–658.
- Trelles-Sticken E, Adelfalk C, Loidl J, Scherthan H. 2005a. Meiotic telomere clustering requires actin for its formation and cohesin for its resolution. *J Cell Biol* **170**: 213–223.
- Trelles-Sticken E, Bonfils S, Sollier J, Geli V, Scherthan H, de La Roche Saint-Andre C. 2005b. Set1- and Clb5-deficiencies disclose the differential regulation of centromere and telomere dynamics in *Saccharomyces cerevisiae* meiosis. *J Cell Sci* **118**: 4985–4994.
- Tsubouchi T, Roeder GS. 2005. A synaptonemal complex protein promotes homology-independent centromere coupling. *Science* **308**: 870–873.
- Tung KS, Hong EJ, Roeder GS. 2000. The pachytene checkpoint prevents accumulation and phosphorylation of the meiosis-specific transcription factor Ndt80. *Proc Natl Acad Sci* **97**: 12187–12192.
- Turner JM, Mahadevaiah SK, Fernandez-Capetillo O, Nussenzweig A, Xu X, Deng CX, Burgoyne PS. 2005. Silencing of unsynapsed meiotic chromosomes in the mouse. *Nat Genet* **37**: 41–47.
- Turner JM, Mahadevaiah SK, Ellis PJ, Mitchell MJ, Burgoyne PS. 2006. Pachytene asynapsis drives meiotic sex chromosome inactivation and leads to substantial postmeiotic repression in spermatids. *Dev Cell* **10**: 521–529.
- Usui T, Kanehara Y. 2013. Elevated Rad53 kinase activity influences formation and interhomolog repair of meiotic DNA double-strand breaks in budding yeast. *Biochem Biophys Res Commun* **441**: 593–599.
- Usui T, Ogawa H, Petrini JH. 2001. A DNA damage response pathway controlled by Tel1 and the Mre11 complex. *Mol Cell* **7**: 1255–1266.
- Vader G, Blitzblau HG, Tame MA, Falk JE, Curtin L, Hochwagen A. 2011. Protection of repetitive DNA borders from self-induced meiotic instability. *Nature* **477**: 115–119.
- van Heemst D, Heyting C. 2000. Sister chromatid cohesion and recombination in meiosis. *Chromosoma* **109**: 10–26.
- Vasileva A, Hopkins KM, Wang X, Weisbach MM, Friedman RA, Wolgemuth DJ, Lieberman HB. 2013. The DNA damage checkpoint protein RAD9A is essential for male meiosis in the mouse. *J Cell Sci* **126**: 3927–3938.
- Vigodner M. 2009. Sumoylation precedes accumulation of phosphorylated H2AX on sex chromosomes during their meiotic inactivation. *Chromosome Res* **17**: 37–45.
- Wang H, Hoog C. 2006. Structural damage to meiotic chromosomes impairs DNA recombination and checkpoint control in mammalian oocytes. *J Cell Biol* **173**: 485–495.
- Wang Y, Chang CY, Wu JF, Tung KS. 2011. Nuclear localization of the meiosis-specific transcription factor Ndt80 is regulated by the pachytene checkpoint. *Mol Biol Cell* **22**: 1878–1886.
- Woglar A, Jantsch V. 2013. Chromosome movement in meiosis I prophase of *Caenorhabditis elegans*. *Chromosoma* **123**: 15–24.
- Woglar A, Daryabeigi A, Adamo A, Habacher C, Machacek T, La Volpe A, Jantsch V. 2013. Matefin/SUN-1 phosphorylation is part of a surveillance mechanism to coordinate chromosome synapsis and recombination with meiotic progression and chromosome movement. *PLoS Genet* **9**: e1003335.
- Wojtasz L, Daniel K, Roig I, Bolcun-Filas E, Xu H, Boonsanay V, Eckmann CR, Cooke HJ, Jasini M, Keeney S, et al. 2009. Mouse HORMAD1 and HORMAD2, two conserved meiotic chromosomal proteins, are depleted from synapsed chromosome axes with the help of TRIP13 AAA-ATPase. *PLoS Genet* **5**: e1000702.
- Wojtasz L, Cloutier JM, Baumann M, Daniel K, Varga J, Fu J, Anastassiadis K, Stewart AF, Remenyi A, Turner JM, et al. 2012. Meiotic DNA double-strand breaks and chromosome asynapsis in mice are monitored by distinct HORMAD2-independent and -dependent mechanisms. *Genes Dev* **26**: 958–973.

V.V. Subramanian and A. Hochwagen

- Wu HY, Burgess SM. 2006. Two distinct surveillance mechanisms monitor meiotic chromosome metabolism in budding yeast. *Curr Biol* **16**: 2473–2479.
- Wu HY, Ho HC, Burgess SM. 2010. Mek1 kinase governs outcomes of meiotic recombination and the checkpoint response. *Curr Biol* **20**: 1707–1716.
- Wynne DJ, Rog O, Carlton PM, Dernburg AF. 2012. Dynein-dependent processive chromosome motions promote homologous pairing in *C. elegans* meiosis. *J Cell Biol* **196**: 47–64.
- Xu L, Weiner BM, Kleckner N. 1997. Meiotic cells monitor the status of the interhomolog recombination complex. *Genes Dev* **11**: 106–118.
- Yin Y, Smolikove S. 2013. Impaired resection of meiotic double-strand breaks channels repair to nonhomologous end joining in *Caenorhabditis elegans*. *Mol Cell Biol* **33**: 2732–2747.
- You Z, Chahwan C, Bailis J, Hunter T, Russell P. 2005. ATM activation and its recruitment to damaged DNA require binding to the C terminus of Nbs1. *Mol Cell Biol* **25**: 5363–5379.
- Zakharyevich K, Ma Y, Tang S, Hwang PY, Boiteux S, Hunter N. 2010. Temporally and biochemically distinct activities of Exo1 during meiosis: Double-strand break resection and resolution of double Holliday junctions. *Mol Cell* **40**: 1001–1015.
- Zanders S, Alani E. 2009. The *pch2Δ* mutation in baker's yeast alters meiotic crossover levels and confers a defect in crossover interference. *PLoS Genet* **5**: e1000571.
- Zanders S, Sonntag Brown M, Chen C, Alani E. 2011. Pch2 modulates chromatid partner choice during meiotic double-strand break repair in *Saccharomyces cerevisiae*. *Genetics* **188**: 511–521.
- Zhang L, Kim KP, Kleckner NE, Storlazzi A. 2011. Meiotic double-strand breaks occur once per pair of (sister) chromatids and, via Mec1/ATR and Tel1/ATM, once per quartet of chromatids. *Proc Natl Acad Sci* **108**: 20036–20041.
- Zou L, Elledge SJ. 2003. Sensing DNA damage through ATRIP recognition of RPA-ssDNA complexes. *Science* **300**: 1542–1548.

Cold Spring Harbor Perspectives in Biology

The Meiotic Checkpoint Network: Step-by-Step through Meiotic Prophase

Vijayalakshmi V. Subramanian and Andreas Hochwagen

Cold Spring Harb Perspect Biol 2014; doi: 10.1101/cshperspect.a016675

Subject Collection [DNA Recombination](#)

Meiotic Recombination: The Essence of Heredity
Neil Hunter

Regulation of Recombination and Genomic Maintenance
Wolf-Dietrich Heyer

Initiation of Meiotic Homologous Recombination: Flexibility, Impact of Histone Modifications, and Chromatin Remodeling
Lóránt Székvölgyi, Kunihiro Ohta and Alain Nicolas

Mechanism and Regulation of Meiotic Recombination Initiation
Isabel Lam and Scott Keeney

Homologous Recombination and Human Health: The Roles of BRCA1, BRCA2, and Associated Proteins
Rohit Prakash, Yu Zhang, Weiran Feng, et al.

Cell Biology of Mitotic Recombination
Michael Lisby and Rodney Rothstein

DNA-Pairing and Annealing Processes in Homologous Recombination and Homology-Directed Repair
Scott W. Morrical

An Overview of the Molecular Mechanisms of Recombinational DNA Repair
Stephen C. Kowalczykowski

Recombination, Pairing, and Synapsis of Homologs during Meiosis
Denise Zickler and Nancy Kleckner

DNA Strand Exchange and RecA Homologs in Meiosis
M. Scott Brown and Douglas K. Bishop

Meiosis and Maternal Aging: Insights from Aneuploid Oocytes and Trisomy Births
Mary Herbert, Dimitrios Kalleas, Daniel Cooney, et al.

Mismatch Repair during Homologous and Homeologous Recombination
Maria Spies and Richard Fishel

Mechanisms of Gene Duplication and Amplification
Andrew B. Reams and John R. Roth

The Role of Double-Strand Break Repair Pathways at Functional and Dysfunctional Telomeres
Ylli Doksanli and Titia de Lange

For additional articles in this collection, see <http://cshperspectives.cshlp.org/cgi/collection/>

**All Modifications and
Oligo Types Synthesized**
Long Oligos • Fluorescent • Chimeric • DNA • RNA • Antisense

Oligo Modifications?
Your wish is our command.

Mediators of Homologous DNA Pairing

Alex Zelensky, Roland Kanaar and Claire Wyman

Regulation of DNA Pairing in Homologous Recombination

James M. Daley, William A. Gaines, YoungHo Kwon, et al.

For additional articles in this collection, see <http://cshperspectives.cshlp.org/cgi/collection/>

**All Modifications and
Oligo Types Synthesized**

Long Oligos • Fluorescent • Chimeric • DNA • RNA • Antisense

Oligo Modifications?
Your wish is our command.