Photoreceptor Cilia and Retinal Ciliopathies

Kinga M. Bujakowska, Qin Liu, and Eric A. Pierce

Ocular Genomics Institute, Massachusetts Eye and Ear Infirmary, Department of Ophthalmology, Harvard Medical School, Boston, Massachusetts 02114

Correspondence: eric_pierce@meei.harvard.edu

Photoreceptors are sensory neurons designed to convert light stimuli into neurological responses. This process, called phototransduction, takes place in the outer segments (OS) of rod and cone photoreceptors. OS are specialized sensory cilia, with analogous structures to those present in other nonmotile cilia. Deficient morphogenesis and/or dysfunction of photoreceptor sensory cilia (PSC) caused by mutations in a variety of photoreceptor-specific and common cilia genes can lead to inherited retinal degenerations (IRDs). IRDs can manifest as isolated retinal diseases or syndromic diseases. In this review, we describe the structure and composition of PSC and different forms of ciliopathies with retinal involvement. We review the genetics of the IRDs, which are monogenic disorders but genetically diverse with regard to causality.

Photoreceptors are sensory neurons designed to convert light stimuli into electrical responses, a process called phototransduction. Phototransduction takes place in the highly specialized compartment of photoreceptors, the outer segment (OS) (Pearing et al. 2013; Molday and Moritz 2015). The OS of the rod and cone photoreceptors differ in structure and protein composition, related to their functional adaptation, in which rods have high sensitivity necessary in dim light and cones are responsible for the high-resolution color vision working in bright light (Lamb and Pugh 2006; Lamb et al. 2007). Research over the past decade on the genetic and molecular components of photoreceptors in vertebrate retinae has led to the clear recognition that photoreceptor OS are specialized sensory cilia (Liu et al. 2007a; Ramamurthy and Cayouette 2009; Khanna 2015). Deficient

morphogenesis and/or dysfunction of photoreceptor sensory cilia (PSC) caused by mutations in a variety of photoreceptor-specific and common cilia genes can lead to a group of clinical manifestations, called inherited retinal degenerations (IRDs). In this review, we will discuss the structure and composition of PSC and different forms of ciliopathies with retinal involvement.

SPECIALIZED PHOTORECEPTOR SENSORY CILIA

In vertebrate retina, the visual function depends on the formation of complex sensory cilia of rod and cone photoreceptors. Photoreceptors are highly polarized neurons, composed of four distinct compartments: the OS, the inner segment (IS), the nucleus and a short axon extending to second order neurons (bipolar and hor-

K.M. Bujakowska et al.

horizontal cells) (Fig. 1) (Kennedy and Malicki 2009; Pearing et al. 2013; Molday and Moritz 2015). OS are ciliary organelles with analogous structure to the primary sensory cilia in other cell types (Rosenbaum and Witman 2002; Liu

et al. 2007a; Ramamurthy and Cayouette 2009; Khanna 2015). The recognition of PSCs as distinct morphological structures is valuable for the study of photoreceptor cell biology and disease pathogenesis. Studies of genes involved in IRDs and other ciliopathies have identified dozens of novel components of the PSCs. A comprehensive proteomic study of mouse PSCs has identified ~2000 proteins in this organelle, out of which hundreds are present in other cilia (Liu et al. 2007a). These findings have greatly improved our understanding of how photoreceptor cilia are built and maintained, and how these processes are disrupted in disease.

Ciliary Backbone of PSC

The structure of the PSC is analogous to other cilia, where the axoneme arises from the basal body through the transition zone (also called the “connecting cilium”) and extends up to two-thirds of the OS (Fig. 1) (De Robertis 1956; Kaplan et al. 1987). The basal body also nucleates the ciliary rootlet, which extends into the IS, and which is covalently linked to the PSC structure (Yang et al. 2005; Liu et al. 2007a). The basal body contains nine triplet microtubules, two of which extend further to form the axoneme and the third anchors transition fibers linking the basal body to the plasma membrane. The nine doublet microtubules in the transition zone are cross-linked to the surrounding plasma membrane by Y-link structures (Besharse et al. 1985; Horst et al. 1990). These Y-link structures are absent in the rest of the axoneme. The transition zone of rods and cones measures ~0.3 μm in diameter and 1–1.5 μm in length, which is fairly consistent throughout the species (Besharse et al. 1985). This structure was originally called the “connecting cilium” by De Robertis in 1956, when he was studying some of the first electron micrographs of photoreceptor cells (De Robertis 1956). However, as an analogy with other primary sensory cilia, we refer to this region as a transition zone. Above the transition zone, disc morphogenesis takes place where the surrounding plasma membrane transforms into the disc precursors through membrane evagina-

Figure 1. Photoreceptor structure. Schematic representation of the rod photoreceptor with sensory cilia components indicated. The drawings to the left of the photoreceptor represent the cross-sectional view of the microtubule structure of the distal axoneme (Ax), proximal Ax, transition zone (TZ), and basal body (BB). RT, rootlet; OS, outer segment; CP, calyceal process; IS, inner segment; Ncl, nucleus; Syn, synapse.

tion (Steinberg et al. 1980; Ding et al. 2015; Pugh 2015). At the distal part of the PSC axoneme, the double microtubules are reduced to singlets (Fig. 1) (Rosenbaum and Witman 2002; Pearson et al. 2013).

Other Structural Components of Outer Segments

PSCs are highly specialized sensory cilia, adapted for light detection by the presence of tightly packed membranous discs containing visual pigments and other phototransduction proteins (Sjostrand 1953; Nickell et al. 2007; Gilliam et al. 2012). PSCs are among the largest of mammalian cilia (Pan et al. 2005) and, like other cilia, they are comprised of a cytoskeleton backbone and a membrane domain, which is distinct from the surrounding plasma membrane (Steinberg et al. 1980; Molday and Molday 1987). In murine rods, the numerous membranous discs in the PSC compartment are stacked at a density of ~ 30 discs per micrometer, which is thought to be constant throughout species (Nickell et al. 2007; Gilliam et al. 2012). Such OS organization provides a large surface area for optimized photon capture and rapid signal transduction reactions to occur. Rhodopsin is the most abundant disc membrane protein, organized as rows of dimers with a density of $\sim 48,000$ monomers per μm^2 (Fotiadis et al. 2003). With this high density in the disc membranes, rhodopsin plays an important structural role apart from being the main visual pigment in the retina (Wang and Deretic 2014). The rim of the photoreceptor discs contains two tetraspanins: Rds/peripherin-2 (PRPH2) and retinal OS membrane protein 1 (ROM1), which facilitate the folding of the OS discs and are crucial for rim formation and sorting of the OS proteins during the OS biogenesis (Molday et al. 1987; Goldberg and Molday 1996; Arikawa et al. 2011). PRPH2 and its homolog ROM1 both form homodimers and then associate together to form tetrameric complexes, exclusively present at the disc rims (Molday et al. 1987; Goldberg and Molday 1996; Arikawa et al. 2011). Two other membrane proteins prominin 1 (PROM1) and cadherin-related family mem-

ber 1 (CDHR1) were associated with the open lamellar evaginations in rod and cone discs in *Xenopus laevis* and mice, respectively (Rattner et al. 2001; Han et al. 2012). PSC are responsible for mediating the sensory transduction of the visual system with a number of proteins involved in this process, including the above-mentioned rhodopsin. Most of these proteins are expressed specifically in PSC and, when mutated, cause nonsyndromic IRDs (Table 1) (Dryja et al. 1990; Farrar et al. 1990; Kajiwarra et al. 1991, 1994; Travis et al. 1991; Bascom et al. 1992; Rosenfeld et al. 1992; Dryja et al. 1993; Maw et al. 2000; Yang et al. 2008). Studies of mutant animals have shown that the above-mentioned proteins are essential for OS disc morphogenesis and maintenance (Sanyal et al. 1980; Clarke et al. 2000; Rattner et al. 2001; Dellett et al. 2015).

PROTEIN TRANSPORT TO PSC

A unique feature of the photoreceptor OS is the high level of its renewal. Each day $\sim 10\%$ of the OS is shed from the distal tip, which is replaced by new disc formation at the base of the PSC (Young 1967). This necessitates a robust system of protein synthesis in the IS and efficient trafficking of selected proteins to the photoreceptor OS.

Intraflagellar Transport in PSC

The axoneme, initiated at the mother centriole, is built and maintained by extending its distal (+) end (Pedersen and Rosenbaum 2008). Because protein synthesis occurs in the IS, the axoneme building blocks need to be transported to the distal end via intraflagellar transport (IFT) (Rosenbaum and Witman 2002; Pedersen and Rosenbaum 2008; Taschner et al. 2012). The anterograde transport from the base to the tip of the axoneme is mediated by IFT complex B (IFT-B), where kinesin-2 is the motor protein (Rosenbaum and Witman 2002). Kinesin-2 is a heterotrimeric protein composed of Kif3A, Kif3B, and KAP, which is further associated with 14 other IFT proteins that bind cargo molecules (Taschner et al. 2012). Once the axoneme

K.M. Bujakowska et al.

Table 1. Genes associated with nonsyndromic retinal degeneration

Gene	Inheritance pattern	Nonsyndromic form	Syndromic form	Notes and references
Genes coding axoneme-associated proteins				
<i>ARL6</i>	AR	RP	BBS	New retina-specific exon present (Chiang et al. 2004; Fan et al. 2004; Aldahmesh et al. 2009; Pretorius et al. 2010)
<i>BBS1</i>	AR	RP	BBS	Nishimura et al. 2001; Mykytyn et al. 2002; Estrada-Cuzcano et al. 2012a
<i>BBS2</i>	AR	RP	BBS	Consugar et al. 2014; Shevach et al. 2015
<i>BBS9</i>	AR	RP	BBS	Nishimura et al. 2005; Abu-Safieh et al. 2012
<i>C2orf71</i>	AR	RP	-	Putative cilia function, exclusive eye expression (Collin et al. 2010; Nishimura et al. 2010; Kevany et al. 2015)
<i>C8orf37</i>	AR	RP	BBS	Estrada-Cuzcano et al. 2012b; Heon et al. 2016; Khan et al. 2016
<i>CEP164</i>	AR	LCA	SLS	Chaki et al. 2012
<i>CEP290</i>	AR	LCA	BBS, JBS, MKS, SLS	den Hollander et al. 2006; Sayer et al. 2006; Valente et al. 2006; Baala et al. 2007a; Helou et al. 2007; Frank et al. 2008; Leitch et al. 2008
<i>CLRN1</i>	AR	RP	USH	Joensuu et al. 2001; Khan et al. 2011
<i>FAM161A</i>	AR	RP	-	Bandah-Rozenfeld et al. 2010; Langmann et al. 2010
<i>IFT140</i>	AR	RP, LCA	JATD, MZSDS	Perrault et al. 2012; Schmidts et al. 2013a; Bifari et al. 2015; Xu et al. 2015
<i>IFT172</i>	AR	RP	BBS, JATD, MZSDS	Halbritter et al. 2013; Bujakowska et al. 2014
<i>IQCB1</i>	AR	LCA	SLS	Otto et al. 2005; Estrada-Cuzcano et al. 2011; Stone et al. 2011
<i>KIZ</i>	AR	RP	-	El Shamieh et al. 2014
<i>LAC5</i>	AR	LCA	-	den Hollander et al. 2007
<i>MAK</i>	AR	RP	-	Ozgül et al. 2011; Tucker et al. 2011
<i>NEK2</i>	AR	RP	-	Nishiguchi et al. 2013
<i>OFD1</i>	XL	RP	OFD, JBS	Ferrante et al. 2001; Coene et al. 2009; Webb et al. 2012
<i>RAB28</i>	AR	CRD	-	Roosing et al. 2013
<i>RP1</i>	AR, AD	RP	-	Guillonneau et al. 1999; Pierce et al. 1999
<i>RP1L1</i>	AR, AD	RP, OMD	-	There are some doubts about this gene being truly associated with IRD, because this gene is highly polymorphic and some mutations were seen in the

Continued

Table 1. Continued

Gene	Inheritance pattern	Nonsyndromic form	Syndromic form	Notes and references
				controls (Bowne et al. 2003; Yamashita et al. 2009; Akahori et al. 2010; Davidson et al. 2013) Hardcastle et al. 1999; Mears et al. 1999
<i>RP2</i>	XL	RP	-	
<i>RPGR</i>	XL	RP, CD, MD	RP with hearing loss and sinorespiratory infections	Extraocular phenotypes may not be related to <i>RPGR</i> , because it mapped to a 43.6-Mb interval with 215 genes including <i>OFD1</i> (Meindl et al. 1996; Roepman et al. 1996; Ayyagari et al. 2002; Yang et al. 2002; Zito et al. 2003)
<i>RPGRIP1</i>	AR	LCA, CRD	-	Dryja et al. 2001; Hameed et al. 2003
<i>SPATA7</i>	AR	LCA, RP	-	Wang et al. 2009
<i>TOPORS</i>	AD	RP	-	Chakarova et al. 2007, 2011
<i>TTC8</i>	AR	RP	BBS	Ansley et al. 2003; Riazuddin et al. 2010
<i>USH2A</i>	AR	RP	USH	Eudy et al. 1998; Rivolta et al. 2000
<i>WDR19</i>	AR	RP	SLS, CED, JATD	Bredrup et al. 2011; Coussa et al. 2013
Other structural OS proteins				
<i>CDHR1</i>	AR	CRD		Henderson et al. 2010; Ostergaard et al. 2010
<i>EYS</i>	AR	RP		Abd El-Aziz et al. 2008; Collin et al. 2008
<i>FSCN2</i>	AD	RP, MD		There are some doubts about this gene being truly associated with IRD, because a frameshift c.208delG is a common polymorphism in the Asian population (Wada et al. 2001, 2003; Zhang et al. 2007; Shin et al. 2010)
<i>PROM1</i>	AR, AD	RP, MD		Maw et al. 2000; Yang et al. 2008
<i>PRPH2</i>	AD, digenic with ROM1	RP, MD		Kajiwara et al. 1991, 1994; Travis et al. 1991
<i>ROM1</i>	AD, digenic with ROM1	RP		Bascom et al. 1992; Kajiwara et al. 1994
<i>TULP1</i>	AR	RP		Banerjee et al. 1998; Hagstrom et al. 1998; Larsson et al. 1998
Genes involved with the POS sensory function (phototransduction cascade and retinoid cycle in the photoreceptors)				
<i>ABCA4</i>	AR	STGD, RP, CRD		Allikmets et al. 1997; Sun and Nathans 1997; Cremers et al. 1998; Martínez-Mir et al. 1998
<i>CNGA1</i>	AR	RP		Dryja et al. 1995
<i>CNGA3</i>	AR	ACHR		Kohl et al. 1998

Continued

K.M. Bujakowska et al.

Table 1. *Continued*

Gene	Inheritance pattern	Nonsyndromic form	Syndromic form	Notes and references
<i>CNGB1</i>	AR	RP		Bareil et al. 2001
<i>CNGB3</i>	AR	ACHM, CD		Kohl et al. 2000
<i>GNAT1</i>	AD, AR	CSNB		Dryja et al. 1996
<i>GNAT2</i>	AR	ACHM		Aligianis et al. 2002; Kohl et al. 2002
<i>GRK</i>	AR	CSNB		Yamamoto et al. 1997
<i>GUCA1A</i>	AD	CD, CRD		Payne et al. 1998; Sokal et al. 1998
<i>GUCA1B</i>	AD	RP, MD		Sato et al. 2005
<i>GUCY2D</i>	AR, AD	LCA, CRD		Perrault et al. 1996; Kelsell et al. 1998
<i>OPN1LW</i>	XL	Deuteranopia, blue cone monochromacy		Nathans et al. 1986; Winderickx et al. 1992; Ayyagari et al. 1999
<i>OPN1MW</i>	XL	Protanopia, blue cone monochromacy		Nathans et al. 1986; Ayyagari et al. 1999
<i>OPN1SW</i>	AD	Tritanopia		Nathans et al. 1992; Weitz et al. 1992a,b
<i>PDE6A</i>	AR	RP		Huang et al. 1995
<i>PDE6B</i>	AR, AD	RP, CSNB		McLaughlin et al. 1993; Gal et al. 1994
<i>PDE6C</i>	AR	CD, ACHM		Thiadens et al. 2009
<i>PDE6G</i>	AR	RP		Dvir et al. 2010
<i>RDH12</i>	AR, AD	LCA, RP		Janecke et al. 2004; Perrault et al. 2004; Fingert et al. 2008
<i>RGS9</i>	AR	Delayed cone adaptation		Nishiguchi et al. 2004
<i>RGS9BP</i>	AR	Delayed cone adaptation		Nishiguchi et al. 2004
<i>RHO</i>	AD, AR	RP, CSNB		Dryja et al. 1990; Farrar et al. 1990; Rosenfeld et al. 1992; Dryja et al. 1993
<i>SAG</i>	AR	RP, CSNB		Fuchs et al. 1995; Nakazawa et al. 1998

ACHM, Achromatopsia; BBS, Bardet–Biedl syndrome; CD, cone dystrophy; CED, cranioectodermal dysplasia, also known as Sensenbrenner syndrome; CRD, cone–rod dystrophy; CSNB, congenital stationary night blindness; JBS, Joubert syndrome; JATD, Jeune asphyxiating thoracic dystrophy; LCA, Leber congenital amaurosis; MKS, Meckel–Gruber syndrome; MZSDS, Mainzer–Saldino syndrome; OFD, oral–facial–digital syndrome; OMD, occult macular dystrophy; RP, retinitis pigmentosa; SLS, Senior–Løken syndrome; STGD, Stargardt disease; USH, Usher syndrome.

and other PSC components have been delivered to the tip of the cilium, the IFT-B components are recycled back to the base of the cilium by retrograde transport mediated by IFT complex A (IFT-A) (Rosenbaum and Witman 2002). Dynein-2 is the motor protein of IFT-A and it is associated with six other IFT proteins (Taschner et al. 2012). Apart from the IFT complexes, Bardet–Biedl syndrome proteins (BBSome) are

also involved in the transport of membrane proteins to the cilium (Taschner et al. 2012; Williams et al. 2014).

Because 10% of the PSC is shed and renewed every day, the necessity for the retrograde transport in this cell type was not clear. However, identification of IRD patients with mutations in genes coding for retrograde transport proteins (e.g., *TTC21B*) and pro-

teins involved in switching from anterograde to retrograde IFT direction (e.g., *IFT172*) underlines the importance of transport in both directions for the development and maintenance of PSC (Liu et al. 2010; Davis et al. 2011; Halbritter et al. 2013; Bujakowska et al. 2014).

Transport of Membrane Proteins to PSC

Even though the plasma membrane surrounding the photoreceptor cilium is continuous with the plasma membrane of the cell body, its protein composition is different. In addition, the membranous discs in the rod photoreceptors are distinct from the surrounding plasma membrane (Steinberg et al. 1980; Molday and Molday 1987). This selective protein content in PSC membranes is established by diffusional barriers present at the base of the cilium and within the transition zone (Pearring et al. 2013; Wang and Deretic 2014; Khanna 2015). The molecular composition of the diffusional barrier is not fully understood, although certain proteins like Septin 2 and CEP290 are thought to play an important role (Pearring et al. 2013; Wang and Deretic 2014).

Because rhodopsin is the most abundant protein in the PSC, its photoreceptor OS transport has been studied in detail. After synthesis in the IS endoplasmic reticulum and transport through the Golgi and *trans*-Golgi network, rhodopsin is sorted into vesicles destined for the OS. This is achieved thanks to the presence of specific sequence signatures (e.g., VXPX and FR motifs), which facilitate interaction with a ciliary targeting molecules Arf4 and ASAP1 (Deretic et al. 2005; Wang and Deretic 2014). Further interaction with FIP3 and small GTPases Rab6, Rab8, and Rab11 directs the vesicle to the base of the OS for fusion with the membrane (Deretic et al. 2005; Pearring et al. 2013; Wang and Deretic 2014). Further, rhodopsin molecules are transported through the transition zone to the site of the disc morphogenesis by two motor proteins, kinesin II and myosin VIIa, as shown in *Kif3a* and *Myo7a* knockout mice (Liu et al. 1999; Williams 2002). As mentioned before, kinesin II mediates

a microtubule-dependent anterograde IFT (Rosenbaum and Witman 2002). Myosin VIIa is an actin-dependent motor molecule and, in mouse photoreceptors, it locates to the periciliary membrane complex; however, in primates, it locates to the calyceal processes (Sahly et al. 2012). It is, therefore, unclear whether mutations in *Myosin VIIa* in humans also lead to the aberrant opsin trafficking as shown in mice (Liu et al. 1999). Immunoelectron microscope studies of *Rana pipiens* frog photoreceptors, revealed that actin is present not only in calyceal processes but also at the sites of disc morphogenesis, suggesting involvement of actin-mediated transport in protein delivery to the forming discs (Chaitin et al. 1984). Little is known about OS targeting of other OS-specific transmembrane proteins, apart from retinol dehydrogenase (RDH8), which also contains the VXPX motif and PRPH2, which has its own OS-targeting sequence (Pearring et al. 2013).

Photoactivated Protein Diffusion

The base of the OS does not contain a selective barrier for the soluble proteins as shown in mice by the light-activated translocation of phototransduction proteins: transducin, arrestin, and recoverin (Sokolov et al. 2002; Calvert et al. 2006). Furthermore, this translocation is thought to be energy-independent, implying that the protein movement occurs by simple diffusion (Nair et al. 2005; Calvert et al. 2010). The diffusion of the proteins is dependent, however, on the steric interactions between the molecules and cell structures, termed steric volume exclusion, which reduces the entry of larger molecular weight proteins to the OS (Najafi and Calvert 2012). Light-mediated translocation of these proteins is thought to play a role in adaptation to different light conditions, in which for instance concentrating transducin in the rod OS in the darkness amplifies the phototransduction signal, and translocation of arrestin to OS in light conditions terminates transducin activation and accelerates photopigment recovery (Pearring et al. 2013). A neuroprotective role for the light-induced protein translocation has also been suggested (Pearring et al. 2013).

K.M. Bujakowska et al.

RETINAL CILIOPATHIES

Mutations in genes coding for ciliary proteins lead to ciliopathies, rare genetic disorders that may affect one or more organs, including the retina, central nervous system, olfactory epithelium, cardiovascular system, liver, kidney, skeletal system, gonads, and adipose tissue (Goetz and Anderson 2010; Patel and Honoré 2010; Mockel et al. 2011; Waters and Beales 2011). In this review, we will focus on ciliopathies that involve the retina, manifesting most commonly as retinitis pigmentosa (RP) (Hamel 2006; Hartong et al. 2006; Berger et al. 2010) or Leber congenital amaurosis (LCA) (Weleber 2002; Chung and Traboulsi 2009). RP is a condition that primarily affects rod photoreceptors and retinal pigment epithelium. It is the most frequent cause of the IRDs, with a prevalence of $\sim 1/3500$ and accounting for roughly 25% of vision loss in adults (Hamel 2006; Hartong et al. 2006; Berger et al. 2010). It may start in the first or second decade of life, often with nyctalopia and peripheral vision loss as early symptoms, because of the dysfunction of PSCs and photoreceptor cell death in the peripheral retina. In many cases, the disease progresses to include central vision loss as well, because of eventual dysfunction of PSCs and death of photoreceptor cells in the macula (central retina) (Fig. 2) (Hamel 2006; Hartong et al. 2006; Berger et al. 2010). LCA affects rods and cones and leads to vision loss in infancy or early childhood (Weleber 2002; den Hollander et al. 2008; Chung and Traboulsi 2009). LCA is rare, with a population frequency of $\sim 1/50,000$, yet affecting $\sim 20\%$ of children attending schools for the blind (Weleber 2002; Koenekoop 2004; Berger et al. 2010). Other subtypes of IRD are present in ciliopathy patients and often involve cone photoreceptors and the macula (Michaelides et al. 2006; Estrada-Cuzcano et al. 2012c).

Nonsyndromic Retinal Ciliopathies

As mentioned above, photoreceptor OS can be regarded as specialized cilia designed to detect light and to convert this information into a biochemical signal. Therefore, we consider that all

proteins that participate in this sensory function, as well as proteins that build the PSC structure, are in effect cilia proteins. Consequently, we can distinguish two groups of retinal ciliopathies: (1) affecting the structure, and (2) the sensory function of photoreceptor OS.

Mutations in Genes Disrupting POS Structure

There are currently 36 known genes that have been identified to harbor mutations that disrupt PSC structure and can lead to an isolated or syndromic retinal degeneration (Table 1). Thirteen genes that encode axoneme or basal body-associated proteins (*C2ORF71*, *FAM161A*, *KIZ*, *LCA5*, *MAK*, *NEK2*, *RAB28*, *RPGRIP1*, *RP1*, *RP1L1*, *RP2*, *SPATA7*, *TOPORS*) and seven genes coding for other structural PSC components (*CDHR1*, *EYS*, *FSCN2*, *PROM1*, *PRPH2*, *ROM1*, *TULP1*) have been exclusively associated with nonsyndromic retinal degeneration (Kajiwara et al. 1991, 1994; Bascom et al. 1992; Banerjee et al. 1998; Hagstrom et al. 1998; Guillonnet et al. 1999; Hardcastle et al. 1999; Mears et al. 1999; Pierce et al. 1999; Maw et al. 2000; Dryja et al. 2001; Wada et al. 2001, 2003; Chakarova et al. 2007; den Hollander et al. 2007; Abd El-Aziz et al. 2008; Collin et al. 2008; Wang et al. 2009; Akahori et al. 2010; Bandah-Rozenfeld et al. 2010; Collin et al. 2010; Henderson et al. 2010; Langmann et al. 2010; Nishimura et al. 2010; Ostergaard et al. 2010; Ozgöl et al. 2011; Tucker et al. 2011; Estrada-Cuzcano et al. 2012b; Davidson et al. 2013; Nishiguchi et al. 2013; Roosing et al. 2013; El Shamieh et al. 2014). A query of the human proteome map (Kim et al. 2014) shows that nine of these genes (*CDHR1*, *FSCN2*, *MAK*, *PROM1*, *PRPH2*, *ROM1*, *RP1*, *RP1L1*, *TULP1*) are predominantly expressed in the human retina, which corroborates with the retina-specific phenotype. With the exception of *LCA5*, which shows no significant expression in any of the assayed tissues, the remaining genes are also significantly expressed in other human tissues, and it remains unclear why mutations in these genes affect specifically the retina.

Two genes stand apart in IRD ciliopathies, *USH2A* and *CLRN1*. Mutations in these genes

Figure 2. Clinical features of inherited retinal degeneration (IRD). Fundus photos and optical coherence tomography (OCT) images of normal and diseased retinas are shown. (A) Wide-field fundus photo shows the appearance of a normal retina; the optic nerve (ON) is visible. (B) Fundus image from a patient with early retinal degeneration because of retinitis pigmentosa (RP). The arrow shows the pigment changes, which are characteristic of this disorder. (C) Fundus image from a patient with advanced retinal degeneration because of RP. (D) A cross-sectional image of the center of the normal retina (macula; white line in A) obtained with OCT shows normal retinal layers, including the outer nuclear layer (ONL), where photoreceptor cell nuclei are located. The central indentation is normal, and indicates the fovea. The white band showing the ellipsoid zone (EZ, arrow) is generated from the junction of the inner and outer segments of photoreceptor cells, and its presence indicates normal photoreceptor cell and thus PSC structure. (E) The OCT image shows loss of photoreceptor cells peripherally, with the ONL visible only near the fovea. The EZ is evident centrally, but is lost more peripherally, indicating loss of PSCs by the more peripheral photoreceptor cells present. (F) The OCT image shows loss of the ONL and thus all of the photoreceptor cells in the macula of this patient consistent with greatly reduced central vision.

lead to isolated retinal degeneration (Rivolta et al. 2000; Khan et al. 2011) or to deaf-blindness, called Usher syndrome (Eudy et al. 1998; Joensuu et al. 2001). There is only one report of nonsyndromic IRD because of mutations in *CLRN1* (Khan et al. 2011); however, mutations in *USH2A* are the leading cause of the nonsyndromic autosomal recessive IRD, accounting for ~9% of RP patients (Hartong et al. 2006). These two genes will be further discussed in the section on Usher syndrome.

The remaining 12 genes code for axoneme proteins and are associated with isolated retinopathy or syndromic disease (Table 1). In certain cases, the broad spectrum of phenotypes associated with mutations in a given gene can be explained by the primary mutations in the gene, where mutations in the retina-specific transcripts or hypomorphic alleles may lead to the isolated retinal phenotype (Ansley et al. 2003; den Hollander et al. 2006; Coene et al. 2009; Riazuddin et al. 2010; Webb et al. 2012;

Xu et al. 2015). In other cases, the relationship between the primary disease-causing mutation and the phenotype is not clear and epistatic effects of other alleles have been suggested (Badano et al. 2003b, 2006; Estrada-Cuzcano et al. 2012a; Bujakowska et al. 2014). (Both cases will be discussed in detail in the section Broad Phenotypic Spectrum of Ciliopathies and Genetic Modifiers.)

Mutations in Genes Impeding POS Sensory Function

A less obvious subgroup of nonsyndromic retinal ciliopathies are IRDs in which PSC sensory function is affected. This classification is analogous to another ciliopathy, the dominant form of polycystic kidney disease, because of mutations in *PKD1* or *PKD2*, which code for membrane proteins that act as a cilia sensor and a calcium channel, respectively (Ong and Harris 2015). Similarly, genetic mutations that affect

K.M. Bujakowska et al.

phototransduction cascade and retinoid cycle in the photoreceptors are also considered as ciliopathies (Table 1). However, we do not consider as ciliopathies IRDs that are caused by mutations in genes that are expressed in the RPE or code for proteins that function in other compartments of the photoreceptor cell (e.g., splicing factor genes). A comprehensive list of all IRD genes can be found at the Retinal Information Network (RetNet) portal (sph.uth.edu/retnet/home.htm).

Usher Syndrome

Usher syndrome is an autosomal recessive dual impairment of vision and sensorineural hearing with a prevalence of $\sim 1/25,000$ people (Kremer et al. 2006; Millán et al. 2011; Bonnet and El-Amraoui 2012). It is phenotypically and genetically heterogeneous and most of the patients fall into one of the three clinical subtypes of decreasing severity: Usher syndrome type I (USH1), type II (USH2), and type III (USH3).

The most severe form, USH1, is not strictly a ciliopathy because the proteins coded by the six associated genes: *MYO7A* (Weil et al. 1995), *USH1C* (Verpy et al. 2000), *CDH23* (Bolz et al. 2001), *PCDH15* (Ahmed et al. 2001), *USH1G* (Weil et al. 2003), and *CIB2* (Riazuddin et al. 2012), locate to actin-based structures in the periciliary region, called the calyceal processes, and not to cilia themselves (Sahly et al. 2012). The function of these structures is unclear and they might have a structural role supporting the photoreceptor outer segments or be involved in fine tuning of the photoreceptor signaling. Calyceal processes are not present in all of the vertebrates and their absence in rodents may account for the subtle retinal phenotypes in most of the USH1 murine models, in which hearing and vestibular phenotypes are profound (Liu et al. 1999; Di Palma et al. 2001; Libby and Steel 2001; Johnson et al. 2003; Ahmed et al. 2008; Williams 2008; Miyasaka et al. 2013). In the ear, USH1 proteins are involved in the maturation of the stereocilia in the auditory hair cells of the inner-ear cochlea, where they form transient links between the kinocilium and stereocilia during development and the

tip links between the mature stereocilia (Kremer et al. 2006; Bonnet and El-Amraoui 2012; Riazuddin et al. 2012; Sahly et al. 2012).

USH2 is caused by mutations in one of three genes: *USH2A* (Eudy et al. 1998), *GPR98* (Weston et al. 2004), and *DFNB31* (Mburu et al. 2003). The products of these genes locate to the periciliary membrane complex adjacent to the transition zone of the photoreceptor cilia (Liu et al. 2007b; Sahly et al. 2012). In the cochlea, they form the transient ankle links between the developing stereocilia (reviewed in Bonnet and El-Amraoui 2012).

USH3, the least severe form of Usher, is caused by mutations in *CLRN1* or *HARS* (Joensuu et al. 2001; Puffenberger et al. 2012). In mice, *Clarín-1* locates to the base of the photoreceptors cilia and to synaptic ribbons (Zallocchi et al. 2009); however, it does not seem to be essential for the photoreceptor function in rodents, because homozygous *Clrn1* knockout mice show no retinal phenotype (Geller et al. 2009). In the cochlea, *Clarín-1* locates to the apical and basal aspects of stereocilia depending on the developmental stage (Zallocchi et al. 2009). *HARS* codes for a histidyl-tRNA synthetase and, currently, the mechanism of the disease is unknown (Puffenberger et al. 2012).

Of the three Usher types, the most frequent is USH2 (56%–57% cases), in which mutations in the *USH2A* gene are the most common (Millán et al. 2011). USH1 represents 33%–44% of all Usher cases, followed by USH3, with a prevalence of 2% (Yan and Liu 2010; Millán et al. 2011). However, in certain populations (Finland and Ashkenazi Jewish), USH3 reaches 40% of all Usher patients because of founder mutations (Yan and Liu 2010; Millán et al. 2011). Atypical Usher syndromes have also been reported in which the deaf–blindness phenotype is explained by mutations in two different genes: *USH2A* with *PDZD7* and *C2orf71* with *CEP250* (Table 2) (Ebermann et al. 2010; Khateb et al. 2014). Recently, two groups have reported another form of Usher syndrome because of mutations in *CEP78*, which manifest as a cone–rod dystrophy accompanied by sensorineural hearing loss involving mainly high frequencies (Nikopoulos et al. 2016; Sharon et al.

Table 2. Genetic modifiers of ciliopathies

Primary gene	Modifier	Phenotype	Evidence	References
<i>PRPH2/RDS</i> p.(Leu185Pro) <i>ROM1</i>	Digenic inheritance	RP	Three families (11 affected in total)	Kajiwara et al. 1994
p.(Gly80fs) and p.(Leu114fs) <i>BBS2</i>	Digenic inheritance	BBS	In one family, the affected sib carried three alleles, whereas the unaffected sib is the only compound heterozygous for the <i>BBS2</i> mutations; three additional families with a probable triallelic <i>BBS2/BBS6</i> inheritance	Katsanis et al. 2001
p.[Tyr24*] and p.[Gln59*] <i>BBS6</i>				
p.(Gln147*) <i>RPE65</i>		More severe RD	One family with two sibs, targeted analysis of: <i>GUCY2D</i> , <i>RPE65</i> , <i>CRX</i> , <i>AIP1</i> , and <i>RPGRIPI</i>	Silva et al. 2004
p.(Glu102*) <i>BBS1</i>		More severe ciliopathy	In one family of two affected sibs, the sister carrying the additional <i>ALR6</i> allele showed a more severe phenotype	Fan et al. 2004
p.(Met390Arg) <i>BBS1</i> and unknown BBS genotypes		Increased severity of disease, general mutational load	In three families, variant associated with more severe BBS; variant also enriched in BBS patients (14/226) over controls (4/274)	Badano et al. 2006
Mixed ciliopathy cohort	<i>RPGRIPI</i> p.(Ala229Thr)	Presence of RD in ciliopathies	Targeted screening of <i>RPGRIPI</i> ; enrichment of 226Thr variant in ciliopathy patients with RD (43/487) over ciliopathy patients without RD (0/115); functional data, showing a decreased binding activity to RPGR	Khanna et al. 2009
BBS mixed cohort	<i>MKS1</i> p.(Arg123Gln) p.(Asp286Gly) p.(Ile450Thr) p.(Val339Met) (specific to short isoform) <i>TMEM67</i> (c.2241G>A) and p.(Ser320Cys)	More severe ciliopathy phenotype, seizures in five of six patients	Targeted screening of <i>MKS1</i> in BBS cohort, heterozygous potentially modifying changes found in six patients from five families (5/155); functionally, variants ranged from mild hypomorphs to null	Leitch et al. 2008
<i>BBS9</i> and <i>CEP290</i>		General mutational load	Potentially pathogenic alleles found in two families: c.2241G>A affects the canonical splice site and is predicted to lead to exon skipping; p.(Ser320Cys) was functionally null	Leitch et al. 2008
<i>PRPH2</i>	<i>ROM1</i> p.(Arg229His) and/or <i>ABCA4</i> p.(Val2050Leu)	Increased macular involvement in RD	One family, eight affected; microarray mutation detection in 16 genes, full sequencing of <i>PRPH2</i> , <i>ROM1</i> , and <i>ABCA4</i>	Poloschek et al. 2010

Continued

Table 2. *Continued*

Primary gene	Modifier	Phenotype	Evidence	References
<i>USH2A</i>	<i>PDZD7</i> p.(Arg56fs)	More severe RD	One family with two sibs; targeted analysis of <i>PDZD7</i> and other Usher genes; digenic inheritance stated but insufficient genetic data to prove it	Ebermann et al. 2010
<i>CEP290</i>	<i>AHII</i> p.(Asn811Lys) and p.(His758Pro)	More severe neurological phenotype	Targeted screening of <i>AHII</i> in eight patients; variants detected in two patients	Coppieters et al. 2010
<i>NPHP1</i> and unknown NPHP genotypes	<i>AHII</i> p.(Arg830Trp)	Presence of RD	Targeted screening of <i>AHII</i> in 153 NPHP ± RD patients	Louie et al. 2010
Mixed MKS and BBS cohort	<i>C2ORF86</i> (various changes)	General mutational load	Targeted screening of <i>C2ORF86</i> ; enrichment of nonsynonymous coding changes in patients (6/192) versus controls (0/384)	Kim et al. 2010
<i>RPGR</i>	<i>RPGRIP1L</i> p.(Arg744Gln) and <i>IQCB1</i> p.(Ile393Asn)	Severity of RD	Targeted screening of <i>RPGRIP1L</i> , <i>RPGRIP1L</i> , <i>CEP290</i> , and <i>IQCB1</i> in 98 male patients; the results were marginally significant	Fahim et al. 2011
Mixed ciliopathy cohort	<i>TTC21B</i> (various changes)	General mutational load	Targeted screening of <i>TTC21B</i> ; enrichment of pathogenic changes in ciliopathy patients (28/555) over controls (4/305)	Davis et al. 2011
<i>C2orf71</i> p.(Gln1097*)	<i>CEP250</i> p.(Arg1155*)	More severe retinal degeneration + hearing loss	Homozygosity mapping in family with seven affected, subsequent WES in two affected; homozygous <i>CEP250</i> mutation leads to an early-onset severe hearing loss with a mild retinal degeneration and an additional homozygous stop mutation in <i>C2orf71</i> exacerbated the retinal phenotype in three individuals	Khateb et al. 2014
Mixed BBS cohort	<i>NPHP1</i> whole gene deletion and p.(Arg5Leu)	General mutational load in ciliopathies	Targeted analysis of <i>NPHP1</i> in a BBS cohort of 200 families, mutations enriched in the patient population compared with control (incidence of 1.5% (deletion) and 2.5% (missense); functional data for <i>NPHP1</i> -BBS genes interaction)	Lindstrand et al. 2014
<i>PRPH2</i> (c.828+3A>T)	<i>PRPH2</i> p.[(Glu304;Lys310;Gly338)] haplotype in <i>trans</i> with the causal mutation	More severe retinal phenotype	p.[(Glu304;Lys310;Gly338)] haplotype in <i>trans</i> with the splice site mutation was associated with a more severe phenotype as investigated in 62 patients	Shankar et al. 2016

BBS, Bardet–Biedl syndrome; RD, retinal degeneration; RP, retinitis pigmentosa.

2016). Hearing loss can also be part of other syndromes, such as Altrom, as will be discussed later (Mockel et al. 2011).

Other Syndromic IRDs

Based on the presence of particular symptoms, ciliopathies are subdivided into different subtypes, traditionally named after clinicians who first described them. Here, we will review the major syndromes, which involve the retina. Even though these conditions are considered as distinct clinical entities, it is being increasingly recognized that there is a large phenotypic overlap between these diseases, in which characteristic features of two different syndromes are present in the same patient or distinct ciliopathies co-occur in single families (Lehman et al. 2010; Zaki et al. 2011; Valente et al. 2014). Traditional naming of these conditions is therefore often inaccurate and depends on the clinicians' training and their specialty. To overcome this bias, we believe that it is crucial to include in the syndrome naming the underlying molecular cause of the disease (e.g., *AHI1*-associated ciliopathy). However, for the purpose of this review, we describe each syndrome as traditionally called and present the genes associated with them.

Senior–Løken syndrome (SLS) is an autosomal recessive disease characterized by juvenile nephronophthisis (NPHP) and early-onset retinal degeneration (Løken et al. 1961; Senior et al. 1961). NPHP is a medullary cystic kidney disease leading to the end-stage renal failure later in childhood or in adolescence (Ronquillo et al. 2012). About 10% of NPHP patients also have retinal degeneration (Otto et al. 2005; Mockel et al. 2011). So far, eight genes are associated with SLS (*CEP164*, *CEP290*, *IQCB1*, *NPHP1*, *NPHP4*, *SDCCAG8*, *TRAF3IP1*, *WDR19*), although other NPHP-associated genes are mutated in different syndromes involving the retina (Caridi et al. 1998; Otto et al. 2002, 2005, 2010; Sayer et al. 2006; Chaki et al. 2012; Coussa et al. 2013; Bizet et al. 2015). There is a considerable overlap between SLS and other ciliopathies, where almost all SLS genes are associated with other diseases (Fig. 3).

Joubert syndrome (JBS) is a neurological condition characterized by a distinctive abnormality of the midbrain–hindbrain junction and cerebellar vermis hypoplasia, presenting as the molar tooth sign (MTS) on brain imaging (Maria et al. 1997; Valente et al. 2013). These neurological defects correlate with the clinical presentation of hypotonia, ataxia, abnormal breathing, developmental delay, and abnormal ocular movements. JBS patients may also present with retinal degeneration, renal or hepatic defects, polydactyly, and orofacial dysmorphism (Mockel et al. 2011; Valente et al. 2013). The prevalence of JBS is estimated to be between 1/80,000 and 1/100,000 of live births (Valente et al. 2014). Mutations in 26 genes have been reported to cause JBS (*AHI1*, *ARL13B*, *B9D1*, *C5orf42*, *CC2D2A*, *CEP104*, *CEP290*, *CEP41*, *CSPP1*, *INPP5E*, *KIAA0556*, *KIAA0586*, *KIF7*, *NPHP1*, *OFD1*, *RPGRIP1L*, *SRTD1*, *TCTN2*, *TCTN3*, *TECT1*, *TMEM67*, *TMEM138*, *TMEM216*, *TMEM231*, *TMEM237*, *ZNF423*) (Dixon-Salazar et al. 2004; Ferland et al. 2004; Parisi et al. 2004; Sayer et al. 2006; Baala et al. 2007b; Delous et al. 2007; Cantagrel et al. 2008; Gorden et al. 2008; Noor et al. 2008; Bielas et al. 2009; Coene et al. 2009; Edvardson et al. 2010; Dafinger et al. 2011; Garcia-Gonzalo et al. 2011; Huang et al. 2011; Chaki et al. 2012; Lee et al. 2012a,b; Srour et al. 2012a,b; Thomas et al. 2012; Romani et al. 2014; Shaheen et al. 2014; Thomas et al. 2014; Tuz et al. 2014; Sanders et al. 2015). All except one of the above genes cause autosomal-recessive or X-linked disease; *ZNF423* has been associated with a dominant JBS form, although this association showed limited genetic evidence (Chaki et al. 2012). *ZNF423* is also the only JBS gene, which is not associated with the cilium but with the DNA damage response pathway (Chaki et al. 2012).

Meckel–Gruber syndrome, also known as Meckel syndrome (MKS) is a neonatal lethal autosomal recessive disorder defined by the malformation of the central nervous system (occipital encephalocele), cystic kidneys, and liver fibrosis (Wright et al. 1994; Logan et al. 2011). Other features that may be present are postaxial or preaxial polydactyly, skeletal dys-

[illegible]

plasia, cleft lip/palate, microphthalmia, optic nerve coloboma, heart defects, genital anomalies, and complete or partial situs inversus (Logan et al. 2011). Mutations in 12 genes have been associated with MKS (*B9D1*, *B9D2*, *CC2D2A*, *CEP290*, *KIF14*, *MKS1*, *NPHP3*, *RPGRIP1L*, *TCTN2*, *TMEM67*, *TMEM216*, *TMEM231*) (Kyttälä et al. 2006; Smith et al. 2006; Baala et al. 2007a; Delous et al. 2007; Bergmann et al. 2008; Tallila et al. 2008; Valente et al. 2010; Dowdle et al. 2011; Hopp et al. 2011; Shaheen et al. 2011, 2013; Filges et al. 2014). There is a considerable genetic overlap between JBS and MKS, in which eight of the genes are shared between the two syndromes and co-occurrence of the two diseases was reported in the same families (Valente et al. 2014). The incidence of MKS varies among populations and it has been estimated as 1/13,250 in the United

Bardet–Biedl syndrome (BBS) is an autosomal recessive condition defined by rod–cone degeneration, postaxial polydactyly, central obesity, mental retardation, hypogonadism, and renal dysfunction (Beales et al. 1999; Mockel et al. 2011). Other features such as hepatic fibrosis, diabetes mellitus, endocrinological disturbances, heart disease, and short stature may also be present (Beales et al. 1999). Twenty genes have been associated with BBS (*ARL6*, *BBIP1*, *BBS1*, *BBS2*, *BBS4*, *BBS5*, *BBS7*, *BBS9*, *BBS10*, *BBS12*, *CEP290*, *IFT27*, *IFT172*, *LZTFL1*, *MKKS*, *MKSI*, *SDCCAG8*, *TRIM32*, *TTC8*, *WDPCP*) (Katsanis et al. 2000; Slavotinek et al. 2000; Mykytyn et al. 2001, 2002; Nishimura et al. 2001, 2005; Ansley et al. 2003; Badano et al. 2003a; Chiang et al. 2004, 2006;

Fan et al. 2004; Li et al. 2004; Stoetzel et al. 2006, 2007; Leitch et al. 2008; Kim et al. 2010; Otto et al. 2010; Marion et al. 2012; Aldahmesh et al. 2014; Bujakowska et al. 2014; Scheidecker et al. 2014). Apart from the BBS genes associated with the nonsyndromic IRD (Table 1), a homozygous nonsense mutation (p.S701X) in *BBS12* was shown to lead to a late-onset retinal degeneration and postaxial polydactyly but no other BBS-associated clinical features (Pawlik et al. 2010).

There are two phenotypically similar diseases to BBS: Alstrom syndrome (ALMS) and MORM syndrome (mental retardation, truncal obesity, retinal degeneration, and micropenis). ALMS is characterized by cone-rod degeneration, sensorineural hearing loss, childhood obesity, and type 2 diabetes mellitus. ALMS patients often present with cardiomyopathy and other features such as renal, pulmonary, or hepatic disease may also be present. In contrast to BBS, ALMS is not associated with mental retardation, polydactyly, or hypogonadism (Mockel et al. 2011). Mutations in only one gene, *ALMS1*, have been associated with Alstrom syndrome (Collin et al. 2002). MORM has been described in only one Pakistani family of 14 individuals, in whom a homozygous truncating mutation in *INPP5E*, was found to cause the disease (Hampshire et al. 2006; Jacoby et al. 2009).

Short-rib thoracic dysplasia (SRTD) with or without polydactyly regroups syndromes formerly known as Mainzer-Saldino (MZSDS), Jeune asphyxiating thoracic dystrophy (JATD), and Ellis-van Creveld (EVC) syndromes. SRTDs are autosomal recessive skeletal ciliopathies, characterized by short ribs, constricted thoracic cage, shortened tubular bones, and a “trident” appearance of the acetabular roof (Huber and Cormier-Daire 2012). The severely constricted thoracic cage leads to respiratory insufficiency, often resulting in death in infancy. Other features that may be present are polydactyly, cleft lip/palate, retinal degeneration, and anomalies of the brain, heart, kidneys, liver, pancreas, intestines, and genitalia (Waters and Beales 2011; Huber and Cormier-Daire 2012). There is a phenotypic and genetic overlap be-

tween SRTDs and cranioectodermal dysplasia (CED), also known as Sensenbrenner syndrome. CED is characterized by sagittal craniosynostosis, narrow thorax, short limbs, brachydactyly, protuberant abdomen, and facial and ectodermal anomalies (Huber and Cormier-Daire 2012). Seventeen genes have been associated with these diseases (*CEP120*, *DYNC2H1*, *EVC*, *EVC2*, *IFT52*, *IFT122*, *IFT140*, *IFT172*, *IFT43*, *IFT80*, *KIAA0586*, *NEK1*, *TTC21B*, *WDR19*, *WDR34*, *WDR35*, *WDR60*) (Ruiz-Perez et al. 2000; Galdzicka et al. 2002; Beales et al. 2007; Dagoneau et al. 2009; Gilissen et al. 2010; Walczak-Sztulpa et al. 2010; Arts et al. 2011; Bredrup et al. 2011; Davis et al. 2011; Mill et al. 2011; Thiel et al. 2011; Perrault et al. 2012; Halbritter et al. 2013; McInerney-Leo et al. 2013; Schmidts et al. 2013b; Alby et al. 2015; Shaheen et al. 2015; Girisha et al. 2016). Interestingly, some of these genes have also been implicated with a nonsyndromic disease, for example, *TTC21B* and *WDR19* in NPHP (Bredrup et al. 2011; Davis et al. 2011) or *IFT172* and *IFT140* in RP (Fig. 3) (Bujakowska et al. 2014; Bifari et al. 2015; Xu et al. 2015).

BROAD PHENOTYPIC SPECTRUM OF CILIOPATHIES AND GENETIC MODIFIERS

One of the important aspects of the IRDs is that mutations in the same gene can lead to variable phenotypes (Ferrante et al. 2001; Ansley et al. 2003; Sayer et al. 2006; Perrault et al. 2007; Frank et al. 2008; Leitch et al. 2008; Coene et al. 2009; Riazuddin et al. 2010; Bujakowska et al. 2012; Estrada-Cuzcano et al. 2012a; Webb et al. 2012). In some cases, the severity of disease can be explained by the primary disease-causing mutation. For example, a splice site mutation of a retina-specific exon in *TTC8* leads to a nonsyndromic RP (Riazuddin et al. 2010), whereas the gene is most commonly associated with BBS (Ansley et al. 2003). The position of a mutation may also determine the phenotype as in the case of truncating mutations in *OFD1*. Nonsense mutations downstream from exon 17 lead to an X-linked dominant oral-facial-digital type 1 (OFD1) syndrome, manifesting with malformations of face, oral cavity, and digits

K.M. Bujakowska et al.

in affected females and lethal in males (Ferrante et al. 2001). However, truncating mutations upstream of exon 17 lead to an X-linked recessive JBS (Coene et al. 2009). In addition, hypomorphic alleles can arise by mutations activating cryptic splice sites, which leads to severely reduced levels of wild-type transcripts as in the case of *CEP290* (den Hollander et al. 2006) and *OFD1* (Webb et al. 2012).

In many cases, however, even a precise genetic diagnosis does not yield a clear genotype–phenotype correlation and the severity of disease can vary greatly even between patients with the same genetic cause of disease. Examples of this include family members that share the same 3bp deletion (c.461_463del) in the *PRPH2* gene but show phenotypes varying from RP involving the peripheral retina to macular disease involving only the central retina (Weleber et al. 1993). Similarly, individuals with mutations in the *RP1* gene show variable phenotypes, ranging from near normal to profoundly affected by retinal degeneration (Jacobson et al. 2000; Berson et al. 2001). Several genetic modifiers have already been identified in IRD disease (Table 2), in which extreme examples are cases of digenic inheritance of nonsyndromic IRD (Kajiwar et al. 1994) and BBS (Katsanis et al. 2001) or the rescuing effect of the wild-type *PRPF31* allele in the dominant *PRPF31*-associated disease (McGee et al. 1997; Vithana et al. 2003; Rose et al. 2016). Even though more than a dozen of genetic modifiers of IRD disease severity have been reported, our knowledge about these variants is still limited because the studies were conducted on a limited number of patients (sometimes single families) targeting a small number of genes and functional validation was not always performed (Table 2). In addition, no study has yet shown the validity of the previously reported modifiers and therefore they remain to be scrutinized by future research.

CONCLUSIONS

In summary, mutations in many different genes can cause retinal ciliopathies, reflecting the diversity of protein functions required for normal PSC function. As indicated, it is increasingly

clear that the phenotypes ascribed to specific genetic forms of disease overlap, and thus a revised system of disease definitions that includes the genetic etiology in the disease name would improve our understanding of these disorders, and their description for patients and clinicians. Further, as we have attempted to illustrate, studies of retinal ciliopathies have provided insights into syndromic disorders, and cilia function in general. Given the ubiquitous presence of cilia on mammalian cells, we anticipate that further study of these disorders and their pathogenesis will continue to inform us about cilia function broadly, and to be informed by the results of cilia in other contexts.

REFERENCES

- Abd El-Aziz MM, Barragan I, O'Driscoll CA, Goodstadt L, Prigmore E, Borrego S, Mena M, Pieras JI, El-Ashry ME, Safieh LA, et al. 2008. *EYS*, encoding an ortholog of *Drosophila* spacemaker, is mutated in autosomal recessive retinitis pigmentosa. *Nat Genet* **40**: 1285–1287.
- Abu-Safieh L, Al-Anazi S, Al-Abdi L, Hashem M, Alkuraya H, Alamr M, Sirelkhatim MO, Al-Hassnan Z, Alkuraya B, Mohamed JY, et al. 2012. In search of triallelism in Bardet–Biedl syndrome. *Eur J Hum Genet* **20**: 420–427.
- Ahmed ZM, Riazuddin S, Bernstein SL, Ahmed Z, Khan S, Griffith AJ, Morell RJ, Friedman TB, Wilcox ER. 2001. Mutations of the protocadherin gene *PCDH15* cause Usher syndrome type 1F. *Am J Hum Genet* **69**: 25–34.
- Ahmed ZM, Kjellstrom S, Haywood-Watson RJJ, Bush RA, Hampton LL, Battley JE, Riazuddin S, Frolenkov G, Sieving PA, Friedman TB. 2008. Double homozygous waltzer and Ames waltzer mice provide no evidence of retinal degeneration. *Mol Vis* **14**: 2227–2236.
- Akatori M, Tsunoda K, Miyake Y, Fukuda Y, Ishiura H, Tsuji S, Usui T, Hatase T, Nakamura M, Ohde H, et al. 2010. Dominant mutations in *RP1L1* are responsible for occult macular dystrophy. *Am J Hum Genet* **87**: 424–429.
- Alby C, Piquand K, Huber C, Megarbané A, Ichkou A, Legendre M, Pelluard F, Encha-Ravazi F, Abi-Tayeh G, Bessières B, et al. 2015. Mutations in *KIAA0586* cause lethal ciliopathies ranging from a hydroletharus phenotype to short-rib polydactyly syndrome. *Am J Hum Genet* **97**: 311–318.
- Aldahmesh MA, Safieh LA, Alkuraya H, Al-Rajhi A, Shamseldin H, Hashem M, Alzahrani F, Khan AO, Alqahtani F, Rahbeeni Z, et al. 2009. Molecular characterization of retinitis pigmentosa in Saudi Arabia. *Mol Vis* **15**: 2464–2469.
- Aldahmesh MA, Li Y, Alhashem A, Anazi S, Alkuraya H, Hashem M, Awaji AA, Sogaty S, Alkharashi A, Alzahrani S, et al. 2014. *IFT27*, encoding a small GTPase component of IFT particles, is mutated in a consanguineous

Photoreceptor Cilia and Retinal Ciliopathies

- family with Bardet–Biedl syndrome. *Hum Mol Genet* **23**: 3307–3315.
- Aligianis IA, Forsheew T, Johnson S, Michaelides M, Johnson CA, Trembath RC, Hunt DM, Moore AT, Maher ER. 2002. Mapping of a novel locus for achromatopsia (*ACHM4*) to 1p and identification of a germline mutation in the alpha subunit of cone transducin (*GNAT2*). *J Med Genet* **39**: 656–660.
- Allikmets R, Singh N, Sun H, Shroyer NF, Hutchinson A, Chidambaram A, Gerrard B, Baird L, Stauffer D, Peiffer A, et al. 1997. A photoreceptor cell-specific ATP-binding transporter gene (*ABCR*) is mutated in recessive Starardt macular dystrophy. *Nat Genet* **15**: 236–246.
- Ansley SJ, Badano JL, Blacque OE, Hill J, Hoskins BE, Leitch CC, Kim JC, Ross AJ, Eichers ER, Teslovich TM, et al. 2003. Basal body dysfunction is a likely cause of pleiotropic Bardet–Biedl syndrome. *Nature* **425**: 628–633.
- Arikawa K, Molday LL, Molday RS, Williams DS. 2011. Localization of peripherin/RDS in the disk membranes of cone and rod photoreceptors: Relationship to disk membrane morphogenesis and retinal degeneration. *J Cell Biol* **116**: 659–667.
- Arts HH, Bongers EMHF, Mans DA, van Beersum SEC, Oud MM, Bolat E, Spruijt L, Cornelissen EAM, Schuurs-Hoeijmakers JHM, de Leeuw N, et al. 2011. *C14ORF179* encoding IFT43 is mutated in Sensenbrenner syndrome. *J Med Genet* **48**: 390–395.
- Ayyagari R, Kakuk LE, Coats CL, Bingham EL, Toda Y, Feliuss J, Sieving P. 1999. Bilateral macular atrophy in blue cone monochromacy (BCM) with loss of the locus control region (LCR) and part of the red pigment gene. *Mol Vis* **5**: 13.
- Ayyagari R, Demirci FY, Liu J, Bingham EL, Stringham H, Kakuk LE, Boehnke M, Gorin MB, Richards JE, Sieving PA. 2002. X-linked recessive atrophic macular degeneration from *RPGR* mutation. *Genomics* **80**: 166–171.
- Baala L, Audollent S, Martinovic J, Ozilou C, Babron M-C, Sivanandamoorthy S, Saunier S, Salomon R, Gonzales M, Rattenberry E, et al. 2007a. Pleiotropic effects of *CEP290* (*NPHP6*) mutations extend to Meckel syndrome. *Am J Hum Genet* **81**: 170–179.
- Baala L, Romano S, Khaddour R, Saunier S, Smith UM, Audollent S, Ozilou C, Faivre L, Laurent N, Foliguet B, et al. 2007b. The Meckel–Gruber syndrome gene, *MKS3*, is mutated in Joubert syndrome. *Am J Hum Genet* **80**: 186–194.
- Badano JL, Ansley SJ, Leitch CC, Lewis RA, Lupski JR, Katsanis N. 2003a. Identification of a novel Bardet–Biedl syndrome protein, BBS7, that shares structural features with BBS1 and BBS2. *Am J Hum Genet* **72**: 650–658.
- Badano JL, Kim JC, Hoskins BE, Lewis RA, Ansley SJ, Cutler DJ, Castellon C, Beales PL, Leroux MR, Katsanis N. 2003b. Heterozygous mutations in *BBS1*, *BBS2* and *BBS6* have a potential epistatic effect on Bardet–Biedl patients with two mutations at a second BBS locus. *Hum Mol Genet* **12**: 1651–1659.
- Badano JL, Leitch CC, Ansley SJ, May-Simera H, Lawson S, Lewis RA, Beales PL, Dietz HC, Fisher S, Katsanis N. 2006. Dissection of epistasis in oligogenic Bardet–Biedl syndrome. *Nature* **439**: 326–330.
- Bandah-Rozenfeld D, Mizrahi-Meissonnier L, Farhy C, Obolensky A, Chowers I, Pe'er J, Merin S, Ben-Yosef T, Ashery-Padan R, Banin E, et al. 2010. Homozygosity mapping reveals null mutations in *FAM161A* as a cause of autosomal-recessive retinitis pigmentosa. *Am J Hum Genet* **87**: 382–391.
- Banerjee P, Kleyn PW, Knowles JA, Lewis CA, Ross BM, Parano E, Kovats SG, Lee JJ, Penchaszadeh GK, Ott J, et al. 1998. *TULP1* mutation in two extended Dominican kindreds with autosomal recessive retinitis pigmentosa. *Nat Genet* **18**: 177–179.
- Bareil C, Hamel CP, Delague V, Arnaud B, Demaille J, Claustres M. 2001. Segregation of a mutation in *CNGB1* encoding the β -subunit of the rod cGMP-gated channel in a family with autosomal recessive retinitis pigmentosa. *Hum Genet* **108**: 328–334.
- Bascom RA, Manara S, Collins L, Molday RS, Kalnins VI, McInnes RR. 1992. Cloning of the cDNA for a novel photoreceptor membrane protein (*rom-1*) identifies a disk rim protein family implicated in human retinopathies. *Neuron* **8**: 1171–1184.
- Beales PL, Elcioglu N, Woolf AS, Parker D, Flintner FA. 1999. New criteria for improved diagnosis of Bardet–Biedl syndrome: Results of a population survey. *J Med Genet* **36**: 437–446.
- Beales PL, Bland E, Tobin JL, Bacchelli C, Tuysuz B, Hill J, Rix S, Pearson CG, Kai M, Hartley J, et al. 2007. *IFT80*, which encodes a conserved intraflagellar transport protein, is mutated in Jeune asphyxiating thoracic dystrophy. *Nat Genet* **39**: 727–729.
- Berger W, Kloeckener-Gruissem B, Neidhardt J. 2010. The molecular basis of human retinal and vitreoretinal diseases. *Prog Retin Eye Res* **29**: 335–375.
- Bergmann C, Fliegauf M, Brühl NO, Frank V, Olbrich H, Kirschner J, Schermer B, Schmedding I, Kispert A, Kränzl B, et al. 2008. Loss of nephrocystin-3 function can cause embryonic lethality, Meckel–Gruber-like syndrome, situs inversus, and renal-hepatic-pancreatic dysplasia. *Am J Hum Genet* **82**: 959–970.
- Berson EL, Grimsby JL, Adams SM, McGee TL, Sweklo E, Pierce EA, Sandberg MA, Dryja TP. 2001. Clinical features and mutations in patients with dominant retinitis pigmentosa-1 (*RPI*). *Invest Ophthalmol Vis Sci* **42**: 2217–2224.
- Besharse JC, Forestner DM, Defoe DM. 1985. Membrane assembly in retinal photoreceptors. III: Distinct membrane domains of the connecting cilium of developing rods. *J Neurosci* **5**: 1035–1048.
- Bielas SL, Silhavy JL, Brancati F, Kisseleva MV, Al-Gazali L, Sztriha L, Bayoumi RA, Zaki MS, Abdel-Aleem A, Rosti RO, et al. 2009. Mutations in *INPP5E*, encoding inositol polyphosphate-5-phosphatase E, link phosphatidylinositol signaling to the ciliopathies. *Nat Genet* **41**: 1032–1036.
- Bifari IN, Elkhamary SM, Bolz HJ, Khan AO. 2015. The ophthalmic phenotype of *IFT140*-related ciliopathy ranges from isolated to syndromic congenital retinal dystrophy. *Br J Ophthalmol* **6**: 829–833.
- Bizet AA, Becker-Heck A, Ryan R, Weber K, Filhol E, Krug P, Halbritter J, Delous M, Lasbennes M-C, Linghu B, et al. 2015. Mutations in *TRAF3IP1/IFT54* reveal a new role for IFT proteins in microtubule stabilization. *Nat Commun* **6**: 8666.

K.M. Bujakowska et al.

- Bolz H, von Brederlow B, Ramírez A, Bryda EC, Kutsche K, Nothwang HG, Seeliger M, del C-Salcedó Cabrera M, Vila MC, Molina OP, et al. 2001. Mutation of *CDH23*, encoding a new member of the cadherin gene family, causes Usher syndrome type 1D. *Nat Genet* **27**: 108–112.
- Bonnet C, El-Amraoui A. 2012. Usher syndrome (sensori-neural deafness and retinitis pigmentosa): Pathogenesis, molecular diagnosis and therapeutic approaches. *Curr Opin Neurol* **25**: 42–49.
- Bowne SJ, Daiger SP, Malone KA, Heckenlively JR, Kennan A, Humphries P, Hughbanks-Wheaton D, Birch DG, Liu Q, Pierce EA, et al. 2003. Characterization of *RP1L1*, a highly polymorphic paralog of the retinitis pigmentosa 1 (*RP1*) gene. *Mol Vis* **9**: 129–137.
- Bredrup C, Saunier S, Oud MM, Fiskerstrand T, Hoischen A, Brackman D, Leh SM, Midtbo M, Filhol E, Bole-Feysot C, et al. 2011. Ciliopathies with skeletal anomalies and renal insufficiency due to mutations in the IFT-A gene *WDR19*. *Am J Hum Genet* **89**: 634–643.
- Bujakowska K, Audo I, Mohand-Saïd S, Lancelot ME, Antonio A, Germain A, Leveillard T, Letexier M, Saraiva JB, Lonjou C, et al. 2012. *CRB1* mutations in inherited retinal dystrophies. *Hum Mutat* **33**: 306–315.
- Bujakowska KM, Zhang Q, Siemiatkowska AM, Liu Q, Place E, Falk MJ, Consugar M, Lancelot ME, Antonio A, Lonjou C, et al. 2014. Mutations in *IFT172* cause isolated retinal degeneration and Bardet–Biedl syndrome. *Hum Mol Genet* **24**: 230–242.
- Calvert PD, Strissel KJ, Schiesser WE, Pugh EN, Arshavsky VY. 2006. Light-driven translocation of signaling proteins in vertebrate photoreceptors. *Trends Cell Biol* **16**: 560–568.
- Calvert PD, Schiesser WE, Pugh EN. 2010. Diffusion of a soluble protein, photoactivatable GFP, through a sensory cilium. *J Gen Physiol* **135**: 173–196.
- Cantagrel V, Silhavy JL, Bielas SL, Swistun D, Marsh SE, Bertrand JY, Audollent S, Attié-Bitach T, Holden KR, Dobyns WB, et al. 2008. Mutations in the cilia gene *ARL13B* lead to the classical form of Joubert syndrome. *Am J Hum Genet* **83**: 170–179.
- Caridi G, Murer L, Bellantuono R, Sorino P, Caringella DA, Gusmano R, Ghiggeri GM. 1998. Renal–retinal syndromes: Association of retinal anomalies and recessive nephronophthisis in patients with homozygous deletion of the *NPH1* locus. *Am J Kidney Dis* **32**: 1059–1062.
- Chaitin MH, Schneider BG, Hall MO, Papermaster DS. 1984. Actin in the photoreceptor connecting cilium: Immunocytochemical localization to the site of outer segment disk formation. *J Cell Biol* **99**: 239–247.
- Chakarova CF, Papaioannou MG, Khanna H, Lopez I, Wa-seem N, Shah A, Theis T, Friedman J, Maubaret C, Bujakowska K, et al. 2007. Mutations in *TOPORS* cause autosomal dominant retinitis pigmentosa with perivascular retinal pigment epithelium atrophy. *Am J Hum Genet* **81**: 1098–1103.
- Chakarova CF, Khanna H, Shah AZ, Patil SB, Sedmak T, Murga-Zamalloa CA, Papaioannou MG, Nagel-Wolfrum K, Lopez I, Munro P, et al. 2011. *TOPORS*, implicated in retinal degeneration, is a cilia-centrosomal protein. *Hum Mol Genet* **20**: 975–987.
- Chaki M, Airik R, Ghosh AK, Giles RH, Chen R, Slaats GG, Wang H, Hurd TW, Zhou W, Cluckey A, et al. 2012. Exome capture reveals *ZNF423* and *CEP164* mutations, linking renal ciliopathies to DNA damage response signaling. *Cell* **150**: 533–548.
- Chiang AP, Nishimura D, Searby C, Elbedour K, Carmi R, Ferguson AL, Secrist J, Braun T, Casavant T, Stone EM, et al. 2004. Comparative genomic analysis identifies an ADP-ribosylation factor-like gene as the cause of Bardet–Biedl syndrome (BBS3). *Am J Hum Genet* **75**: 475–484.
- Chiang AP, Beck JS, Yen HJ, Tayeh MK, Scheetz TE, Swiderski RE, Nishimura DY, Braun T, Kim KY, Huang J, et al. 2006. Homozygosity mapping with SNP arrays identifies *TRIM32*, an E3 ubiquitin ligase, as a Bardet–Biedl syndrome gene (BBS11). *Proc Natl Acad Sci* **103**: 6287–6292.
- Chung DC, Traboulsi EI. 2009. Leber congenital amaurosis: Clinical correlations with genotypes, gene therapy trials update, and future directions. *J AAPOS* **13**: 587–592.
- Clarke G, Goldberg AF, Vidgen D, Collins L, Ploder L, Schwarz L, Molday LL, Rossant J, Szél A, Molday RS, et al. 2000. Rom-1 is required for rod photoreceptor viability and the regulation of disk morphogenesis. *Nat Genet* **25**: 67–73.
- Coene KLM, Roepman R, Doherty D, Afroze B, Kroes HY, Letteboer SJE, Ngu LH, Budny B, van Wijk E, Gorden NT, et al. 2009. OFD1 is mutated in X-linked Joubert syndrome and interacts with *LCA5*-encoded lebercilin. *Am J Hum Genet* **85**: 465–481.
- Collin GB, Marshall JD, Ikeda A, So WV, Russell-Eggitt I, Maffei P, Beck S, Boerkoel CF, Siculo N, Martin M, et al. 2002. Mutations in *ALMS1* cause obesity, type 2 diabetes and neurosensory degeneration in Alström syndrome. *Nat Genet* **31**: 74–78.
- Collin RWJ, Littink KW, Klevering BJ, van den Born LI, Koenekoop RK, Zonneveld MN, Blokland EAW, Strom TM, Hoyng CB, den Hollander AI, et al. 2008. Identification of a 2 Mb human ortholog of *Drosophila* eyes shut/spacemaker that is mutated in patients with retinitis pigmentosa. *Am J Hum Genet* **83**: 594–603.
- Collin RWJ, Safieh C, Littink KW, Shalev SA, Garzozzi HJ, Rizel L, Abbasi AH, Cremers FPM, den Hollander AI, Klevering BJ, et al. 2010. Mutations in *C2ORF71* cause autosomal-recessive retinitis pigmentosa. *Am J Hum Genet* **86**: 783–788.
- Consugar MB, Navarro-Gomez D, Place EM, Bujakowska KM, Sousa ME, Fonseca-Kelly ZD, Taub DG, Janessian M, Wang DY, Au ED, et al. 2014. Panel-based genetic diagnostic testing for inherited eye diseases is highly accurate and reproducible, and more sensitive for variant detection, than exome sequencing. *Genet Med* **17**: 253–261.
- Coppieters F, Casteels I, Meire F, De Jaegere S, Hooghe S, van Regemortel N, Van Esch H, Matuleviciene A, Nunes L, Meerschaert V, et al. 2010. Genetic screening of LCA in Belgium: Predominance of *CEP290* and identification of potential modifier alleles in *AHI1* of *CEP290*-related phenotypes. *Hum Mutat* **31**: E1709–E1766.
- Coussa RG, Otto EA, Gee HY, Arthurs P, Ren H, Lopez I, Keser V, Fu Q, Faingold R, Khan A, et al. 2013. *WDR19*: An ancient, retrograde, intraflagellar ciliary protein is mutated in autosomal recessive retinitis pigmentosa and in Senior–Løken syndrome. *Clin Genet* **84**: 150–159.

Photoreceptor Cilia and Retinal Ciliopathies

- Cremers FP, van de Pol DJ, van Driel M, den Hollander AI, van Haren FJ, Knoers NV, Tijmes N, Bergen AA, Rohrschneider K, Blankenagel A, et al. 1998. Autosomal recessive retinitis pigmentosa and cone-rod dystrophy caused by splice site mutations in the Stargardt's disease gene *ABCR*. *Hum Mol Genet* **7**: 355–362.
- Dafinger C, Liebau MC, Elsayed SM, Hellenbroich Y, Boltshauser E, Korenke GC, Fabretti F, Janecke AR, Ebermann I, Nürnberg G, et al. 2011. Mutations in *KIF7* link Joubert syndrome with Sonic Hedgehog signaling and microtubule dynamics. *J Clin Invest* **121**: 2662–2667.
- Dagoneau N, Goulet M, Geneviève D, Sznajder Y, Martinovic J, Smithson S, Huber C, Baujat G, Flori E, Tecco L, et al. 2009. *DYNC2H1* mutations cause asphyxiating thoracic dystrophy and short rib-polydactyly syndrome, type III. *Am J Hum Genet* **84**: 706–711.
- Davidson AE, Sergouniotis PI, Mackay DS, Wright GA, Waaseem NH, Michaelides M, Holder GE, Robson AG, Moore AT, Plagnol V, et al. 2013. *RP1L1* variants are associated with a spectrum of inherited retinal diseases including retinitis pigmentosa and occult macular dystrophy. *Hum Mutat* **34**: 506–514.
- Davis EE, Zhang Q, Liu Q, Diplas BH, Davey LM, Hartley J, Stoezel C, Szymanska K, Ramaswami G, Logan CV, et al. 2011. *TTC21B* contributes both causal and modifying alleles across the ciliopathy spectrum. *Nat Genet* **43**: 189–196.
- Dellett M, Sasai N, Nishide K, Becker S, Papadaki V, Astrid Limb G, Moore AT, Kondo T, Ohnuma SI. 2015. Genetic background and light-dependent progression of photoreceptor cell degeneration in *Prominin-1* knockout mice. *Investig Ophthalmol Vis Sci* **56**: 164–176.
- Delous M, Baala L, Salomon R, Laclef C, Vierkotten J, Tory K, Golzio C, Lacoste T, Besse L, Ozilou C, et al. 2007. The ciliary gene *RPGRIP1L* is mutated in cerebello-oculorenal syndrome (Joubert syndrome type B) and Meckel syndrome. *Nat Genet* **39**: 875–881.
- den Hollander AI, Koenekoop RK, Yzer S, Lopez I, Arends ML, Voesenek KEJ, Zonneveld MN, Strom TM, Meitinger T, Brunner HG, et al. 2006. Mutations in the *CEP290* (*NPHP6*) gene are a frequent cause of Leber congenital amaurosis. *Am J Hum Genet* **79**: 556–561.
- den Hollander AI, Koenekoop RK, Mohamed MD, Arts HH, Boldt K, Towns KV, Sedmak T, Beer M, Nagel-Wolfrum K, McKibbin M, et al. 2007. Mutations in *LCA5*, encoding the ciliary protein lebercilin, cause Leber congenital amaurosis. *Nat Genet* **39**: 889–895.
- den Hollander AI, Roepman R, Koenekoop RK, Cremers FPM. 2008. Leber congenital amaurosis: Genes, proteins and disease mechanisms. *Prog Retin Eye Res* **27**: 391–419.
- Deretic D, Williams AH, Ransom N, Morel V, Hargrave PA, Arendt A. 2005. Rhodopsin C terminus, the site of mutations causing retinal disease, regulates trafficking by binding to ADP-ribosylation factor 4 (ARF4). *Proc Natl Acad Sci* **102**: 3301–3306.
- De Robertis E. 1956. Electron microscope observations on the submicroscopic organization of the retinal rods. *J Biophys Biochem Cytol* **2**: 319–330.
- Ding JD, Salinas RY, Arshavsky VY. 2015. Discs of mammalian rod photoreceptors form through the membrane evagination mechanism. *J Cell Biol* **211**: 495–502.
- Di Palma F, Holme RH, Bryda EC, Belyantseva IA, Pellegrino R, Kachar B, Steel KP, Noben-Trauth K. 2001. Mutations in *Cdh23*, encoding a new type of cadherin, cause stereocilia disorganization in waltzer, the mouse model for Usher syndrome type 1D. *Nat Genet* **27**: 103–107.
- Dixon-Salazar T, Silhavy JL, Marsh SE, Louie CM, Scott LC, Gururaj A, Al-Gazali L, Al-Tawari AA, Kayserili H, Sztriha L, et al. 2004. Mutations in the *AHI1* gene, encoding joubertin, cause Joubert syndrome with cortical polymicrogyria. *Am J Hum Genet* **75**: 979–987.
- Dowdle WE, Robinson JF, Kneist A, Sirerol-Piquer MS, Frints SGM, Corbit KC, Zaghoul NA, Zaghoul NA, van Lijnschoten G, Mulders L, et al. 2011. Disruption of a ciliary B9 protein complex causes Meckel syndrome. *Am J Hum Genet* **89**: 94–110.
- Dryja TP, McGee TL, Reichel E, Hahn LB, Cowley GS, Yandell DW, Sandberg MA, Berson EL. 1990. A point mutation of the rhodopsin gene in one form of retinitis pigmentosa. *Nature* **343**: 364–366.
- Dryja TP, Berson EL, Rao VR, Oprian DD. 1993. Heterozygous missense mutation in the rhodopsin gene as a cause of congenital stationary night blindness. *Nat Genet* **4**: 280–283.
- Dryja TP, Finn JT, Peng YW, McGee TL, Berson EL, Yau KW. 1995. Mutations in the gene encoding the α subunit of the rod cGMP-gated channel in autosomal recessive retinitis pigmentosa. *Proc Natl Acad Sci* **92**: 10177–10181.
- Dryja TP, Hahn LB, Reboul T, Arnaud B. 1996. Missense mutation in the gene encoding the α subunit of rod transducin in the Nougaret form of congenital stationary night blindness. *Nat Genet* **13**: 358–360.
- Dryja TP, Adams SM, Grimsby JL, McGee TL, Hong DH, Li T, Andréasson S, Berson EL. 2001. Null *RPGRIP1* alleles in patients with Leber congenital amaurosis. *Am J Hum Genet* **68**: 1295–1298.
- Dvir L, Srour G, Abu-Ras R, Miller B, Shalev S, Ben-Yosef T. 2010. Autosomal-recessive early-onset retinitis pigmentosa caused by a mutation in *PDE6G*, the gene encoding the gamma subunit of rod cGMP phosphodiesterase. *Am J Hum Genet* **87**: 258–264.
- Ebermann I, Phillips JB, Liebau MC, Koenekoop RK, Schermer B, Lopez I, Schäfer E, Roux AF, Dafinger C, Bernd A, et al. 2010. *PDZD7* is a modifier of retinal disease and a contributor to digenic Usher syndrome. *J Clin Invest* **120**: 1812–1823.
- Edvardson S, Shaag A, Zenvirt S, Erlich Y, Hannon GJ, Shanske AL, Gomori JM, Ekstein J, Elpeleg O. 2010. Joubert syndrome 2 (JBTS2) in Ashkenazi Jews is associated with a *TMEM216* mutation. *Am J Hum Genet* **86**: 93–97.
- El Shamieh S, Neuillé M, Terray A, Orhan E, Condroyer C, Démontant V, Michiels C, Antonio A, Boyard F, Lancelot ME, et al. 2014. Whole-exome sequencing identifies *KIZ* as a ciliary gene associated with autosomal-recessive rod-cone dystrophy. *Am J Hum Genet* **94**: 625–633.
- Estrada-Cuzcano A, Koenekoop RK, Coppieters F, Kohl S, Lopez I, Collin RW, De Baere E, Roeleveld D, Marek J, Bernd A, et al. 2011. *IQCB1* mutations in patients with Leber congenital amaurosis. *Invest Ophthalmol Vis Sci* **52**: 834–839.
- Estrada-Cuzcano A, Koenekoop RK, Senechal A, De Baere EBW, de Ravel T, Banfi S, Kohl S, Ayuso C, Sharon D, Hoyng CB, et al. 2012a. *BBS1* mutations in a wide spec-

K.M. Bujakowska et al.

- trum of phenotypes ranging from nonsyndromic retinitis pigmentosa to Bardet–Biedl syndrome. *Arch Ophthalmol* **130**: 1425–1432.
- Estrada-Cuzcano A, Neveling K, Kohl S, Banin E, Rotenstreich Y, Sharon D, Falik-Zaccai TC, Hipp S, Roepman R, Wissinger B, et al. 2012b. Mutations in *C8orf37*, encoding a ciliary protein, are associated with autosomal-recessive retinal dystrophies with early macular involvement. *Am J Hum Genet* **90**: 102–109.
- Estrada-Cuzcano A, Roepman R, Cremers FPM, den Hollander AI, Mans D. 2012c. Non-syndromic retinal ciliopathies: Translating gene discovery into therapy. *Hum Mol Genet* **21**: R111–R124.
- Eudy JD, Weston MD, Yao S, Hoover DM, Rehm HL, MaEdmonds M, Yan D, Ahmad I, Cheng JJ, Ayuso C, et al. 1998. Mutation of a gene encoding a protein with extracellular matrix motifs in Usher syndrome type IIa. *Science* **280**: 1753–1757.
- Fahim AT, Bowne SJ, Sullivan LS, Webb KD, Williams JT, Wheaton DK, Birch DG, Daiger SP. 2011. Allelic heterogeneity and genetic modifier loci contribute to clinical variation in males with X-linked retinitis pigmentosa due to *RPGR* mutations. *PLoS ONE* **6**: e23021.
- Fan Y, Esmail MA, Ansley SJ, Blacque OE, Borojevich K, Ross AJ, Moore SJ, Badano JL, May-Simera H, Compton DS, et al. 2004. Mutations in a member of the Ras superfamily of small GTP-binding proteins causes Bardet–Biedl syndrome. *Nat Genet* **36**: 989–993.
- Farrar GJ, McWilliam P, Bradley DG, Kenna P, Lawler M, Sharp EM, Humphries MM, Eiberg H, Conneally PM, Trofatter JA. 1990. Autosomal dominant retinitis pigmentosa: Linkage to rhodopsin and evidence for genetic heterogeneity. *Genomics* **8**: 35–40.
- Ferland RJ, Eyaad W, Collura RV, Tully LD, Hill RS, Al-Nouri D, Al-Rumayyan A, Topcu M, Gascon G, Bodell A, et al. 2004. Abnormal cerebellar development and axonal decussation due to mutations in *AH11* in Joubert syndrome. *Nat Genet* **36**: 1008–1013.
- Ferrante MI, Giorgio G, Feather SA, Bulfone A, Wright V, Ghiani M, Selicorni A, Gammato L, Scolari F, Woolf AS, et al. 2001. Identification of the gene for oral-facial-digital type I syndrome. *Am J Hum Genet* **68**: 569–576.
- Filges I, Nosova E, Bruder E, Tercanli S, Townsend K, Gibson WT, Röthlisberger B, Heinemann K, Hall JG, Gregory-Evans CY, et al. 2014. Exome sequencing identifies mutations in *KIF14* as a novel cause of an autosomal recessive lethal fetal ciliopathy phenotype. *Clin Genet* **86**: 220–228.
- Fingert JH, Oh K, Chung M, Scheetz TE, Andorf JL, Johnson RM, Sheffield VC, Stone EM. 2008. Association of a novel mutation in the retinol dehydrogenase 12 (*RDH12*) gene with autosomal dominant retinitis pigmentosa. *Arch Ophthalmol* **126**: 1301–1307.
- Fotiadi D, Liang Y, Filipek S, Saperstein DA, Engel A, Palczewski K. 2003. Rhodopsin dimers in native disc membranes. *Nature* **421**: 127–128.
- Frank V, den Hollander AI, Brückle NO, Zonneveld MN, Nürnberg G, Becker C, Du Bois G, Kendziorra H, Roosing S, Senderek J, et al. 2008. Mutations of the *CEP290* gene encoding a centrosomal protein cause Meckel–Gruber syndrome. *Hum Mutat* **29**: 45–52.
- Fuchs S, Nakazawa M, Maw M, Tamai M, Oguchi Y, Gal A. 1995. A homozygous 1-base pair deletion in the arrestin gene is a frequent cause of Oguchi disease in Japanese. *Nat Genet* **10**: 360–362.
- Gal A, Orth U, Baehr W, Schwinger E, Rosenberg T. 1994. Heterozygous missense mutation in the rod cGMP phosphodiesterase beta-subunit gene in autosomal dominant stationary night blindness. *Nat Genet* **7**: 64–68.
- Galdzicka M, Patnala S, Hirshman MG, Cai JF, Nitowsky H, Egeland JA, Ginns EI. 2002. A new gene, *EVC2*, is mutated in Ellis–van Creveld syndrome. *Mol Genet Metab* **77**: 291–295.
- Garcia-Gonzalo FR, Corbit KC, Sirel-Piquer MS, Ramaswami G, Otto EA, Noriega TR, Seol AD, Robinson JF, Bennett CL, Josifova DJ, et al. 2011. A transition zone complex regulates mammalian ciliogenesis and ciliary membrane composition. *Nat Genet* **43**: 776–784.
- Geller SE, Guerin KI, Visel M, Pham A, Lee ES, Dror AA, Avraham KB, Hayashi T, Ray CA, Reh TA, et al. 2009. *CLRN1* is nonessential in the mouse retina but is required for cochlear hair cell development. *PLoS Genet* **5**: 17–19.
- Gilissen C, Arts HH, Hoischen A, Spruijt L, Mans DA, Arts P, van Lier B, Stehouwer M, van Reeuwijk J, Kant SG, et al. 2010. Exome sequencing identifies *WDR35* variants involved in Sensenbrenner syndrome. *Am J Hum Genet* **87**: 418–423.
- Gilliam JC, Chang JT, Sandoval IM, Zhang Y, Li T, Pittler SJ, Chiu W, Wensel TG. 2012. Three-dimensional architecture of the rod sensory cilium and its disruption in retinal neurodegeneration. *Cell* **151**: 1029–1041.
- Girisha KM, Shukla A, Trujillano D, Bhavani GS, Hebbar M, Kadavigere R, Rolfs A. 2016. A homozygous nonsense variant in *IFT52* is associated with a human skeletal ciliopathy. *Clin Genet* doi: 10.1111/cge.12762.
- Goetz SC, Anderson KV. 2010. The primary cilium: A signalling centre during vertebrate development. *Nat Rev Genet* **11**: 331–344.
- Goldberg AFX, Molday RS. 1996. Subunit composition of the peripherin/RDS–rom-1 disk rim complex from rod photoreceptors: Hydrodynamic evidence for a tetrameric quaternary structure. *Biochemistry* **35**: 6144–6149.
- Gorden NT, Arts HH, Parisi MA, Coene KLM, Lettewer SJF, van Beersum SEC, Mans DA, Hikida A, Eckert M, Knutzen D, et al. 2008. *CC2D2A* is mutated in Joubert syndrome and interacts with the ciliopathy-associated basal body protein CEP290. *Am J Hum Genet* **83**: 559–571.
- Guillonnet X, Piriev NI, Danciger M, Kozak CA, Cideciyan AV, Jacobson SG, Farber DB. 1999. A nonsense mutation in a novel gene is associated with retinitis pigmentosa in a family linked to the *RP1* locus. *Hum Mol Genet* **8**: 1541–1546.
- Hagstrom SA, North MA, Nishina PL, Berson EL, Dryja TP. 1998. Recessive mutations in the gene encoding the tubby-like protein *TULP1* in patients with retinitis pigmentosa. *Nat Genet* **18**: 174–176.
- Halbritter J, Bizet AA, Schmidts M, Porath JD, Braun DA, Gee HY, McInerney-Leo AM, Krug P, Filhol E, Davis EE, et al. 2013. Defects in the IFT-B component *IFT172* cause Jeune and Mainzer–Saldino syndromes in humans. *Am J Hum Genet* **93**: 1–11.
- Hameed A, Abid A, Aziz A, Ismail M, Mehdi SQ, Khaliq S. 2003. Evidence of *RPGRIP1* gene mutations associated

- with recessive cone-rod dystrophy. *J Med Genet* **40**: 616–619.
- Hamel C. 2006. Retinitis pigmentosa. *Orphanet J Rare Dis* **1**: 40.
- Hampshire DJ, Ayub M, Springell K, Roberts E, Jafri H, Rashid Y, Bond J, Riley JH, Woods CG. 2006. MORM syndrome (mental retardation, truncal obesity, retinal dystrophy and micropenis), a new autosomal recessive disorder, links to 9q34. *Eur J Hum Genet* **14**: 543–548.
- Han Z, Anderson DW, Papermaster DS. 2012. Prominin-1 localizes to the open rims of outer segment lamellae in *Xenopus laevis* rod and cone photoreceptors. *Investig Ophthalmol Vis Sci* **53**: 361–373.
- Hardcastle AJ, Thiselton DL, Van Maldergem L, Saha BK, Jay M, Plant C, Taylor R, Bird AC, Bhattacharya S. 1999. Mutations in the *RP2* gene cause disease in 10% of families with familial X-linked retinitis pigmentosa assessed in this study. *Am J Hum Genet* **64**: 1210–1215.
- Hartong DT, Berson EL, Dryja TP. 2006. Retinitis pigmentosa. *Lancet* **368**: 1795–1809.
- Helou J, Otto EA, Attanasio M, Allen SJ, Parisi MA, Glass I, Utsch B, Hashmi S, Fazzi E, Omran H, et al. 2007. Mutation analysis of *NPHP6/CEP290* in patients with Joubert syndrome and Senior–Løken syndrome. *J Med Genet* **44**: 657–663.
- Henderson RH, Li Z, Abd El Aziz MM, Mackay DS, Eljinini MA, Zeidan M, Moore AT, Bhattacharya SS, Webster AR. 2010. Biallelic mutation of protocadherin-21 (*PCDH21*) causes retinal degeneration in humans. *Mol Vis* **16**: 46–52.
- Heon E, Kim G, Qin S, Garrison JE, Tavares E, Vincent A, Nuangchamnong N, Scott CA, Slusarski DC, Sheffield VC. 2016. Mutations in *C8ORF37* cause Bardet–Biedl syndrome (BBS21). *Hum Mol Genet* doi: 10.1093/hmg/ddw096.
- Hopp K, Heyer CM, Hommerding CJ, Henke SA, Sundsbak JL, Patel S, Patel B, Consugar MB, Czarnecki PG, Gliem TJ, et al. 2011. *B9D1* is revealed as a novel Meckel syndrome (MKS) gene by targeted exon-enriched next-generation sequencing and deletion analysis. *Hum Mol Genet* **20**: 2524–2534.
- Horst CJ, Johnson IV, Besharse JC. 1990. Transmembrane assemblage of the photoreceptor connecting cilium and motile cilium transition zone contain a common immunologic epitope. *Cell Motil Cytoskeleton* **17**: 329–344.
- Huang SHH, Pittler SJJ, Huang X, Oliveira L, Berson ELL, Dryja TPP. 1995. Autosomal recessive retinitis pigmentosa caused by mutations in the α subunit of rod cGMP phosphodiesterase. *Nat Genet* **11**: 468–471.
- Huang L, Szymanska K, Jensen VL, Janecke AR, Innes AM, Davis EE, Frosk P, Li C, Willer JR, Chodirker BN, et al. 2011. *TMEM237* is mutated in individuals with a Joubert syndrome related disorder and expands the role of the TMEM family at the ciliary transition zone. *Am J Hum Genet* **89**: 713–730.
- Huber C, Cormier-Daire V. 2012. Ciliary disorder of the skeleton. *Am J Med Genet Semin Med Genet* **160C**: 165–174.
- Jacobson SG, Cideciyan AV, Iannaccone A, Weleber RG, Fishman GA, Maguire AM, Affatigato LM, Bennett J, Pierce EA, Danciger M, et al. 2000. Disease expression of *RP1* mutations causing autosomal dominant retinitis pigmentosa. *Invest Ophthalmol Vis Sci* **41**: 1898–1908.
- Jacoby M, Cox JJ, Gayral S, Hampshire DJ, Ayub M, Blockmans M, Pernot E, Kisseleva MV, Compère P, Schiffmann SN, et al. 2009. *INPP5E* mutations cause primary cilium signaling defects, ciliary instability and ciliopathies in human and mouse. *Nat Genet* **41**: 1027–1031.
- Janecke AR, Thompson DA, Utermann G, Becker C, Hubner CA, Schmid E, McHenry CL, Nair AR, Ruschendorf F, Heckenlively J, et al. 2004. Mutations in *RDH12* encoding a photoreceptor cell retinol dehydrogenase cause childhood-onset severe retinal dystrophy. *Nat Genet* **36**: 850–854.
- Joensuu T, Hämäläinen R, Yuan B, Johnson C, Tegelberg S, Gasparini P, Zelante L, Pirvola U, Pakarinen L, Lehesjoki AE, et al. 2001. Mutations in a novel gene with transmembrane domains underlie Usher syndrome type 3. *Am J Hum Genet* **69**: 673–684.
- Johnson KR, Gagnon LH, Webb LS, Peters LL, Hawes NL, Zheng QY, Chang B, Zheng QY. 2003. Mouse models of *USH1C* and *DFNB18*: Phenotypic and molecular analyses of two new spontaneous mutations of the *Ush1c* gene. *Hum Mol Genet* **12**: 3075–3086.
- Kajiwar K, Hahn LB, Mukai S, Travis GH, Berson EL, Dryja TP. 1991. Mutations in the human retinal degeneration slow gene in autosomal dominant retinitis pigmentosa. *Nature* **354**: 480–483.
- Kajiwar K, Berson EL, Dryja TP. 1994. Digenic retinitis pigmentosa due to mutations at the unlinked peripherin/RDS and ROM1 loci. *Science* **264**: 1604–1608.
- Kaplan MW, Iwata RT, Sears RC. 1987. Lengths of immunolabeled ciliary microtubules in frog photoreceptor outer segments. *Exp Eye Res* **44**: 623–632.
- Katsanis N, Beales PL, Woods MO, Lewis RA, Green JS, Parfrey PS, Ansley SJ, Davidson WS, Lupski JR. 2000. Mutations in *MKKS* cause obesity, retinal dystrophy and renal malformations associated with Bardet–Biedl syndrome. *Nat Genet* **26**: 67–70.
- Katsanis N, Ansley SJ, Badano JL, Eichers ER, Lewis RA, Hoskins BE, Scambler PJ, Davidson WS, Beales PL, Lupski JR. 2001. Triallelic inheritance in Bardet–Biedl syndrome, a Mendelian recessive disorder. *Science* **293**: 2256–2259.
- Kellsell RE, Gregory-Evans K, Payne AM, Perrault I, Kaplan J, Yang RB, Garbers DL, Bird AC, Moore AT, Hunt DM. 1998. Mutations in the retinal guanylate cyclase (*RETGC-1*) gene in dominant cone–rod dystrophy. *Hum Mol Genet* **7**: 1179–1184.
- Kennedy B, Malicki J. 2009. What drives cell morphogenesis—A look inside the vertebrate photoreceptor. *Dev Dyn* **238**: 2115–2138.
- Kevany BM, Zhang N, Jastrzebska B, Palczewski K. 2015. Animals deficient in *C2Orf71*, an autosomal recessive retinitis pigmentosa-associated locus, develop severe early-onset retinal degeneration. *Hum Mol Genet* **24**: 2627–2640.
- Khan MI, Kersten FJ, Azam M, Collin RWJ, Hussain A, Shah STA, Keunen JEE, Kremer H, Cremers FPM, Qamar R, et al. 2011. *CLRN1* mutations cause nonsyndromic retinitis pigmentosa. *Ophthalmology* **118**: 1444–1448.
- Khan AO, Decker E, Bachmann N, Bolz HJ, Bergmann C. 2016. *C8orf37* is mutated in Bardet–Biedl syndrome and

K.M. Bujakowska et al.

- constitutes a locus allelic to non-syndromic retinal dystrophies. *Ophthalmic Genet* **8**: 1–4.
- Khanna H. 2015. Photoreceptor sensory cilium: Traversing the ciliary gate. *Cells* **4**: 674–686.
- Khanna H, Davis EE, Murga-Zamalloa CA, Estrada-Cuzcano A, Lopez I, den Hollander AI, Zonneveld MN, Othman MI, Waseem N, Chakarova CF, et al. 2009. A common allele in RPGRIP1L is a modifier of retinal degeneration in ciliopathies. *Nat Genet* **41**: 739–745.
- Khateb S, Zelinger L, Mizrahi-Meissonnier L, Ayuso C, Koenekoop RK, Laxer U, Gross M, Banin E, Sharon D. 2014. A homozygous nonsense *CEP250* mutation combined with a heterozygous nonsense *C2orf71* mutation is associated with atypical Usher syndrome. *J Med Genet* **51**: 460–469.
- Kim SK, Shindo A, Park TJ, Oh EC, Ghosh S, Gray RS, Lewis RA, Johnson CA, Attie-Bittach T, Katsanis N, et al. 2010. Planar cell polarity acts through septins to control collective cell movement and ciliogenesis. *Science* **329**: 1337–40.
- Kim MS, Pinto SM, Getnet D, Nirujogi RS, Manda SS, Chaerkady R, Madugundu AK, Kelkar DS, Isserlin R, Jain S, et al. 2014. A draft map of the human proteome. *Nature* **509**: 575–581.
- Koenekoop RK. 2004. An overview of leber congenital amaurosis: A model to understand human retinal development. *Surv Ophthalmol* **49**: 379–398.
- Kohl S, Marx T, Giddings I, Jägle H, Jacobson SG, Apfelstedt-Sylla E, Zrenner E, Sharpe LT, Wissinger B. 1998. Total colourblindness is caused by mutations in the gene encoding the α -subunit of the cone photoreceptor cGMP-gated cation channel. *Nat Genet* **19**: 257–259.
- Kohl S, Baumann B, Broghammer M, Jägle H, Sieving P, Kellner U, Spegal R, Anastasi M, Zrenner E, Sharpe LT, et al. 2000. Mutations in the *CNGB3* gene encoding the β -subunit of the cone photoreceptor cGMP-gated channel are responsible for achromatopsia (ACHM3) linked to chromosome 8q21. *Hum Mol Genet* **9**: 2107–2116.
- Kohl S, Baumann B, Rosenberg T, Kellner U, Lorenz B, Vadalà M, Jacobson SG, Wissinger B. 2002. Mutations in the cone photoreceptor G-protein α -subunit gene *GNAT2* in patients with achromatopsia. *Am J Hum Genet* **71**: 422–425.
- Kremer H, van Wijk E, Märker T, Wolfrum U, Roepman R. 2006. Usher syndrome: Molecular links of pathogenesis, proteins and pathways. *Hum Mol Genet* **15**: R262–R270.
- Kyttälä M, Tallila J, Salonen R, Kopra O, Kohlschmidt N, Paavola-Sakki P, Peltonen L, Kestilä M. 2006. MKS1, encoding a component of the flagellar apparatus basal body proteome, is mutated in Meckel syndrome. *Nat Genet* **38**: 155–157.
- Lamb TD, Pugh EN. 2006. Phototransduction, dark adaptation, and rhodopsin regeneration the proctor lecture. *Invest Ophthalmol Vis Sci* **47**: 5137–5152.
- Lamb TD, Collin SP, Pugh EN. 2007. Evolution of the vertebrate eye: Opsins, photoreceptors, retina and eye cup. *Nat Rev Neurosci* **8**: 960–976.
- Langmann T, Di Gioia SA, Rau I, Stöhr H, Maksimovic NS, Corbo JC, Renner AB, Zrenner E, Kumaramanickavel G, Karlstetter M, et al. 2010. Nonsense mutations in *FAM161A* cause RP28-associated recessive retinitis pigmentosa. *Am J Hum Genet* **87**: 376–381.
- Larsson NG, Wang J, Wilhelmsson H, Oldfors A, Rustin P, Lewandoski M, Barsh GS, Clayton DA. 1998. Recessive mutations in *TULP1* in RP. *Nat Genet* **18**: 231–236.
- Lee JE, Silhavy JL, Zaki MS, Schroth J, Bielas SL, Marsh SE, Olvera J, Brancati F, Iannicelli M, Ikegami K, et al. 2012a. *CEP41* is mutated in Joubert syndrome and is required for tubulin glutamylation at the cilium. *Nat Genet* **44**: 193–199.
- Lee JH, Silhavy JL, Lee JE, Al-Gazali L, Thomas S, Davis EE, Bielas SL, Hill KJ, Iannicelli M, Brancati F, et al. 2012b. Evolutionarily assembled *cis*-regulatory module at a human ciliopathy locus. *Science* **335**: 966–969.
- Lehman AM, Eyedoux P, Doherty D, Glass IA, Chitayat D, Chung BYH, Langlois S, Yong SL, Lowry RB, Hildebrandt F, et al. 2010. Co-occurrence of Joubert syndrome and Jeune asphyxiating thoracic dystrophy. *Am J Med Genet Part A* **152**: 1411–1419.
- Leitch CC, Zaghloul NA, Davis EE, Stoetzel C, Diaz-Font A, Rix S, Alfadhel M, Al-Fadhel M, Lewis RA, Eyaid W, et al. 2008. Hypomorphic mutations in syndromic encephalocele genes are associated with Bardet–Biedl syndrome. *Nat Genet* **40**: 443–448.
- Li JB, Gerdes JM, Haycraft CJ, Fan Y, Teslovich TM, May-Simera H, Li H, Blacque OE, Li L, Leitch CC, et al. 2004. Comparative genomics identifies a flagellar and basal body proteome that includes the *BBS5* human disease gene. *Cell* **117**: 541–552.
- Libby RT, Steel KP. 2001. Electroretinographic anomalies in mice with mutations in *Myo7a*, the gene involved in human Usher syndrome type 1B. *Invest Ophthalmol Vis Sci* **42**: 770–778.
- Lindstrand A, Davis EE, Carvalho CMB, Pehlivan D, Willer JR, Tsai IC, Ramanathan S, Zuppan C, Sabo A, Muzny D, et al. 2014. Recurrent CNVs and SNVs at the *NPHP1* locus contribute pathogenic alleles to Bardet–Biedl syndrome. *Am J Hum Genet* **94**: 745–754.
- Liu X, Udovichenko IP, Brown SD, Steel KP, Williams DS. 1999. Myosin VIIa participates in opsin transport through the photoreceptor cilium. *J Neurosci* **19**: 6267–6274.
- Liu Q, Tan G, Levenkova N, Li T, Pugh EN, Rux JJ, Speicher DW, Pierce EA. 2007a. The proteome of the mouse photoreceptor sensory cilium complex. *Mol Cell Proteomics* **6**: 1299–1317.
- Liu X, Bulgakov OV, Darrow KN, Pawlyk B, Adamian M, Liberman MC, Li T. 2007b. Usherin is required for maintenance of retinal photoreceptors and normal development of cochlear hair cells. *Proc Natl Acad Sci* **104**: 4413–4418.
- Liu Q, Zhang Q, Pierce EA. 2010. Photoreceptor sensory cilia and inherited retinal degeneration. *Adv Exp Med Biol* **664**: 223–232.
- Logan CV, Abdel-Hamed Z, Johnson CA. 2011. Molecular genetics and pathogenic mechanisms for the severe ciliopathies: Insights into neurodevelopment and pathogenesis of neural tube defects. *Mol Neurobiol* **43**: 12–26.
- Løken AC, Hanssen O, Halvorsen S, Jolster NJ. 1961. Hereditary renal dysplasia and blindness. *Acta Paediatr* **50**: 177–184.
- Louie CM, Caridi G, Lopes VS, Brancati F, Kispert A, Lancaster MA, Schlossman AM, Otto EA, Leitges M, Gröne HJ, et al. 2010. *AHI1* is required for photoreceptor outer

- segment development and is a modifier for retinal degeneration in nephronophthisis. *Nat Genet* **42**: 175–180.
- Maria BL, Hoang KB, Tusa RJ, Mancuso AA, Hamed LM, Quisling RG, Hove MT, Fennell EB, Booth-Jones M, Ringdahl DM, et al. 1997. “Joubert syndrome” revisited: Key ocular motor signs with magnetic resonance imaging correlation. *J Child Neurol* **12**: 423–430.
- Marion V, Stutzmann F, Gérard M, De Melo C, Schaefer E, Claussmann A, Hellé S, Delague V, Souied E, Barrey C, et al. 2012. Exome sequencing identifies mutations in *LZTFL1*, a BBSome and smoothed trafficking regulator, in a family with Bardet–Biedl syndrome with situs inversus and insertional polydactyly. *J Med Genet* **49**: 317–321.
- Martínez-Mir A, Paloma E, Allikmets R, Ayuso C, del Río T, Dean M, Vilageliu L, González-Duarte R, Balcells S. 1998. Retinitis pigmentosa caused by a homozygous mutation in the Stargardt disease gene *ABCR*. *Nat Genet* **18**: 11–12.
- Maw MA, Corbeil D, Koch J, Hellwig A, Wilson-Wheeler JC, Bridges RJ, Kumaramanickavel G, John S, Nancarrow D, Röper K, et al. 2000. A frameshift mutation in prominin (mouse)-like 1 causes human retinal degeneration. *Hum Mol Genet* **9**: 27–34.
- Mburu P, Mustapha M, Varela A, Weil D, El-Amraoui A, Holme RH, Rump A, Hardisty RE, Blanchard S, Coimbra RS, et al. 2003. Defects in whirlin, a PDZ domain molecule involved in stereocilia elongation, cause deafness in the whirler mouse and families with DFNB31. *Nat Genet* **34**: 421–428.
- McGee TL, Devoto M, Ott J, Berson EL, Dryja TP. 1997. Evidence that the penetrance of mutations at the RP11 locus causing dominant retinitis pigmentosa is influenced by a gene linked to the homologous RP11 allele. *Am J Hum Genet* **61**: 1059–1066.
- McInerney-Leo AM, Schmidts M, Cortés CR, Leo PJ, Gener B, Courtney AD, Gardiner B, Harris JA, Lu Y, Marshall M, et al. 2013. Short-rib polydactyly and Jeune syndromes are caused by mutations in *WDR60*. *Am J Hum Genet* **93**: 515–523.
- McLaughlin MEE, Sandberg MAA, Berson ELL, Dryja TPP. 1993. Recessive mutations in the gene encoding the β -subunit of rod phosphodiesterase in patients with retinitis pigmentosa. *Nat Genet* **4**: 130–134.
- Mears AJ, Gieser L, Yan D, Chen C, Fahrner S, Hiriyan S, Fujita R, Jacobson SG, Sieving PA, Swaroop A. 1999. Protein-truncation mutations in the *RP2* gene in a North American cohort of families with X-linked retinitis pigmentosa. *Am J Hum Genet* **64**: 897–900.
- Meindl A, Dry K, Herrmann K, Manson F, Ciccodicola A, Edgar A, Carvalho MR, Achatz H, Hellebrand H, Lennon A, et al. 1996. A gene (*RPGR*) with homology to the *RCC1* guanine nucleotide exchange factor is mutated in X-linked retinitis pigmentosa (RP3). *Nat Genet* **13**: 35–42.
- Michaelides M, Hardcastle AJ, Hunt DM, Moore AT. 2006. Progressive cone and cone–rod dystrophies: Phenotypes and underlying molecular genetic basis. *Surv Ophthalmol* **51**: 232–258.
- Mill P, Lockhart PJ, Fitzpatrick E, Mountford HS, Hall EA, Reijns MAM, Keighren M, Bahlo M, Bromhead CJ, Budd B, et al. 2011. Human and mouse mutations in *WDR35* cause short-rib polydactyly syndromes due to abnormal ciliogenesis. *Am J Hum Genet* **88**: 508–515.
- Millán JM, Aller E, Jaijo T, Blanco-Kelly F, Gimenez-Pardo A, Ayuso C. 2011. An update on the genetics of Usher syndrome. *J Ophthalmol* **2011**: 417217.
- Miyasaka Y, Suzuki S, Ohshiba Y, Watanabe K, Sagara Y, Yasuda SP, Matsuoka K, Shitara H, Yonekawa H, Komina R, et al. 2013. Compound heterozygosity of the functionally null *Cdh23*^{v-ngt} and hypomorphic *Cdh23*^{ah1} alleles leads to early-onset progressive hearing loss in mice. *Exp Anim* **62**: 333–346.
- Mockel A, Perdomo Y, Stutzmann F, Letsch J, Marion V, Dollfus H. 2011. Retinal dystrophy in Bardet–Biedl syndrome and related syndromic ciliopathies. *Prog Retin Eye Res* **30**: 258–274.
- Molday RS, Molday LL. 1987. Differences in the protein composition of bovine retinal rod outer segment disk and plasma membranes isolated by a ricin–gold–dextran density perturbation method. *J Cell Biol* **105**: 2589–2601.
- Molday RS, Moritz OL. 2015. Photoreceptors at a glance. *J Cell Sci* **128**: 4039–4045.
- Molday RS, Hicks D, Molday L. 1987. Peripherin: A rim specific membrane protein of rod outer segment discs. *Investig Ophthalmology Vis Sci* **28**: 50–61.
- Myktyyn K, Braun T, Carmi R, Haider NB, Searby CC, Shastri M, Beck G, Wright AE, Iannaccone A, Elbedour K, et al. 2001. Identification of the gene that, when mutated, causes the human obesity syndrome BBS4. *Nat Genet* **28**: 188–191.
- Myktyyn K, Nishimura DY, Searby CC, Shastri M, Yen H, Beck JS, Braun T, Streb LM, Cornier AS, Cox GF, et al. 2002. Identification of the gene (*BBS1*) most commonly involved in Bardet–Biedl syndrome, a complex human obesity syndrome. *Nat Genet* **31**: 435–438.
- Nair KS, Hanson SM, Mendez A, Gurevich EV, Kennedy MJ, Shestopalov VI, Vishnivetskiy SA, Chen J, Hurley JB, Gurevich VV, et al. 2005. Light-dependent redistribution of arrestin in vertebrate rods is an energy-independent process governed by protein–protein interactions. *Neuron* **46**: 555–567.
- Najafi M, Calvert PD. 2012. Transport and localization of signaling proteins in ciliated cells. *Vision Res* **75**: 11–18.
- Nakazawa M, Wada Y, Tamai M. 1998. Arrestin gene mutations in autosomal recessive retinitis pigmentosa. *Arch Ophthalmol* **116**: 498–501.
- Nathans J, Thomas D, Hogness DS. 1986. Molecular genetics of human color vision: The genes encoding blue, green, and red pigments. *Science* **232**: 193–202.
- Nathans J, Merbs S, Sung CH, Weitz C, Wang Y. 1992. Molecular genetics of human visual pigments. *Annu Rev Genet* **26**: 403–424.
- Nickell S, Park PSH, Baumeister W, Palczewski K. 2007. Three-dimensional architecture of murine rod outer segments determined by cryoelectron tomography. *J Cell Biol* **177**: 917–925.
- Nikopoulos K, Farinelli P, Royer-Bertrand B, Bedoni N, Kjellström U, Andreasson S, Tsilimbaris MK, Tsika C, Blazaki S, Rivolta C. 2016. Whole exome sequencing reveals CEP78 as a novel disease gene for cone-rod dys-

K.M. Bujakowska et al.

- trophy. *ARVO Annual Meeting*, Abstract 1423. Seattle, WA, May 1–6.
- Nishiguchi KM, Sandberg MA, Kooijman AC, Martemyanov KA, Pott JWR, Hagstrom SA, Arshavsky VY, Berson EL, Dryja TP. 2004. Defects in RGS9 or its anchor protein R9AP in patients with slow photoreceptor deactivation. *Nature* **427**: 75–78.
- Nishiguchi KM, Tearle RG, Liu YP, Oh EC, Miyake N, Benaglio P, Harper S, Koskiniemi-Kuendig H, Venturini G, Sharon D, et al. 2013. Whole genome sequencing in patients with retinitis pigmentosa reveals pathogenic DNA structural changes and *NEK2* as a new disease gene. *Proc Natl Acad Sci* **110**: 16139–16144.
- Nishimura DY, Searby CC, Carmi R, Elbedour K, Van Maldergem L, Fulton AB, Lam BL, Powell BR, Swiderski RE, Bugge KE, et al. 2001. Positional cloning of a novel gene on chromosome 16q causing Bardet–Biedl syndrome (BBS2). *Hum Mol Genet* **10**: 865–874.
- Nishimura DY, Swiderski RE, Searby CC, Berg EM, Ferguson AL, Hennekam R, Merin S, Weleber RG, Biesecker LG, Stone EM, et al. 2005. Comparative genomics and gene expression analysis identifies *BBS9*, a new Bardet–Biedl syndrome gene. *Am J Hum Genet* **77**: 1021–1033.
- Nishimura DY, Baye LM, Perveen R, Searby CC, Avila-Fernandez A, Pereiro I, Ayuso C, Valverde D, Bishop PN, Manson FDC, et al. 2010. Discovery and functional analysis of a retinitis pigmentosa gene, *C2ORF71*. *Am J Hum Genet* **86**: 686–695.
- Noor A, Windpassinger C, Patel M, Stachowiak B, Mikhailov A, Azam M, Irfan M, Siddiqui ZK, Naeem F, Paterson AD, et al. 2008. CC2D2A, encoding a coiled-coil and C2 domain protein, causes autosomal-recessive mental retardation with retinitis pigmentosa. *Am J Hum Genet* **82**: 1011–1018.
- Ong ACM, Harris PC. 2015. A polycystin-centric view of cyst formation and disease: The polycystins revisited. *Kidney Int* **88**: 699–710.
- Ostergaard E, Batbayli M, Duno M, Vilhelmsen K, Rosenberg T. 2010. Mutations in *PCDH21* cause autosomal recessive cone–rod dystrophy. *J Med Genet* **47**: 665–669.
- Otto E, Hoefele J, Ruf R, Mueller AM, Hiller KS, Wolf MTE, Schuermann MJ, Becker A, Birkenhäger R, Sudbrak R, et al. 2002. A gene mutated in nephronophthisis and retinitis pigmentosa encodes a novel protein, nephroretinin, conserved in evolution. *Am J Hum Genet* **71**: 1161–1117.
- Otto EA, Loeys B, Khanna H, Hellemans J, Sudbrak R, Fan S, Muerb U, O’Toole JF, Helou J, Attanasio M, et al. 2005. Nephrocystin-5, a ciliary IQ domain protein, is mutated in Senior–Løken syndrome and interacts with RPGR and calmodulin. *Nat Genet* **37**: 282–288.
- Otto EA, Hurd TW, Airik R, Chaki M, Zhou W, Stoetzel C, Patil SB, Levy S, Ghosh AK, Murga-Zamalloa CA, et al. 2010. Candidate exome capture identifies mutation of *SDCCAG8* as the cause of a retinal–renal ciliopathy. *Nat Genet* **42**: 840–850.
- Ozgül RK, Siemiatkowska AM, Yücel D, Myers CA, Collin RWJ, Zonneveld MN, Beryozkin A, Banin E, Hoyng CB, van den Born LJ, et al. 2011. Exome sequencing and cis-regulatory mapping identify mutations in *MAK*, a gene encoding a regulator of ciliary length, as a cause of retinitis pigmentosa. *Am J Hum Genet* **89**: 253–264.
- Pan J, Wang Q, Snell WJ. 2005. Cilium-generated signaling and cilia-related disorders. *Lab Invest* **85**: 452–463.
- Parisi MA, Bennett CL, Eckert ML, Dobyns WB, Gleeson JG, Shaw DWW, McDonald R, Eddy A, Chance PF, Glass IA. 2004. The *NPHP1* gene deletion associated with juvenile nephronophthisis is present in a subset of individuals with Joubert syndrome. *Am J Hum Genet* **75**: 82–91.
- Patel A, Honoré E. 2010. Polycystins and renovascular mechanosensory transduction. *Nat Rev Nephrol* **6**: 530–538.
- Pawlik B, Mir A, Iqbal H, Li Y, Nürnberg G, Becker C, Qamar R, Nürnberg P, Wollnik B. 2010. A novel familial *BBS12* mutation associated with a mild phenotype: Implications for clinical and molecular diagnostic strategies. *Mol Syndromol* **1**: 27–34.
- Payne AM, Downes SM, Bessant DA, Taylor R, Holder GE, Warren MJ, Bird AC, Bhattacharya SS. 1998. A mutation in guanylate cyclase activator 1A (*GUCA1A*) in an autosomal dominant cone dystrophy pedigree mapping to a new locus on chromosome 6p21.1. *Hum Mol Genet* **7**: 273–277.
- Pearring JN, Salinas RY, Baker SA, Arshavsky VY. 2013. Protein sorting, targeting and trafficking in photoreceptor cells. *Prog Retin Eye Res* **36**: 24–51.
- Pedersen LB, Rosenbaum JL. 2008. Intraflagellar transport (IFT) role in ciliary assembly, resorption and signalling. *Curr Top Dev Biol* **85**: 23–61.
- Perrault I, Rozet JM, Calvas P, Gerber S, Camuzat A, Dollfus H, Châtelain S, Souied E, Ghazi I, Leowski C, et al. 1996. Retinal-specific guanylate cyclase gene mutations in Leber’s congenital amaurosis. *Nat Genet* **14**: 461–464.
- Perrault I, Hanein S, Gerber S, Barbet F, Ducrocq D, Dollfus H, Hamel C, Dufier JL, Munnich A, Kaplan J, et al. 2004. Retinal dehydrogenase 12 (*RDH12*) mutations in leber congenital amaurosis. *Am J Hum Genet* **75**: 639–646.
- Perrault I, Delphin N, Hanein S, Gerber S, Dufier JL, Roche O, Defoort-Dhellemmes S, Dollfus H, Fazzi E, Munnich A, et al. 2007. Spectrum of *NPHP6/CEP290* mutations in Leber congenital amaurosis and delineation of the associated phenotype. *Hum Mutat* **28**: 416.
- Perrault I, Saunier S, Hanein S, Filhol E, Bizet AA, Collins F, Salih MAM, Gerber S, Delphin N, Bigot K, et al. 2012. Mainzer–Saldino syndrome is a ciliopathy caused by *IFT140* mutations. *Am J Hum Genet* **90**: 864–870.
- Pierce EA, Quinn T, Meehan T, McGee TL, Berson EL, Dryja TP. 1999. Mutations in a gene encoding a new oxygen-regulated photoreceptor protein cause dominant retinitis pigmentosa. *Nat Genet* **22**: 248–254.
- Poloschek CM, Bach M, Lagrèze WA, Glaus E, Lemke JR, Berger W, Neidhardt J, Lagreze WA, Glaus E, Lemke JR, et al. 2010. *ABCA4* and *ROM1*: Implications for modification of the *PRPH2*-associated macular dystrophy phenotype. *Invest Ophthalmol Vis Sci* **51**: 4253–4265.
- Pretorius PR, Baye LM, Nishimura DY, Searby CC, Bugge K, Yang B, Mullins RF, Stone EM, Sheffield VC, Slusarski DC. 2010. Identification and functional analysis of the vision-specific BBS3 (*ARL6*) long isoform. *PLoS Genet* **6**: e1000884.
- Puffenberger EG, Jinks RN, Sougnéz C, Cibulskis K, Willert RA, Achilly NP, Cassidy RP, Fiorentini CJ, Heiken KF, Lawrence JJ, et al. 2012. Genetic mapping and exome sequencing identify variants associated with five novel diseases. *PLoS ONE* **7**: e28936.

- Pugh EN. 2015. Photoreceptor disc morphogenesis: The classical evagination model prevails. *J Cell Biol* **211**: 2–4.
- Ramamurthy V, Cayouette M. 2009. Development and disease of the photoreceptor cilium. *Clin Genet* **76**: 137–145.
- Rattner A, Smallwood PM, Williams J, Cooke C, Savchenko A, Lyubarsky A, Pugh EN, Nathans J. 2001. A photoreceptor-specific cadherin is essential for the structural integrity of the outer segment and for photoreceptor survival. *Neuron* **32**: 775–786.
- Riazuddin SA, Iqbal M, Wang Y, Masuda T, Chen Y, Bowne S, Sullivan LS, Waseem NH, Bhattacharya S, Daiger SP, et al. 2010. A splice-site mutation in a retina-specific exon of *BBS8* causes nonsyndromic retinitis pigmentosa. *Am J Hum Genet* **86**: 805–812.
- Riazuddin S, Belyantseva IA, Giese APJ, Lee K, Indzhukulian AA, Nandamuri SP, Yousaf R, Sinha GP, Lee S, Terrell D, et al. 2012. Alterations of the CIB2 calcium- and integrin-binding protein cause Usher syndrome type 1J and non-syndromic deafness DFNB48. *Nat Genet* **44**: 1265–1271.
- Rivolta C, Sweklo EA, Berson EL, Dryja TP. 2000. Missense mutation in the *USH2A* gene: Association with recessive retinitis pigmentosa without hearing loss. *Am J Hum Genet* **66**: 1975–1978.
- Roepman R, van Duijnhoven G, Rosenberg T, Pinckers AJ, Bleeker-Wagemakers LM, Bergen AA, Post J, Beck A, Reinhardt R, Ropers HH, et al. 1996. Positional cloning of the gene for X-linked retinitis pigmentosa 3: Homology with the guanine-nucleotide-exchange factor RCC1. *Hum Mol Genet* **5**: 1035–1041.
- Romani M, Micalizzi A, Kraoua I, Dotti MT, Cavallin M, Sztriha L, Ruta R, Mancini F, Mazza T, Castellana S, et al. 2014. Mutations in *B9D1* and *MKSI* cause mild Joubert syndrome: Expanding the genetic overlap with the lethal ciliopathy Meckel syndrome. *Orphanet J Rare Dis* **9**: 72.
- Ronquillo CC, Bernstein PS, Baehr W. 2012. Senior-Løken syndrome: A syndromic form of retinal dystrophy associated with nephronophthisis. *Vision Res* **75**: 88–97.
- Roosing S, Rohrschneider K, Beryozkin A, Sharon D, Weischuh N, Staller J, Kohl S, Zelinger L, Peters TA, Neveling K, et al. 2013. Mutations in *RAB28*, encoding a farnesylated small GTPase, are associated with autosomal-recessive cone–rod dystrophy. *Am J Hum Genet* **93**: 110–117.
- Rose AM, Shah AZ, Venturini G, Krishna A, Chakravarti A, Rivolta C, Bhattacharya SS. 2016. Transcriptional regulation of *PRPF31* gene expression by *MSR1* repeat elements causes incomplete penetrance in retinitis pigmentosa. *Sci Rep* **6**: 19450.
- Rosenbaum JL, Witman GB. 2002. Intraflagellar transport. *Nat Rev Mol Cell Biol* **3**: 813–825.
- Rosenfeld PJ, Cowley GS, McGee TL, Sandberg MA, Berson EL, Dryja TP. 1992. A null mutation in the rhodopsin gene causes rod photoreceptor dysfunction and autosomal recessive retinitis pigmentosa. *Nat Genet* **1**: 209–213.
- Ruiz-Perez VL, Ide SE, Strom TM, Lorenz B, Wilson D, Woods K, King L, Francomano C, Freisinger P, Spranger S, et al. 2000. Mutations in a new gene in Ellis–van Creveld syndrome and Weyers acrodermal dysostosis. *Nat Genet* **24**: 283–286.
- Sahly I, Dufour E, Schietroma C, Michel V, Bahloul A, Perfettini I, Pepermans E, Estivalet A, Carette D, Aghaie A, et al. 2012. Localization of Usher 1 proteins to the photoreceptor calyceal processes, which are absent from mice. *J Cell Biol* **199**: 381–399.
- Sanders AAWM, de Vrieze E, Alazami AM, Alzahrani F, Malarkey EB, Soroush N, Tebbe L, Kuhns S, van Dam TJP, Alhashem A, et al. 2015. KIAA0556 is a novel ciliary basal body component mutated in Joubert syndrome. *Genome Biol* **16**: 293.
- Sanyal S, De Ruiter A, Hawkins R. 1980. Development and degeneration of retina in *rd* mutant mice: Light microscopy. *J Comp Neurol* **194**: 193–207.
- Sato M, Nakazawa M, Usui T, Tanimoto N, Abe H, Ohguro H. 2005. Mutations in the gene coding for guanylate cyclase-activating protein 2 (*GUCA1B* gene) in patients with autosomal dominant retinal dystrophies. *Graefes Arch Clin Exp Ophthalmol* **243**: 235–242.
- Sayer JA, Otto EA, O'Toole JF, Nurnberg G, Kennedy MA, Becker C, Hennies HC, Helou J, Attanasio M, Fausett BV, et al. 2006. The centrosomal protein nephrocystin-6 is mutated in Joubert syndrome and activates transcription factor ATF4. *Nat Genet* **38**: 674–681.
- Scheidecker S, Etard C, Pierce NW, Geoffroy V, Schaefer E, Muller J, Chennen K, Flori E, Pelletier V, Poch O, et al. 2014. Exome sequencing of Bardet–Biedl syndrome patient identifies a null mutation in the BBSome subunit *BBIP1* (*BBS18*). *J Med Genet* **51**: 132–136.
- Schmidts M, Frank V, Eisenberger T, Al Turki S, Bizet AA, Antony D, Rix S, Decker C, Bachmann N, Bald M, et al. 2013a. Combined NGS approaches identify mutations in the intraflagellar transport gene *IFT140* in skeletal ciliopathies with early progressive kidney disease. *Hum Mutat* **34**: 714–724.
- Schmidts M, Vodopiutz J, Christou-Savina S, Cortés CR, McInerney-Leo AM, Emes RD, Arts HH, Tüysüz B, D'Silva J, Leo PJ, et al. 2013b. Mutations in the gene encoding IFT dynein complex component WDR34 cause Jeune asphyxiating thoracic dystrophy. *Am J Hum Genet* **93**: 932–944.
- Senior B, Friedmann AI, Braudo JL. 1961. Juvenile familial nephropathy with tapetoretinal degeneration. A new oculorenal dystrophy. *Am J Ophthalmol* **52**: 625–633.
- Shaheen R, Faqeih E, Seidahmed MZ, Sunker A, Alali FE, AlQahtani K, Alkuraya FS. 2011. A *TCTN2* mutation defines a novel Meckel–Gruber syndrome locus. *Hum Mutat* **32**: 573–578.
- Shaheen R, Ansari S, Mardawi EAL, Alshammari MJ, Alkuraya FS. 2013. Mutations in *TMEM231* cause Meckel–Gruber syndrome. *J Med Genet* **50**: 160–162.
- Shaheen R, Shamseldin HE, Loucks CM, Seidahmed MZ, Ansari S, Ibrahim Khalil M, Al-Yacoub N, Davis EE, Mola NA, Szymanska K, et al. 2014. Mutations in *CSPPI*, encoding a core centrosomal protein, cause a range of ciliopathy phenotypes in humans. *Am J Hum Genet* **94**: 73–79.
- Shaheen R, Schmidts M, Faqeih E, Hashem A, Lausch E, Holder I, Superti-Furga A, Mitchison HM, Almoisheer A, Alamro R, et al. 2015. A founder *CEP120* mutation in Jeune asphyxiating thoracic dystrophy expands the role of centriolar proteins in skeletal ciliopathies. *Hum Mol Genet* **24**: 1410–1419.
- Shankar SP, Hughbanks-Wheaton DK, Birch DG, Sullivan LS, Conneely KN, Bowne SJ, Stone EM, Daiger SP. 2016. Autosomal dominant retinal dystrophies caused by a

K.M. Bujakowska et al.

- founder splice site mutation, c.828+3A>T, in *PRPH2* and protein haplotypes in *trans* as modifiers. *Investig Ophthalmol Vis Sci* **57**: 349.
- Sharon D, Namburi P, Ratnapriya R, Lazar C, Obolensky A, Ben-Yosef T, Pras E, Gross M, Banin E, Swaroop A. 2016. Whole exome sequencing reveals a homozygous splicing mutation in *CEP78* as the cause of atypical Usher syndrome in Eastern Jewish patients. *ARVO Annual Meeting*, Abstract B0253. Seattle, WA, May 1–6.
- Shevach E, Ali M, Mizrahi-Meissonnier L, McKibbin M, El-Asrag M, Watson CM, Inglehearn CF, Ben-Yosef T, Blumenfeld A, Jalas C, et al. 2015. Association between missense mutations in the *BBS2* gene and nonsyndromic retinitis pigmentosa. *JAMA Ophthalmol* **133**: 312.
- Shin JB, Longo-Guess CM, Gagnon LH, Saylor KW, Dumont RA, Spinelli KJ, Pagana JM, Wilmarth PA, David LL, Gillespie PG, et al. 2010. The R109H variant of fascin-2, a developmentally regulated actin crosslinker in hair-cell stereocilia, underlies early-onset hearing loss of DBA/2J mice. *J Neurosci* **30**: 9683–9694.
- Silva E, Dharmaraj S, Li YY, Pina AL, Carter RC, Loyer M, Traboulsi E, Theodossiadi G, Koenekoop R, Sundin O, et al. 2004. A missense mutation in *GUCY2D* acts as a genetic modifier in *RPE65*-related Leber congenital amaurosis. *Ophthalmic Genet* **25**: 205–217.
- Sjostrand FS. 1953. The ultrastructure of the outer segments of rods and cones of the eye as revealed by the electron microscope. *J Cell Physiol* **42**: 15–44.
- Slavotinek AM, Stone EM, Myktyyn K, Heckenlively JR, Green JS, Heon E, Musarella MA, Parfrey PS, Sheffield VC, Biesecker LG. 2000. Mutations in *MKKS* cause Bardet–Biedl syndrome. *Nat Genet* **26**: 15–16.
- Smith UM, Consugar M, Tee LJ, McKee BM, Maina EN, Whelan S, Morgan NV, Goranson E, Gissen P, Lilliquist S, et al. 2006. The transmembrane protein meckelin (*MKS3*) is mutated in Meckel–Gruber syndrome and the *wpk* rat. *Nat Genet* **38**: 191–196.
- Sokal I, Li N, Surgucheva I, Warren MJ, Payne AM, Bhattacharya SS, Baehr W, Palczewski K. 1998. GCAP1 (Y99C) mutant is constitutively active in autosomal dominant cone dystrophy. *Mol Cell* **2**: 129–133.
- Sokolov M, Lyubarsky AL, Strissel KJ, Savchenko AB, Govardovskii VI, Pugh EN, Arshavsky VY. 2002. Massive light-driven translocation of transducin between the two major compartments of rod cells: A novel mechanism of light adaptation. *Neuron* **34**: 95–106.
- Srour M, Hamdan FF, Schwartzentruber JA, Patry L, Ospina LH, Shevell MI, Désilets V, Dobrzyńska S, Mathonnet G, Lemysre E, et al. 2012a. Mutations in *TMEM231* cause Joubert syndrome in French Canadians. *J Med Genet* **49**: 636–641.
- Srour M, Schwartzentruber J, Hamdan FF, Ospina LH, Patry L, Labuda D, Massicotte C, Dobrzyńska S, Capo-Chichi J-M, Papillon-Cavanagh S, et al. 2012b. Mutations in *C5ORF42* cause Joubert syndrome in the French Canadian population. *Am J Hum Genet* **90**: 693–700.
- Steinberg RH, Fisher SK, Anderson DH. 1980. Disc morphogenesis in vertebrate photoreceptors. *J Comp Neurol* **190**: 501–518.
- Stoetzel C, Laurier V, Davis EE, Muller J, Rix S, Badano JL, Leitch CC, Salem N, Chouery E, Corbani S, et al. 2006. *BBS10* encodes a vertebrate-specific chaperonin-like protein and is a major BBS locus. *Nat Genet* **38**: 521–524.
- Stoetzel C, Muller J, Laurier V, Davis EE, Zaghoul NA, Vicaire S, Jacquelin C, Plewniak F, Leitch CC, Sarda P, et al. 2007. Identification of a novel BBS gene (*BBS12*) highlights the major role of a vertebrate-specific branch of chaperonin-related proteins in Bardet–Biedl syndrome. *Am J Hum Genet* **80**: 1–11.
- Stone EM, Cideciyan AV, Aleman TS, Scheetz TE, Sumaroka A, Ehlinger MA, Schwartz SB, Fishman GA, Traboulsi EI, Lam BL, et al. 2011. Variations in *NPHP5* in patients with nonsyndromic leber congenital amaurosis and Senior–Løken syndrome. *Arch Ophthalmol* **129**: 81–87.
- Sun H, Nathans J. 1997. Stargardt's ABCR is localized to the disc membrane of retinal rod outer segments. *Nat Genet* **17**: 15–16.
- Tallila J, Jakkula E, Peltonen L, Salonen R, Kestilä M. 2008. Identification of *CC2D2A* as a Meckel syndrome gene adds an important piece to the ciliopathy puzzle. *Am J Hum Genet* **82**: 1361–1367.
- Taschner M, Bhogaraju S, Lorentzen E. 2012. Architecture and function of IFT complex proteins in ciliogenesis. *Differentiation* **83**: S12–S22.
- Thiadens AAHJ, den Hollander AI, Roosing S, Nabuurs SB, Zekveld-Vroon RC, Collin RWJ, De Baere E, Koenekoop RK, van Schooneveld MJ, Strom TM, et al. 2009. Homozygosity mapping reveals *PDE6C* mutations in patients with early-onset cone photoreceptor disorders. *Am J Hum Genet* **85**: 240–247.
- Thiel C, Kessler K, Giessel A, Dimmler A, Shalev SA, von der Haar S, Zenker M, Zahnleiter D, Stöss H, Beinder E, et al. 2011. *NEK1* mutations cause short-rib polydactyly syndrome type majewski. *Am J Hum Genet* **88**: 106–114.
- Thomas S, Legendre M, Saunier S, Bessières B, Alby C, Bonnière M, Toutain A, Loeuillet L, Szymanska K, Jossic F, et al. 2012. *TCTN3* mutations cause Mohr–Majewski syndrome. *Am J Hum Genet* **91**: 372–378.
- Thomas S, Wright KJ, Le Corre S, Micalizzi A, Romani M, Abhyankar A, Saada J, Perrault I, Amiel J, Litzler J, et al. 2014. A homozygous *PDE6D* mutation in Joubert syndrome impairs targeting of farnesylated INPP5E protein to the primary cilium. *Hum Mutat* **35**: 137–146.
- Travis GH, Sutcliffe JG, Bok D. 1991. The *retinal degeneration slow (rds)* gene product is a photoreceptor disc membrane-associated glycoprotein. *Neuron* **6**: 61–70.
- Tucker BA, Scheetz TE, Mullins RF, DeLuca AP, Hoffmann JM, Johnston RM, Jacobson SG, Sheffield VC, Stone EM. 2011. Exome sequencing and analysis of induced pluripotent stem cells identify the cilia-related gene *male germ cell-associated kinase (MAK)* as a cause of retinitis pigmentosa. *Proc Natl Acad Sci* **108**: E569–E576.
- Tuz K, Bachmann-Gagescu R, O'Day DR, Hua K, Isabella CR, Phelps IG, Stolarski AE, O'Roak BJ, Dempsey JC, Lourenco C, et al. 2014. Mutations in *CSPP1* cause primary cilia abnormalities and Joubert syndrome with or without Jeune asphyxiating thoracic dystrophy. *Am J Hum Genet* **94**: 62–72.
- Valente EM, Silhavy JL, Brancati F, Barrano G, Krishnaswami SR, Castori M, Lancaster MA, Boltshauser E, Boccone L, Al-Gazali L, et al. 2006. Mutations in *CEP290*,

Photoreceptor Cilia and Retinal Ciliopathies

- which encodes a centrosomal protein, cause pleiotropic forms of Joubert syndrome. *Nat Genet* **38**: 623–625.
- Valente EM, Logan CV, Mougou-Zerelli S, Lee JH, Silhavy JL, Brancati F, Iannicelli M, Travaglini L, Romani S, Illi B, et al. 2010. Mutations in *TMEM216* perturb ciliogenesis and cause Joubert, Meckel and related syndromes. *Nat Genet* **42**: 619–625.
- Valente EM, Dallapiccola B, Bertini E. 2013. Joubert syndrome and related disorders. *Handb Clin Neurol* **113**: 1879–1888.
- Valente EM, Rosti RO, Gibbs E, Gleeson JG. 2014. Primary cilia in neurodevelopmental disorders. *Nat Rev Neurol* **10**: 27–36.
- Verpy E, Leibovici M, Zwaenepoel I, Liu XZ, Gal A, Salem N, Mansour A, Blanchard S, Kobayashi I, Keats BJ, et al. 2000. A defect in harmonin, a PDZ domain-containing protein expressed in the inner ear sensory hair cells, underlies Usher syndrome type 1C. *Nat Genet* **26**: 51–55.
- Vithana EN, Safieh LA, Pelosini L, Winchester E, Hornan D, Bird AC, Hunt DM, Bustin SA, Bhattacharya SS. 2003. Expression of *PRPF31* mRNA in patients with autosomal dominant retinitis pigmentosa: A molecular clue for incomplete penetrance? *Invest Ophthalmol Vis Sci* **44**: 4204–4209.
- Wada Y, Abe T, Takeshita T, Sato H, Yanashima K, Tamai M. 2001. Mutation of human retinal fascin gene (*FSCN2*) causes autosomal dominant retinitis pigmentosa. *Invest Ophthalmol Vis Sci* **42**: 2395–2400.
- Wada Y, Abe T, Itabashi T, Sato H, Kawamura M, Tamai M. 2003. Autosomal dominant macular degeneration associated with 208delG mutation in the *FSCN2* gene. *Arch Ophthalmol* **121**: 1613–1620.
- Walczak-Sztulpa J, Eggenschwiler J, Osborn D, Brown DA, Emma F, Klingenberg C, Hennekam RC, Torre G, Garshasbi M, Tzschach A, et al. 2010. Cranioectodermal Dysplasia, Sensenbrenner syndrome, is a ciliopathy caused by mutations in the *IFT122* gene. *Am J Hum Genet* **86**: 949–956.
- Wang J, Deretic D. 2014. Molecular complexes that direct rhodopsin transport to primary cilia. *Prog Retin Eye Res* **38**: 1–19.
- Wang H, den Hollander AI, Moayed Y, Abulimiti A, Li Y, Collin RWJ, Hoyng CB, Lopez I, Abboud EB, Al-Rajhi AA, et al. 2009. Mutations in *SPATA7* cause Leber congenital amaurosis and juvenile retinitis pigmentosa. *Am J Hum Genet* **84**: 380–387.
- Waters AM, Beales PL. 2011. Ciliopathies: An expanding disease spectrum. *Pediatr Nephrol* **26**: 1039–1056.
- Webb TR, Parfitt DA, Gardner JC, Martinez A, Bevilacqua D, Davidson AE, Zito I, Thiselton DL, Ressa JHC, Aperi M, et al. 2012. Deep intronic mutation in *OFD1*, identified by targeted genomic next-generation sequencing, causes a severe form of X-linked retinitis pigmentosa (RP23). *Hum Mol Genet* **21**: 3647–3654.
- Weil D, Blanchard S, Kaplan J, Guilford P, Gibson G, Walsh J, Mburu P, Varela A, Levilliers J, Weston M, et al. 1995. Defective myosin VIIA gene responsible for Usher syndrome type 1B. *Nature* **374**: 60–61.
- Weil D, El-Amraoui A, Masmoudi S, Mustapha M, Kikkawa Y, Laine S, Delmaghani S, Adato A, Nadifi S, Ben Zina Z, et al. 2003. Usher syndrome type I G (USH1G) is caused by mutations in the gene encoding SANS, a protein that associates with the USH1C protein, harmonin. *Hum Mol Genet* **12**: 463–471.
- Weitz CJ, Miyake Y, Shinzato K, Montag E, Zrenner E, Went LN, Nathans J. 1992a. Human tritanopia associated with two amino acid substitutions in the blue-sensitive opsin. *Am J Hum Genet* **50**: 498–507.
- Weitz CJ, Went LN, Nathans J. 1992b. Human tritanopia associated with a third amino acid substitution in the blue-sensitive visual pigment. *Am J Hum Genet* **51**: 444–446.
- Weleber RG. 2002. Infantile and childhood retinal blindness: A molecular perspective (The Franceschetti Lecture). *Ophthalmic Genet* **23**: 71–97.
- Weleber RG, Carr RE, Murphey WH, Sheffield VC, Stone EM. 1993. Phenotypic variation including retinitis pigmentosa, pattern dystrophy, and fundus flavimaculatus in a single family with a deletion of codon 153 or 154 of the peripherin/RDS gene. *Arch Ophthalmol* **111**: 1531–1542.
- Weston MD, Luijendijk MWJ, Humphrey KD, Möller C, Kimberling WJ. 2004. Mutations in the *VLGR1* gene implicate G-protein signaling in the pathogenesis of Usher syndrome type II. *Am J Hum Genet* **74**: 357–366.
- Williams DS. 2002. Transport to the photoreceptor outer segment by myosin VIIa and kinesin II. *Vision Res* **42**: 455–462.
- Williams D. 2008. Usher syndrome: Animal models, retinal function of Usher proteins, and prospects for gene therapy. *Vision Res* **48**: 433–441.
- Williams CL, McIntyre JC, Norris SR, Jenkins PM, Zhang L, Pei Q, Verhey K, Martens JR. 2014. Direct evidence for BBSome-associated intraflagellar transport reveals distinct properties of native mammalian cilia. *Nat Commun* **5**: 5813.
- Winderickx J, Sanocki E, Lindsey DT, Teller DY, Motulsky AG, Deeb SS. 1992. Defective colour vision associated with a missense mutation in the human green visual pigment gene. *Nat Genet* **1**: 251–256.
- Wright C, Healicorn R, English C, Burn J. 1994. Meckel syndrome: What are the minimum diagnostic criteria? *J Med Genet* **31**: 482–485.
- Xu M, Yang L, Wang F, Li HH, Wang X, Wang W, Ge Z, Wang K, Zhao L, Li HH, et al. 2015. Mutations in human *IFT140* cause non-syndromic retinal degeneration. *Hum Genet* **134**: 1069–1078.
- Yamamoto S, Sippel KC, Berson EL, Dryja TP. 1997. Defects in the rhodopsin kinase gene in the Oguchi form of stationary night blindness. *Nat Genet* **15**: 175–178.
- Yamashita T, Liu J, Gao J, LeNoue S, Wang C, Kaminoh J, Bowne SJ, Sullivan LS, Daiger SP, Zhang K, et al. 2009. Essential and synergistic roles of RP1 and RP1L1 in rod photoreceptor axoneme and retinitis pigmentosa. *J Neurosci* **29**: 9748–9760.
- Yan D, Liu XZ. 2010. Genetics and pathological mechanisms of Usher syndrome. *J Hum Genet* **55**: 327–335.
- Yang Z, Peachey NS, Moshfeghi DM, Thirumalaichary S, Chorich L, Shugart YY, Fan K, Zhang K. 2002. Mutations in the *RPGR* gene cause X-linked cone dystrophy. *Hum Mol Genet* **11**: 605–611.

K.M. Bujakowska et al.

- Yang J, Gao J, Adamian M, Wen XH, Pawlyk B, Zhang L, Sanderson MJ, Zuo J, Makino CL, Li T. 2005. The ciliary rootlet maintains long-term stability of sensory cilia. *Mol Cell Biol* **25**: 4129–4137.
- Yang Z, Chen Y, Lillo C, Chien J, Yu Z, Michaelides M, Klein M, Howes KA, Li Y, Kaminoh Y, et al. 2008. Mutant prominin 1 found in patients with macular degeneration disrupts photoreceptor disk morphogenesis in mice. *J Clin Invest* **118**: 2908–2916.
- Young RW. 1967. The renewal of photoreceptor cell outer segments. *J Cell Biol* **33**: 61–72.
- Zaki MS, Sattar S, Massoudi RA, Gleeson JG. 2011. Co-occurrence of distinct ciliopathy diseases in single families suggests genetic modifiers. *Am J Med Genet Part A* **155**: 3042–3049.
- Zallocchi M, Meehan DT, Delimont D, Askew C, Garige S, Gratton MA, Rothermund-Franklin CA, Cosgrove D. 2009. Localization and expression of clarin-1, the *Clrn1* gene product, in auditory hair cells and photoreceptors. *Hear Res* **255**: 109–120.
- Zhang Q, Li S, Xiao X, Jia X, Guo X. 2007. The 208delG mutation in *FSCN2* does not associate with retinal degeneration in Chinese individuals. *Invest Ophthalmol Vis Sci* **48**: 530–533.
- Zito I, Downes SM, Patel RJ, Cheetham ME, Ebenezer ND, Jenkins SA, Bhattacharya SS, Webster AR, Holder GE, Bird AC, et al. 2003. *RPGR* mutation associated with retinitis pigmentosa, impaired hearing, and sinorespiratory infections. *J Med Genet* **40**: 609–615.

Cold Spring Harbor Perspectives in Biology

Photoreceptor Cilia and Retinal Ciliopathies

Kinga M. Bujakowska, Qin Liu and Eric A. Pierce

Cold Spring Harb Perspect Biol 2017; doi: 10.1101/cshperspect.a028274 originally published online March 13, 2017

Subject Collection [Cilia](#)

Ciliary Mechanisms of Cyst Formation in Polycystic Kidney Disease

Ming Ma, Anna-Rachel Gallagher and Stefan Somlo

Photoreceptor Cilia and Retinal Ciliopathies

Kinga M. Bujakowska, Qin Liu and Eric A. Pierce

G-Protein-Coupled Receptor Signaling in Cilia

Kirk Mykytyn and Candice Askwith

Evolution of Cilia

David R. Mitchell

Transition Zone Migration: A Mechanism for Cytoplasmic Ciliogenesis and Postaxonemal Centriole Elongation

Tomer Avidor-Reiss, Andrew Ha and Marcus L. Basiri

Cilia and Obesity

Christian Vaisse, Jeremy F. Reiter and Nicolas F. Barbari

Posttranslational Modifications of Tubulin and Cilia

Dorota Wloga, Ewa Joachimiak, Panagiota Louka, et al.

Cilia in Left–Right Symmetry Breaking

Kyosuke Shinohara and Hiroshi Hamada

Discovery, Diagnosis, and Etiology of Craniofacial Ciliopathies

Elizabeth N. Schock and Samantha A. Brugmann

Axoneme Structure from Motile Cilia

Takashi Ishikawa

Cilia and Ciliopathies in Congenital Heart Disease

Nikolai T. Klena, Brian C. Gibbs and Cecilia W. Lo

Sperm Sensory Signaling

Dagmar Wachten, Jan F. Jikeli and U. Benjamin Kaupp

Primary Cilia and Coordination of Receptor Tyrosine Kinase (RTK) and Transforming Growth Factor β (TGF- β) Signaling

Søren T. Christensen, Stine K. Morthorst, Johanne B. Mogensen, et al.

Primary Cilia and Mammalian Hedgehog Signaling

Fiona Bangs and Kathryn V. Anderson

For additional articles in this collection, see <http://cshperspectives.cshlp.org/cgi/collection/>

**All Modifications and
Oligo Types Synthesized**
Long Oligos • Fluorescent • Chimeric • DNA • RNA • Antisense

Oligo Modifications?
Your wish is our command.

Radial Spokes—A Snapshot of the Motility Regulation, Assembly, and Evolution of Cilia and Flagella

Xiaoyan Zhu, Yi Liu and Pinfen Yang

Cilia and Mucociliary Clearance

Ximena M. Bustamante-Marin and Lawrence E. Ostrowski

For additional articles in this collection, see <http://cshperspectives.cshlp.org/cgi/collection/>

**All Modifications and
Oligo Types Synthesized**

Oligo Modifications?
Your wish is our command.

Long Oligos • Fluorescent • Chimeric • DNA • RNA • Antisense